

Reduction by symmetries in singular quantum-mechanical problems: general scheme and application to Aharonov-Bohm model

A.G. Smirnov^{a)}

*I. E. Tamm Theory Department, P. N. Lebedev Physical Institute, Leninsky prospect 53,
Moscow 119991, Russia*

We develop a general technique for finding self-adjoint extensions of a symmetric operator that respect a given set of its symmetries. Problems of this type naturally arise when considering two- and three-dimensional Schrödinger operators with singular potentials. The approach is based on constructing a unitary transformation diagonalizing the symmetries and reducing the initial operator to the direct integral of a suitable family of partial operators. We prove that symmetry preserving self-adjoint extensions of the initial operator are in a one-to-one correspondence with measurable families of self-adjoint extensions of partial operators obtained by reduction. The general scheme is applied to the three-dimensional Aharonov-Bohm Hamiltonian describing the electron in the magnetic field of an infinitely thin solenoid. We construct all self-adjoint extensions of this Hamiltonian, invariant under translations along the solenoid and rotations around it, and explicitly find their eigenfunction expansions.

I. INTRODUCTION

The spectral analysis of Schrödinger operators can be often facilitated if some symmetries of the problem are known. This happens if the spectral problem for the operators representing symmetries can be explicitly solved. In this case, the initial operator can be represented as a direct sum (or, more generally, direct integral) of partial operators corresponding to fixed eigenvalues of symmetries. In this way, two- and three-dimensional problems can be often reduced to solving certain one-dimensional Schrödinger equations (this is the case, for example, for spherically symmetric problems).

The present paper is concerned with such a reduction by symmetries in the case of singular quantum-mechanical problems. It is well known^{1,2} that strong singularities in the potential may lead to the lack of self-adjointness of the corresponding Schrödinger operator on its natural domain. As a result, the quantum model is no longer fixed uniquely by the potential and different quantum dynamics described by various self-adjoint extensions of the initial Schrödinger operator are possible. In this paper, we are interested in the extensions that respect some given set of symmetries of the initial operator. We propose a general technique for the reduction of such symmetry preserving extensions and apply it to the spectral analysis of the three-dimensional Aharonov-Bohm Hamiltonian describing the electron in the magnetic field of an infinitely thin solenoid.

In a general form, the problem can be posed as follows. Suppose H is a closed operator in a separable Hilbert space $\mathfrak{H}$ and $\mathfrak{X}$ is a subset of the algebra $L(\mathfrak{H})$ of linear bounded everywhere defined operators in $\mathfrak{H}$. We assume that $\mathfrak{X}$ is involutive (i.e., the adjoint T^* of every $T \in \mathfrak{X}$ belongs to $\mathfrak{X}$) and consists of pairwise commuting operators. Further, we assume $\mathfrak{X}$ is a set of symmetries of H , by which we mean that H commutes with elements of $\mathfrak{X}$ in the sense of the next definition.

Definition I.1. A linear operator H in $\mathfrak{H}$ with the domain D_H is said to commute with $T \in L(\mathfrak{H})$ if $T\Psi \in D_H$ and $HT\Psi = TH\Psi$ for any $\Psi \in D_H$.

Our aim is to show how spectral decompositions of symmetries can be used to represent closed (and, in particular, self-adjoint) symmetry preserving extensions of H as direct integrals of suitable partial operators. To clarify main ideas, we assume for a while that $\mathfrak{H}$ falls into an orthogonal direct sum of eigenspaces of symmetries. That is, we suppose that there is a countable family $\{\mathfrak{S}(s)\}_{s \in \mathcal{S}}$ of nontrivial closed pairwise orthogonal subspaces of $\mathfrak{H}$ such that

$$\mathfrak{H} = \bigoplus_{s \in \mathcal{S}} \mathfrak{S}(s) \quad (1)$$

and

$$T\Psi = g_T(s)\Psi, \quad \Psi \in \mathfrak{S}(s),$$

^{a)}Electronic mail: smirnov@lpi.ru

for every $s \in \mathcal{S}$ and $T \in \mathfrak{X}$, where $g_T(s)$ are some complex numbers (since $\mathfrak{S}(s)$ are nontrivial, the eigenvalues $g_T(s)$ are defined uniquely). Let P_s denote the orthogonal projection of $\mathfrak{H}$ onto $\mathfrak{S}(s)$. We would like to have a decomposition of the form

$$H = \bigoplus_{s \in \mathcal{S}} \mathcal{H}(s), \quad (2)$$

where $\mathcal{H}(s)$ are closed operators in $\mathfrak{S}(s)$ for all $s \in \mathcal{S}$ and $\bigoplus_{s \in \mathcal{S}} \mathcal{H}(s)$ is, by definition, the operator in $\mathfrak{H}$ whose graph consists of all $(\Psi, \tilde{\Psi}) \in \mathfrak{H} \oplus \mathfrak{H}$ such that $(P_s \Psi, P_s \tilde{\Psi})$ is in the graph of $\mathcal{H}(s)$ for every $s \in \mathcal{S}$. As the operator $\bigoplus_{s \in \mathcal{S}} \mathcal{H}(s)$ obviously commutes with all projections P_s , equality (2) can hold only if H commutes with P_s for every $s \in \mathcal{S}$. In fact, the latter condition is also sufficient for the existence of decomposition (2) (see Theorems 2 and 3 in Sec. 45 of Ref. 3). Given a decomposition of form (2), it can be easily shown that $\mathcal{H}(s)$ is actually the restriction of H to the domain $D_H \cap \mathfrak{S}(s)$ for every $s \in \mathcal{S}$. Thus, the partial operators $\mathcal{H}(s)$ are uniquely determined by H .

In general, the condition that $\mathfrak{S}(s)$ are eigenspaces of symmetries is insufficient to guarantee that H commutes with P_s for all $s \in \mathcal{S}$ (otherwise we could choose $\mathfrak{X}$ to be the one-element set containing the identity operator in $\mathfrak{H}$ and consider the decomposition $\mathfrak{H} = \mathfrak{H}' \oplus \mathfrak{H}'^\perp$, where $\mathfrak{H}'$ is an arbitrary closed subspace of $\mathfrak{H}$; it would follow that every closed operator in $\mathfrak{H}$ commutes with the orthogonal projection onto $\mathfrak{H}'$). The commutation between H and P_s can be ensured, however, if we require that

$$P_s \in \mathcal{A}(\mathfrak{X}) \quad (3)$$

for every $s \in \mathcal{S}$, where $\mathcal{A}(\mathfrak{X})$ is the smallest strongly closed subalgebra of $L(\mathfrak{H})$ containing the set $\mathfrak{X}$ and the identity operator in $\mathfrak{H}$ (in other words, $\mathcal{A}(\mathfrak{X})$ is the strong closure of the algebra generated by $\mathfrak{X}$ and the identity operator). Indeed, let $\mathcal{M}$ be the subset of $L(\mathfrak{H})$ consisting of all operators commuting with H . Since the sum and product of any two operators in $L(\mathfrak{H})$ commuting with H also commute with H , we conclude that $\mathcal{M}$ is an algebra. Moreover, it easily follows from the closedness of H that $\mathcal{M}$ is closed in the strong operator topology (see Lemma 1 in Ref. 4) and, hence, $\mathcal{A}(\mathfrak{X}) \subset \mathcal{M}$. Condition (3) therefore implies that $P_s \in \mathcal{M}$ or, in other words, that H commutes with P_s for all $s \in \mathcal{S}$.

We shall say that an eigenspace decomposition of form (1) is exact for $\mathfrak{X}$ if condition (3) is fulfilled for all $s \in \mathcal{S}$. Given an exact decomposition for $\mathfrak{X}$, it is easy to describe symmetry preserving closed extensions of H . Indeed, suppose (2) holds and a closed extension $\tilde{\mathcal{H}}(s)$ of $\mathcal{H}(s)$ is given for every $s \in \mathcal{S}$. Then the operator $\tilde{H} = \bigoplus_{s \in \mathcal{S}} \tilde{\mathcal{H}}(s)$ is clearly a closed extension of H commuting with all elements of $\mathfrak{X}$. Conversely, if $\tilde{H}$ is a closed extension of H commuting with all elements of $\mathfrak{X}$, then the above considerations applied to $\tilde{H}$ instead of H show that $\tilde{H}$ falls into the direct sum of its partial operators $\tilde{\mathcal{H}}(s)$, which are obviously closed extensions of $\mathcal{H}(s)$. Moreover, $\tilde{H}$ is densely defined if and only if all $\tilde{\mathcal{H}}(s)$ are densely defined, in which case we have $\tilde{H}^* = \bigoplus_{s \in \mathcal{S}} \tilde{\mathcal{H}}(s)^*$. Hence, $\tilde{H}$ is self-adjoint if and only if $\tilde{\mathcal{H}}(s)$ is self-adjoint for every $s \in \mathcal{S}$.

Thus, given an exact decomposition for $\mathfrak{X}$, the operator H falls into the direct sum of its partial operators $\mathcal{H}(s)$, and the symmetry preserving closed (resp., self-adjoint) extensions of H are precisely the direct sums of closed (resp., self-adjoint) extensions of $\mathcal{H}(s)$.

Exactness condition (3) can be given another, equivalent, formulation that is better suited for verifying in concrete applications. More specifically, we claim that (3) holds for all $s \in \mathcal{S}$ if and only if the following condition is fulfilled:

(E) For every $s, s' \in \mathcal{S}$ such that $s \neq s'$, there is $T \in \mathfrak{X}$ such that $g_T(s) \neq g_T(s')$.

Indeed, suppose (E) holds. Since $\mathfrak{X}$ is involutive, the elements of $\mathcal{A}(\mathfrak{X})$ can be characterized using von Neumann's bicommutant theorem (see Sec. III for details): an operator in $L(\mathfrak{H})$ belongs to $\mathcal{A}(\mathfrak{X})$ if and only if it commutes with every element of commutant of $\mathfrak{X}$, i.e., with every operator in $L(\mathfrak{H})$ that commutes with all elements of $\mathfrak{X}$. To prove (3), we have to verify this condition for P_s . Let $R \in L(\mathfrak{H})$ commute with all elements of $\mathfrak{X}$. Since $TP_s = P_s T = g_T(s)P_s$ for every $s \in \mathcal{S}$ and $T \in \mathfrak{X}$, we have $P_{s'} R T P_s = g_T(s) P_{s'} R P_s = g_T(s') P_{s'} R P_s$ for all $T \in \mathfrak{X}$ and $s, s' \in \mathcal{S}$. In view of (E), this implies that $P_{s'} R P_s = 0$ for all $s, s' \in \mathcal{S}$ such that $s \neq s'$. As $\Psi = \sum_{s \in \mathcal{S}} P_s \Psi$ for any $\Psi \in \mathfrak{H}$, we obtain

$$P_s R \Psi = \sum_{s' \in \mathcal{S}} P_s R P_{s'} \Psi = P_s R P_s \Psi = \sum_{s' \in \mathcal{S}} P_{s'} R P_s \Psi = R P_s \Psi$$

for every $s \in \mathcal{S}$ and $\Psi \in \mathfrak{H}$. Thus, P_s commutes with R for every $s \in \mathcal{S}$, and (3) is proved. We now show that, conversely, (3) implies (E). Suppose, to the contrary, that (3) holds and there exist $s_1, s_2 \in \mathcal{S}$ such that $s_1 \neq s_2$ and $g_T(s_1) = g_T(s_2)$ for every $T \in \mathfrak{X}$. We then have $\tilde{P} T = T \tilde{P} = g_T(s_1) \tilde{P}$ for every $T \in \mathfrak{X}$, where $\tilde{P} = P_{s_1} + P_{s_2}$ is the orthogonal projection of $\mathfrak{H}$ onto $\mathfrak{S}(s_1) \oplus \mathfrak{S}(s_2)$. It follows that every element

of $\mathfrak{X}$ commutes with $\tilde{P}R\tilde{P}$ for every $R \in L(\mathfrak{H})$. In view of (3) and the characterization of elements of $\mathcal{A}(\mathfrak{X})$ given above, this implies that P_{s_1} commutes with $\tilde{P}R\tilde{P}$ for every $R \in L(\mathfrak{H})$, which is obviously false. We thus arrive at a contradiction and our claim is proved.

The above discussion is based on the assumption that the Hilbert space $\mathfrak{H}$ falls into a direct sum of eigenspaces of symmetries. In general, this assumption does not hold and one has to use direct integral decompositions instead. More specifically, we can try to find a positive measure ν , a ν -measurable family $\mathfrak{S}$ of Hilbert spaces, and a unitary operator $V: \mathfrak{H} \rightarrow \int^\oplus \mathfrak{S}(s) d\nu(s)$ such that every $T \in \mathcal{X}$ is representable in the form $V^{-1}\mathcal{T}_g^{\nu, \mathfrak{S}}V$ for some ν -measurable complex function g , where $\mathcal{T}_g^{\nu, \mathfrak{S}}$ is the operator of multiplication by g in $\int^\oplus \mathfrak{S}(s) d\nu(s)$ (we refer the reader to Appendix B for the notions related to direct integrals of Hilbert spaces). A triple $(\nu, \mathfrak{S}, V)$ satisfying this condition is said to be a diagonalization for $\mathfrak{X}$. Since countable direct sums of separable Hilbert spaces can be always identified with suitable direct integrals (see Sec. B3 and formula (B2)), the eigenspace decompositions considered above can be viewed as a particular case of diagonalizations, with the role of ν and V played by the counting measure⁵ on $\mathcal{S}$ and the natural identification between $\mathfrak{H}$ and $\bigoplus_{s \in \mathcal{S}} \mathfrak{S}(s)$ respectively.

Given a diagonalization $(\nu, \mathfrak{S}, V)$, we would like to have a decomposition of H of the form

$$H = V^{-1} \int^\oplus \mathcal{H}(s) d\nu(s) V, \quad (4)$$

where $\mathcal{H}$ is a ν -measurable family of closed operators in $\mathfrak{S}$. This decomposition is an analogue of (2) in the direct integral setting. As we have seen, such a decomposition, in general, does not exist even in the direct sum case. It turns out, however, that this problem does not arise if we confine ourselves to *exact* diagonalizations that are singled out by the requirement that

$$V^{-1}\mathcal{T}_g^{\nu, \mathfrak{S}}V \in \mathcal{A}(\mathfrak{X}) \quad (5)$$

for every ν -measurable ν -essentially bounded function g . It will be shown that, in contrast to direct sum decompositions, exact diagonalizations exist for every involutive set $\mathfrak{X} \subset L(\mathfrak{H})$ of pairwise commuting operators. Exactness condition (5) (which is an analogue of (3)) implies that VHV^{-1} commutes with $\mathcal{T}_g^{\nu, \mathfrak{S}}$ for every ν -measurable ν -essentially bounded g since, as shown above, H commutes with all elements of $\mathcal{A}(\mathfrak{X})$. This allows us to apply the von Neumann's reduction theory⁶ (or, more precisely, its generalization^{7,8} for the case of unbounded operators) to VHV^{-1} and obtain decomposition (4) as a result. Moreover, given an exact diagonalization $(\nu, \mathfrak{S}, V)$ for $\mathfrak{X}$ and a decomposition of form (4), the symmetry preserving closed (resp., self-adjoint) extensions of H are precisely the operators of the form

$$V^{-1} \int^\oplus \tilde{\mathcal{H}}(s) d\nu(s) V,$$

where $\tilde{\mathcal{H}}$ is a ν -measurable family of operators in $\mathfrak{S}$ such that $\tilde{\mathcal{H}}(s)$ is a closed (resp., self-adjoint) extension of the partial operator $\mathcal{H}(s)$ for ν -a.e. s .⁹ Condition (5) is, as a rule, inconvenient for concrete applications because it involves the algebra $\mathcal{A}(\mathfrak{X})$ rather than the set $\mathfrak{X}$ itself. We shall see, however, that a simple exactness criterion similar to condition (E) can be obtained under very mild restrictions on the measure ν (Theorem IV.2). This generalized condition (E) can usually be easily verified for concrete examples.

The results formulated above suggest that, in general, one can find the symmetry preserving self-adjoint extensions of H by doing the following steps:

- (I) Find an exact diagonalization $(\nu, \mathfrak{S}, V)$ for $\mathfrak{X}$.
- (II) Compute partial operators $\mathcal{H}(s)$ satisfying (4).
- (III) Find self-adjoint extensions of the partial operators $\mathcal{H}(s)$.

In practice, the operator H often comes as the closure of some non-closed operator $\check{H}$. While $\check{H}$ is usually given by some explicit formula, finding an explicit description of H may be a difficult task. We shall see, however, that every symmetry of $\check{H}$ is also a symmetry of H (Lemma III.1). Moreover, finding partial operators of H can be effectively reduced to some computations involving $\check{H}$ (Proposition V.3). For this reason, the knowledge of $\check{H}$ is actually sufficient for doing steps (I)-(III) for H .

We apply the general construction described above to the three-dimensional model of an electron in the magnetic field of an infinitely thin solenoid. The Hamiltonian for this model is formally given by the differential expression

$$\sum_{j=1}^3 (-i\partial_{x_j} + \mathcal{A}_j(x))^2, \quad (6)$$

where $x = (x_1, x_2, x_3) \in \mathbb{R}^3$ and the vector potential $\mathcal{A} = (\mathcal{A}_1, \mathcal{A}_2, \mathcal{A}_3)$ has the form

$$\mathcal{A}_1(x) = -\frac{\phi x_2}{x_1^2 + x_2^2}, \quad \mathcal{A}_2(x) = \frac{\phi x_1}{x_1^2 + x_2^2}, \quad \mathcal{A}_3(x) = 0.$$

Here, the real parameter ϕ is equal to the flux of the magnetic field through the solenoid divided by 2π .

This model was originally considered by Aharonov and Bohm¹⁰. Self-adjoint extensions in the two-dimensional variant of this model, as well as their eigenfunction expansions and corresponding scattering amplitudes, were analysed in Refs. 11–13.

The vector potential $\mathcal{A}$ is smooth on the domain $\mathcal{O} = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1^2 + x_2^2 > 0\}$, which is obtained by excluding the x_3 -axis from $\mathbb{R}^3$. Hence, (6) naturally determines an operator $\mathcal{H}^\phi$ in the space $C^\infty(\mathcal{O})$ of smooth functions on $\mathcal{O}$,

$$\begin{aligned} (\mathcal{H}^\phi \Phi)(x) &= \sum_{j=1}^3 (-i\partial_{x_j} + \mathcal{A}_j(x))^2 \Phi(x) = \\ &= \left(-\Delta + \frac{2i\phi}{x_1^2 + x_2^2} (x_2 \partial_{x_1} - x_1 \partial_{x_2}) + \frac{\phi^2}{x_1^2 + x_2^2} \right) \Phi(x), \quad \Phi \in C^\infty(\mathcal{O}). \end{aligned} \quad (7)$$

Restricting $\mathcal{H}^\phi$ to the space $C_0^\infty(\mathcal{O})$ of smooth functions on $\mathcal{O}$ with compact support and passing to Λ -equivalence classes, where Λ is the Lebesgue measure on $\mathbb{R}^3$, we obtain a (non-closed) densely defined symmetric operator $\tilde{H}^\phi$ in $L_2(\mathbb{R}^3)$:

$$\begin{aligned} D_{\tilde{H}^\phi} &= \{[\Phi]_\Lambda : \Phi \in C_0^\infty(\mathcal{O})\}, \\ \tilde{H}^\phi[\Phi]_\Lambda &= [\mathcal{H}^\phi \Phi]_\Lambda, \quad \Phi \in C_0^\infty(\mathcal{O}), \end{aligned}$$

where $[\Phi]_\Lambda$ denotes the Λ -equivalence class corresponding to Φ . We define the operator H^ϕ in $L_2(\mathbb{R}^3)$ as the closure of $\tilde{H}^\phi$,

$$H^\phi = \overline{\tilde{H}^\phi}. \quad (8)$$

Let $\mathcal{G}$ be the Abelian group of linear operators in $\mathbb{R}^3$ generated by translations along the x_3 -axis and rotations around the x_3 -axis. Given $G \in \mathcal{G}$, we denote by T_G the unitary operator in $L_2(\mathbb{R}^3)$ taking Ψ to $[\Psi \circ G^{-1}]_\Lambda$ for any $\Psi \in L_2(\mathbb{R}^3)$. Clearly, $G \rightarrow T_G$ is a linear representation of $\mathcal{G}$ in $L_2(\mathbb{R}^3)$. It is straightforward to check that $\tilde{H}^\phi$ (and, hence, H^ϕ) commutes with T_G for any $G \in \mathcal{G}$. The above abstract scheme can therefore be applied to $H = H^\phi$ and the set $\mathfrak{X}$ consisting of all T_G with $G \in \mathcal{G}$. As a result, we shall describe all self-adjoint extensions of H^ϕ commuting with T_G for any $G \in \mathcal{G}$ and find eigenfunction expansions for such extensions.

The paper is organized as follows. In Sec. II, we formulate the results concerning self-adjoint extensions and eigenfunction expansions for the Aharonov-Bohm model. In Secs. III–V, we elaborate on the general scheme of constructing symmetry preserving extensions outlined above. Sec. III is devoted to preliminaries concerning the commutation properties of operators in Hilbert space. In Sec. IV, we establish the existence of exact diagonalizations and prove an exactness criterion, analogous to condition (E) discussed above. In Sec. V, we consider direct integral decompositions for closed operators possessing a given set of symmetries and describe their symmetry preserving extensions. The aim of Secs. VI–X is to derive eigenfunction expansions for the Aharonov-Bohm model. Sec. VI is concerned with self-adjoint extensions and eigenfunction expansions for one-dimensional Schrödinger operators. In Sec. VII, we treat the measurability questions for families of such operators. Secs. VI and VII provide a background for performing steps (I)–(III) for H^ϕ in Sec VIII. As a result, we represent symmetry preserving self-adjoint extensions of H^ϕ as direct integrals of suitable one-dimensional Schrödinger operators. In Sec IX, we study a special construction, called the direct integral of measures, which can be viewed as a generalization of direct products of measures and arises naturally in the spectral analysis of direct integrals of operators. In Sec. X, we combine the results of Secs. VI, VIII, and IX to obtain the eigenfunction expansions of symmetry preserving self-adjoint extensions of H^ϕ . In Appendices A and B, we give the necessary background material concerning measure theory and direct integral decompositions of operators in Hilbert space respectively.

II. FORMULATION OF RESULTS FOR THE AHARONOV-BOHM MODEL

In this section, we give a precise description of self-adjoint extensions of H^ϕ commuting with T_G for all $G \in \mathcal{G}$ and their eigenfunction expansions. The proofs of results stated below will be given in Sec. X.

Let $S = \mathbb{Z} \times \mathbb{R}$, where $\mathbb{Z}$ is the set of integers. For $\phi \in \mathbb{R}$, let the subset A^ϕ of S be given by

$$A^\phi = \{(m, p) \in S : |m + \phi| < 1\}. \quad (9)$$

For every Borel¹⁴ real function θ on A^ϕ , we shall define a map $\mathcal{W}_\theta^\phi$ on $S \times \mathbb{R}$ such that

- (a) For every $s \in S$ and $E \in \mathbb{R}$, $\mathcal{W}_\theta^\phi(s, E)$ is a smooth complex function on $\mathcal{O}$ that is locally square-integrable on $\mathbb{R}^3$.
- (b) The equalities¹⁵

$$\begin{aligned} \mathcal{H}^\phi \mathcal{W}_\theta^\phi(s, E) &= (E + p^2) \mathcal{W}_\theta^\phi(s, E), \\ -i \partial_{x_3} \mathcal{W}_\theta^\phi(s, E|x) &= p \mathcal{W}_\theta^\phi(s, E|x), \\ -i(x_1 \partial_{x_2} - x_2 \partial_{x_1}) \mathcal{W}_\theta^\phi(s, E|x) &= m \mathcal{W}_\theta^\phi(s, E|x) \end{aligned}$$

hold for every $s = (m, p) \in S$, $E \in \mathbb{R}$, and $x = (x_1, x_2, x_3) \in \mathcal{O}$.

Furthermore, we shall define a positive Borel measure M_θ^ϕ on $S \times \mathbb{R}$ such that the following statement is valid.

Proposition II.1. *Let $\phi \in \mathbb{R}$ and θ be a Borel real function on A^ϕ . Then there is a unique unitary operator $W: L_2(\mathbb{R}^3) \rightarrow L_2(S \times \mathbb{R}, M_\theta^\phi)$ such that*

$$(W\Psi)(s, E) = \int \overline{\mathcal{W}_\theta^\phi(s, E|x)} \Psi(x) dx, \quad \Psi \in L_2^c(\mathbb{R}^3), \quad (10)$$

for M_θ^ϕ -a.e. (s, E) , where the bar means complex conjugation and $L_2^c(\mathbb{R}^3)$ is the subspace of $L_2(\mathbb{R}^3)$ consisting of all its elements vanishing Λ -a.e. outside some compact subset of $\mathbb{R}^3$.

Condition (b) means that $\mathcal{W}_\theta^\phi(m, s)$ is a joint generalized eigenfunction of operator (6) and x_3 -components of momentum and angular momentum for every $s \in S$ and $E \in \mathbb{R}$. Proposition II.1 states that these eigenfunctions constitute a complete set and can be used to expand square-integrable functions on $\mathbb{R}^3$. We shall see (Theorem II.5) that all such eigenfunction expansions correspond to symmetry preserving self-adjoint extensions of H^ϕ , but first we give precise definitions of $\mathcal{W}_\theta^\phi$ and M_θ^ϕ .

It follows from the second and third equalities in condition (b) that

$$\mathcal{W}_\theta^\phi(s, E|x) = \frac{e^{ipx_3}}{2\pi\sqrt{r_x}} \left(\frac{x_1 + ix_2}{r_x} \right)^m \mathcal{J}_\theta^\phi(s, E|r_x) \quad (11)$$

for every $s = (m, p) \in S$, $E \in \mathbb{R}$, and $x \in \mathcal{O}$, where $r_x = \sqrt{x_1^2 + x_2^2}$ and $\mathcal{J}_\theta^\phi(s, E)$ is some smooth function on $\mathbb{R}_+ = (0, \infty)$ (note that $x_1 \partial_{x_2} - x_2 \partial_{x_1}$ and $r_x^{-m}(x_1 + ix_2)^m$ correspond to ∂_φ and $e^{im\varphi}$ respectively in the polar coordinates). For future convenience, we write the factor $(2\pi\sqrt{r_x})^{-1}$ separately rather than subsume it into $\mathcal{J}_\theta^\phi(s, E)$. The first equality in condition (b) is fulfilled if and only if $\mathcal{J}_\theta^\phi(s, E)$ satisfies the one-dimensional Schrödinger equation

$$-\partial_r^2 \mathcal{J}_\theta^\phi(s, E|r) + \frac{(m + \phi)^2 - 1/4}{r^2} \mathcal{J}_\theta^\phi(s, E|r) = E \mathcal{J}_\theta^\phi(s, E|r), \quad r \in \mathbb{R}_+, \quad (12)$$

for every $s = (m, p) \in S$ and $E \in \mathbb{R}$.

To define $\mathcal{J}_\theta^\phi$ explicitly, we introduce some additional notation. For any $E, \kappa \in \mathbb{R}$, we define the function $u^\kappa(E)$ on $\mathbb{R}_+$ by the relation

$$u^\kappa(E|r) = r^{1/2+\kappa} \mathcal{X}_\kappa(r^2 E), \quad r \in \mathbb{R}_+, \quad (13)$$

where the entire function $\mathcal{X}_\kappa$ is given by

$$\mathcal{X}_\kappa(\zeta) = \frac{1}{2^\kappa} \sum_{n=0}^{\infty} \frac{(-1)^n \zeta^n}{\Gamma(\kappa + n + 1) n! 2^{2n}}, \quad \zeta \in \mathbb{C}. \quad (14)$$

For $-1 < \kappa < 1$, $\vartheta \in \mathbb{R}$, and $E \in \mathbb{R}$, we define the function $u_\vartheta^\kappa(E)$ on $\mathbb{R}_+$ by setting

$$u_\vartheta^\kappa(E) = \frac{u^\kappa(E) \sin(\vartheta + \vartheta_\kappa) - u^{-\kappa}(E) \sin(\vartheta - \vartheta_\kappa)}{\sin \pi \kappa}, \quad 0 < |\kappa| < 1, \quad (15)$$

and

$$u_\vartheta^0(E|r) = \lim_{\kappa \rightarrow 0} u_\vartheta^\kappa(E|r) = u^0(E|r) \cos \vartheta + \frac{2}{\pi} \left[\left(\ln \frac{r}{2} + \gamma \right) u^0(E|r) - \sqrt{r} \mathcal{Y}(r^2 E) \right] \sin \vartheta, \quad r \in \mathbb{R}_+, \quad (16)$$

where

$$\vartheta_\kappa = \frac{\pi \kappa}{2}, \quad (17)$$

the entire function $\mathcal{Y}$ is given by

$$\mathcal{Y}(\zeta) = \sum_{n=1}^{\infty} \frac{(-1)^n c_n}{(n!)^2 2^{2n}} \zeta^n, \quad c_n = \sum_{j=1}^n \frac{1}{j}, \quad (18)$$

and $\gamma = \lim_{n \rightarrow \infty} (c_n - \ln n) = 0,577 \dots$ is the Euler constant.¹⁶ We now define $\mathcal{J}_\theta^\phi$ by setting

$$\mathcal{J}_\theta^\phi(s, E) = \begin{cases} u^{|m+\phi|}(E), & s \in S \setminus A^\phi, \\ u_{\theta(s)}^{m+\phi}(E), & s \in A^\phi, \end{cases} \quad (19)$$

for every $s = (m, p) \in S$ and $E \in \mathbb{R}$.

The function $\mathcal{X}_\kappa$ is closely related to Bessel functions: for $\zeta \neq 0$, we have

$$\mathcal{X}_\kappa(\zeta) = \zeta^{-\kappa/2} J_\kappa(\zeta^{1/2}), \quad (20)$$

where J_κ is the Bessel function of the first kind of order κ . Since J_κ satisfies the Bessel equation, it follows that

$$-\partial_r^2 u^{\pm\kappa}(E|r) + \frac{\kappa^2 - 1/4}{r^2} u^{\pm\kappa}(E|r) = E u^{\pm\kappa}(E|r), \quad r \in \mathbb{R}_+, \quad (21)$$

for every $E \neq 0$. By continuity, this is true for $E = 0$ as well. Moreover, (21) is satisfied for $u_\vartheta^\kappa(E)$ in place of $u^{\pm\kappa}(E)$ for every $-1 < \kappa < 1$, $E \in \mathbb{R}$, and $\vartheta \in \mathbb{R}$ (this obviously follows from (15) for $0 < |\kappa| < 1$; for $\kappa = 0$, we can express $u_\vartheta^0(E|r)$ in terms of the Bessel functions J_0 and Y_0 by means of the equality¹⁷ $\pi Y_0(\zeta) = 2(\gamma + \ln(\zeta/2)) J_0(\zeta) - 2\mathcal{Y}(\zeta^2)$ and make use of the Bessel equation). In view of (19), we conclude that (12) holds for every $s \in S$ and $E \in \mathbb{R}$. It follows immediately from (11), (13), (15), (16), and (19) that $\mathcal{W}_\theta^\phi(s, E)$ is locally square-integrable on $\mathbb{R}^3$ for every $s \in S$ and $E \in \mathbb{R}$. Thus, both conditions (a) and (b) are fulfilled for the map $\mathcal{W}_\theta^\phi$ defined by (11) and (19).

It should be noted that equation (12) and condition (a) determine $\mathcal{J}_\theta^\phi(s, E)$ uniquely up to a numerical factor for $s \in S \setminus A^\phi$. At the same time, if $s \in A^\phi$, then there are two linearly independent solutions of (12) compatible with (a) for every $E \in \mathbb{R}$. According to (19), different choices of the function θ on A^ϕ allow us to pick different solutions of (12) and, consequently, lead to different eigenfunction expansions for operator (6).

We now proceed to define the measure M_θ^ϕ . The above definition of $\mathcal{W}_\theta^\phi$ involves the generalized eigenfunctions $u^\kappa(E)$ and $u_\vartheta^\kappa(E)$ of the one-dimensional Schrödinger operator

$$-\partial_r^2 + \frac{\kappa^2 - 1/4}{r^2}. \quad (22)$$

Similarly, M_θ^ϕ will be constructed from one-dimensional measures $\mathcal{V}_\kappa$ and $\mathcal{V}_{\kappa, \vartheta}$ entering eigenfunction expansions associated with differential expression (22) (see Propositions VI.3 and VI.4). For $\kappa \in \mathbb{R}$, the positive Borel measure $\mathcal{V}_\kappa$ on $\mathbb{R}$ is given by

$$d\mathcal{V}_\kappa(E) = \frac{1}{2} \Theta(E) E^{|\kappa|} dE, \quad (23)$$

where Θ is the Heaviside function, i.e., $\Theta(E) = 1$ for $E \geq 0$ and $\Theta(E) = 0$ for $E < 0$. Given $-1 < \kappa < 1$ and $\vartheta \in \mathbb{R}$, we define the positive Borel measure $\mathcal{V}_{\kappa, \vartheta}$ on $\mathbb{R}$ as follows. If $0 < |\kappa| < 1$, we set

$$\mathcal{V}_{\kappa, \vartheta} = \begin{cases} \tilde{\mathcal{V}}_{\kappa, \vartheta}, & \vartheta \in [-|\vartheta_\kappa|, |\vartheta_\kappa|] + \pi\mathbb{Z}, \\ \frac{\pi \sin \pi \kappa |E_{\kappa, \vartheta}|}{2\kappa \sin(\vartheta + \vartheta_\kappa) \sin(\vartheta - \vartheta_\kappa)} \delta_{E_{\kappa, \vartheta}} + \tilde{\mathcal{V}}_{\kappa, \vartheta}, & \vartheta \in (|\vartheta_\kappa|, \pi - |\vartheta_\kappa|) + \pi\mathbb{Z}, \end{cases} \quad (24)$$

where ϑ_κ is defined by (17), the positive Borel measure $\tilde{\mathcal{V}}_{\kappa,\vartheta}$ on $\mathbb{R}$ is given by

$$d\tilde{\mathcal{V}}_{\kappa,\vartheta}(E) = \frac{1}{2} \frac{\Theta(E) \sin^2 \pi \kappa}{E^{-\kappa} \sin^2(\vartheta + \vartheta_\kappa) - 2 \cos \pi \kappa \sin(\vartheta + \vartheta_\kappa) \sin(\vartheta - \vartheta_\kappa) + E^\kappa \sin^2(\vartheta - \vartheta_\kappa)} dE \quad (25)$$

and $\delta_{E_{\kappa,\vartheta}}$ is the Dirac measure at the point

$$E_{\kappa,\vartheta} = - \left(\frac{\sin(\vartheta + \vartheta_\kappa)}{\sin(\vartheta - \vartheta_\kappa)} \right)^{1/\kappa}. \quad (26)$$

For $\kappa = 0$, the measure $\mathcal{V}_{\kappa,\vartheta}$ is defined by taking the limit $\kappa \rightarrow 0$ in formulas (24), (25), and (26). This yields

$$\mathcal{V}_{0,\vartheta} = \begin{cases} \tilde{\mathcal{V}}_{0,\vartheta}, & \vartheta \in \pi\mathbb{Z}, \\ \frac{\pi^2 |E_{0,\vartheta}|}{2 \sin^2 \vartheta} \delta_{E_{0,\vartheta}} + \tilde{\mathcal{V}}_{0,\vartheta}, & \vartheta \notin \pi\mathbb{Z}, \end{cases} \quad (27)$$

where

$$E_{0,\vartheta} = -e^{\pi \cot \vartheta} \quad (28)$$

and the positive Borel measure $\tilde{\mathcal{V}}_{0,\vartheta}$ on $\mathbb{R}$ is given by

$$d\tilde{\mathcal{V}}_{0,\vartheta}(E) = \frac{1}{2} \frac{\Theta(E)}{(\cos \vartheta - \ln E \sin \vartheta / \pi)^2 + \sin^2 \vartheta} dE. \quad (29)$$

The next lemma shows, in particular, that the measure $\mathcal{V}_{\kappa,\vartheta}$ depends continuously on both κ and ϑ .

Lemma II.2. *Let F be a bounded (continuous) Borel function on $\mathbb{R}$ with compact support. Then $(\kappa, \vartheta) \rightarrow \int F(E) d\mathcal{V}_{\kappa,\vartheta}(E)$ is a Borel (resp., continuous) function on $(-1, 1) \times \mathbb{R}$ that is bounded on $[-\alpha, \alpha] \times \mathbb{R}$ for every $0 \leq \alpha < 1$.*

Proof. See Theorem 4 in Ref. 18. □

Given a Borel real function θ on A^ϕ , let μ_θ^ϕ be the measure-valued map on S such that

$$\mu_\theta^\phi(s) = \begin{cases} \mathcal{V}_{|m+\phi|}, & s \in S \setminus A^\phi, \\ \mathcal{V}_{m+\phi, \theta(s)}, & s \in A^\phi, \end{cases} \quad (30)$$

for every $s = (m, p) \in S$.

The next statement follows immediately from Lemma II.2.

Lemma II.3. *Let $\phi \in \mathbb{R}$ and θ be a Borel real function on A^ϕ . For any compact set $K \subset \mathbb{R}$, $s \rightarrow \mu_\theta^\phi(s|K)$ is a Borel function on S that is bounded on every compact subset of S .*

Let ϱ be the counting measure on $\mathbb{Z}$. We define the positive Borel measure ν_0 on $S = \mathbb{Z} \times \mathbb{R}$ by setting $\nu_0 = \varrho \times \lambda$, where λ is the Lebesgue measure on $\mathbb{R}$. For any ν_0 -integrable f , the function $p \rightarrow f(m, p)$ is integrable for every $m \in \mathbb{Z}$ and we have

$$\int_S f(m, p) d\nu_0(m, p) = \sum_{m \in \mathbb{Z}} \int_{-\infty}^{\infty} f(m, p) dp.$$

Proposition II.4. *Let $\phi \in \mathbb{R}$ and θ be a Borel real function on A^ϕ . Then there is a unique positive Borel measure M on $S \times \mathbb{R}$ such that*

$$M(K' \times K) = \int_{K'} \mu_\theta^\phi(s|K) d\nu_0(s) \quad (31)$$

for every compact sets $K' \subset S$ and $K \subset \mathbb{R}$. If f is an M -integrable complex function, then the function $E \rightarrow f(s, E)$ is $\mu_\theta^\phi(s)$ -integrable for ν_0 -a.e. s , the function $s \rightarrow \int f(s, E) d\mu_\theta^\phi(s|E)$ is ν_0 -integrable, and

$$\int f(s, E) dM(s, E) = \int d\nu_0(s) \int f(s, E) d\mu_\theta^\phi(s|E). \quad (32)$$

Note that the right-hand side of (31) is well-defined in view of Lemma II.3. Given $\phi \in \mathbb{R}$ and a Borel real function θ on A^ϕ , we let M_θ^ϕ denote the measure M satisfying the conditions of Proposition II.4. We thus have explicitly defined all entities entering Proposition II.1.

It is worth mentioning that the structure of the set A^ϕ depends on whether $\phi \in \mathbb{Z}$ or not. If $\phi \in \mathbb{Z}$, then we have $A^\phi = \{-\phi\} \times \mathbb{R}$ and defining a Borel function on A^ϕ amounts to defining a single Borel function on $\mathbb{R}$. On the other hand, if $\phi \notin \mathbb{Z}$, then $A^\phi = \{m_\phi, m_\phi + 1\} \times \mathbb{R}$, where $m_\phi \in \mathbb{Z}$ is such that $-\phi - 1 < m_\phi < -\phi$, and every Borel function on A^ϕ is determined by two Borel functions on $\mathbb{R}$. Using functions on A^ϕ instead of functions on $\mathbb{R}$ allows us to treat both cases uniformly.

Given $\phi \in \mathbb{R}$ and a Borel real function θ on A^ϕ , the operator W satisfying the conditions of Proposition II.1 will be denoted by W_θ^ϕ . Let the function $\mathfrak{f}$ on $S \times \mathbb{R}$ be defined by the relation

$$\mathfrak{f}(m, p; E) = p^2 + E. \quad (33)$$

We define the self-adjoint operator H_θ^ϕ in $L_2(\mathbb{R}^3)$ by setting

$$H_\theta^\phi = (W_\theta^\phi)^{-1} \mathcal{T}_\mathfrak{f}^M W_\theta^\phi, \quad (34)$$

where $M = M_\theta^\phi$ and $\mathcal{T}_\mathfrak{f}^M$ is the operator of multiplication by $\mathfrak{f}$ in $L_2(S \times \mathbb{R}, M)$ (see Sec. A 7).

Let $\Phi \in C_0^\infty(\mathcal{O})$, $\Psi = [\Phi]_\Lambda$, and $h_\Psi(s, E)$ be the right-hand side of (10). In view of (7) and the first equality in condition (b), integrating by parts yields

$$h_{\tilde{H}^\phi \Psi}(s, E) = \int \overline{\mathcal{W}_\theta^\phi(s, E|x)} (\mathcal{H}^\phi \Phi)(x) dx = \int (\mathcal{H}^\phi \mathcal{W}_\theta^\phi(s, E))(x) \Psi(x) dx = \mathfrak{f}(s, E) h_\Psi(s, E)$$

for every $s \in S$ and $E \in \mathbb{R}$. It follows that $W_\theta^\phi \tilde{H}^\phi \Psi = \mathcal{T}_\mathfrak{f}^M W_\theta^\phi \Psi$ for every $\Psi \in D_{\tilde{H}^\phi}$, i.e., H_θ^ϕ is a self-adjoint extension of $\tilde{H}^\phi$ and, hence, of H^ϕ .

Further, given $\alpha, \beta \in \mathbb{R}$, let $G_{\alpha\beta} \in \mathcal{G}$ be defined by the relation

$$G_{\alpha\beta} x = (x_1 \cos \alpha - x_2 \sin \alpha, x_1 \sin \alpha + x_2 \cos \alpha, x_3 + \beta), \quad (35)$$

where $x = (x_1, x_2, x_3) \in \mathbb{R}^3$. It follows from (11) that

$$\mathcal{W}_\theta^\phi(s, E|G_{\alpha\beta} x) = e^{im\alpha + ip\beta} \mathcal{W}_\theta^\phi(s, E|x), \quad x \in \mathcal{O},$$

for every $s = (m, p) \in S$ and $E \in \mathbb{R}$. Let $\Psi \in L_2^c(\mathbb{R}^3)$. Setting $\tilde{\Psi} = T_{G_{\alpha\beta}} \Psi$, we obtain

$$h_{\tilde{\Psi}}(s, E) = \int \overline{\mathcal{W}_\theta^\phi(s, E|x)} \Psi(G_{\alpha\beta}^{-1} x) dx = g_{\alpha\beta}(s, E) h_\Psi(s, E)$$

for every $s \in S$ and $E \in \mathbb{R}$, where $g_{\alpha\beta}$ is the function $(m, p; E) \rightarrow e^{-im\alpha - ip\beta}$ on $S \times \mathbb{R}$. As $L_2^c(\mathbb{R}^3)$ is dense in $L_2(\mathbb{R}^3)$, this implies that

$$T_{G_{\alpha\beta}} = (W_\theta^\phi)^{-1} \mathcal{T}_{g_{\alpha\beta}}^M W_\theta^\phi.$$

Since every element of $\mathcal{G}$ is equal to $G_{\alpha\beta}$ for some $\alpha, \beta \in \mathbb{R}$ and $\mathcal{T}_\mathfrak{f}^M$ commutes with $\mathcal{T}_{g_{\alpha\beta}}^M$, it follows from (34) that H_θ^ϕ commutes with T_G for all $G \in \mathcal{G}$.

Let θ and $\tilde{\theta}$ be Borel real functions on A^ϕ such that $\theta(s) - \tilde{\theta}(s) \in \pi\mathbb{Z}$ for ν_0 -a.e. $s \in A^\phi$. By (24)–(29), we have $\mathcal{V}_{\kappa, \vartheta + \pi n} = \mathcal{V}_{\kappa, \vartheta}$ for every $-1 < \kappa < 1$, $\vartheta \in \mathbb{R}$, and $n \in \mathbb{Z}$. In view of (30), it follows that $\mu_\theta^\phi(s) = \mu_{\tilde{\theta}}^\phi(s)$ for ν_0 -a.e. s . The uniqueness statement of Proposition II.4 therefore implies that $M_\theta^\phi = M_{\tilde{\theta}}^\phi$. By (15) and (16), we have $u_{\vartheta + \pi n}^\kappa(z) = e^{i\pi n} u_\vartheta^\kappa(z)$ for every $-1 < \kappa < 1$, $\vartheta \in \mathbb{R}$, $z \in \mathbb{C}$, and $n \in \mathbb{Z}$. It follows from (11) and (19) that

$$\mathcal{W}_\theta^\phi(s, E|x) = f_{\theta\tilde{\theta}}(s, E) \mathcal{W}_{\tilde{\theta}}^\phi(s, E|x) \quad (36)$$

for every $s \in S \setminus N$, $E \in \mathbb{R}$, and $x \in \mathcal{O}$, where $N = \{s \in A^\phi : \theta(s) - \tilde{\theta}(s) \notin \pi\mathbb{Z}\}$ is a ν_0 -null set and the Borel function $f_{\theta\tilde{\theta}}$ on $S \times \mathbb{R}$ is given by

$$f_{\theta\tilde{\theta}}(s, E) = \begin{cases} 1, & s \in S \setminus A^\phi, \\ e^{i(\tilde{\theta}(s) - \theta(s))}, & s \in A^\phi. \end{cases}$$

It easily follows from Proposition II.4 that $N \times \mathbb{R}$ is an M -null set, where $M = M_\theta^\phi = M_{\tilde{\theta}}^\phi$, and, therefore, $f_{\theta\tilde{\theta}}(s, E) = \pm 1$ for M -a.e. (s, E) . Proposition II.1 and formula (36) hence imply that $W_\theta^\phi = \mathcal{T}_{f_{\theta\tilde{\theta}}}^M W_{\tilde{\theta}}^\phi$. In view of (34), we conclude that $H_\theta^\phi = H_{\tilde{\theta}}^\phi$.

Thus, assuming Propositions II.4 and II.1, we have proved the next statement in one direction.

Theorem II.5. *Let $\phi \in \mathbb{R}$. For every Borel real function θ on A^ϕ , the operator H_θ^ϕ is a self-adjoint extension of H^ϕ commuting with T_G for any $G \in \mathcal{G}$. Conversely, every self-adjoint extension of H^ϕ commuting with T_G for any $G \in \mathcal{G}$ is equal to H_θ^ϕ for some Borel real function θ on A^ϕ . Given Borel real functions θ and $\tilde{\theta}$ on A^ϕ , we have $H_\theta^\phi = H_{\tilde{\theta}}^\phi$ if and only if $\theta(s) - \tilde{\theta}(s) \in \pi\mathbb{Z}$ for ν_0 -a.e. $s \in A^\phi$.*

III. COMMUTATION OF OPERATORS AND VON NEUMANN ALGEBRAS

In this section, we give some background material for the treatment of diagonalizations in Sec. IV. It should be noted that, as far as diagonalizations are concerned, the conditions imposed on $\mathfrak{X}$ in Introduction are excessively restrictive. In fact, it suffices to assume (and we do so in this section and Sec. IV) that $\mathfrak{X}$ is an arbitrary set of closed densely defined operators rather than an involutive subset of $L(\mathfrak{H})$. In Sec. V, however, where the reduction by symmetries is treated, the assumptions on $\mathfrak{X}$ are the same as in Introduction.

Lemma III.1. *Let $\mathfrak{H}$ be a Hilbert space, $T \in L(\mathfrak{H})$, and R be an operator in $\mathfrak{H}$ commuting with T . If R is densely defined, then R^* commutes with T^* . If R is closable, then the closure $\bar{R}$ of R commutes with T .*

Proof. Suppose R is densely defined. Let $\Psi \in D_{R^*}$ and $\Phi = R^*\Psi$. Then we have $\langle R\Psi', \Psi \rangle = \langle \Psi', \Phi \rangle$ for any $\Psi' \in D_R$. Hence, we obtain

$$\langle R\Psi', T^*\Psi \rangle = \langle RT\Psi', \Psi \rangle = \langle T\Psi', \Phi \rangle = \langle \Psi', T^*\Phi \rangle, \quad \Psi' \in D_R.$$

This means that $T^*\Psi \in D_{R^*}$ and $R^*T^*\Psi = T^*R^*\Psi$, i.e., R^* commutes with T^* .

Suppose now that R is closable. Let $\Psi \in D_{\bar{R}}$. Then there is a sequence $\Psi_n \in D_R$ such that $\Psi_n \rightarrow \Psi$ and $R\Psi_n \rightarrow \bar{R}\Psi$ in $\mathfrak{H}$. Since R commutes with T , we have $T\Psi_n \in D_R$ for all n . The continuity of T implies that $T\Psi_n \rightarrow T\Psi$ and $RT\Psi_n = TR\Psi_n \rightarrow T\bar{R}\Psi$. This means that $T\Psi \in D_{\bar{R}}$ and $\bar{R}T\Psi = T\bar{R}\Psi$. $\square$

Given a set $\mathfrak{X}$ of closed densely defined operators in a Hilbert space $\mathfrak{H}$, let $\mathfrak{X}'$ denote its commutant, i.e., the subalgebra of $L(\mathfrak{H})$ consisting of all operators commuting with every element of $\mathfrak{X}$. Let $\mathfrak{X}^*$ be the set consisting of the adjoints of the elements of $\mathfrak{X}$. By Lemma III.1, we have

$$(\mathfrak{X}')^* = (\mathfrak{X}^*)'. \quad (37)$$

The set $\mathfrak{X}$ is involutive if and only if $\mathfrak{X}^* = \mathfrak{X}$.

Recall¹⁹ that a subalgebra $\mathcal{M}$ of $L(\mathfrak{H})$ is called a von Neumann algebra if it is involutive and coincides with its bicommutant $\mathcal{M}''$. By the well-known von Neumann's bicommutant theorem (see, e. g., Ref. 19, Sec. I.3.4, Corollaire 2), an involutive subalgebra $\mathcal{M}$ of $L(\mathfrak{H})$ is a von Neumann algebra if and only if it contains the identity operator and is closed in the strong operator topology. It follows from (37) that $\mathfrak{X}'$ is an involutive subalgebra of $L(\mathfrak{H})$ for any involutive set $\mathfrak{X}$ of closed densely defined operators in $\mathfrak{H}$. Moreover, it is easy to show (see Lemma 1 in Ref. 4) that $\mathfrak{X}'$ is always strongly closed and, therefore, is a von Neumann algebra for involutive $\mathfrak{X}$ by the bicommutant theorem.

A closed densely defined operator T in $\mathfrak{H}$ is called affiliated with a von Neumann algebra $\mathcal{M}$ if T commutes with every element of $\mathcal{M}'$. If $\mathfrak{X}$ is a set of closed densely defined operators in $\mathfrak{H}$, then every element of $\mathfrak{X}$ is obviously affiliated with the algebra $\mathcal{A}(\mathfrak{X}) = (\mathfrak{X} \cup \mathfrak{X}^*)''$. As shown by the next lemma, $\mathcal{A}(\mathfrak{X})$ is actually the smallest von Neumann algebra with this property.

Lemma III.2. *Let $\mathfrak{X}$ be a set of closed densely defined operators in a Hilbert space $\mathfrak{H}$ and $\mathcal{M}$ be a von Neumann algebra in $\mathfrak{H}$. Then $\mathcal{A}(\mathfrak{X}) \subset \mathcal{M}$ if and only if every operator in $\mathfrak{X}$ is affiliated with $\mathcal{M}$.*

Proof. If every element of $\mathfrak{X}$ is affiliated with $\mathcal{M}$, then $\mathcal{M}' \subset \mathfrak{X}'$, whence $\mathcal{M}' \subset (\mathfrak{X}^*)'$ by (37). It follows that $\mathcal{M}' \subset \mathfrak{X}' \cap (\mathfrak{X}^*)' = (\mathfrak{X} \cup \mathfrak{X}^*)'$ and, hence, $\mathcal{A}(\mathfrak{X}) \subset \mathcal{M}'' = \mathcal{M}$. Conversely, if $\mathcal{A}(\mathfrak{X}) \subset \mathcal{M}$, then $\mathcal{M}' \subset (\mathfrak{X} \cup \mathfrak{X}^*)' \subset \mathfrak{X}'$ and, hence, every element of $\mathfrak{X}$ is affiliated with $\mathcal{M}$. $\square$

The algebra $\mathcal{A}(\mathfrak{X})$ will be called the von Neumann algebra generated by $\mathfrak{X}$, and $\mathfrak{X}$ will be referred to as a set of generators of $\mathcal{A}(\mathfrak{X})$. If $\mathfrak{X} \subset L(\mathfrak{H})$, then $\mathcal{A}(\mathfrak{X})$ is just the smallest von Neumann algebra containing $\mathfrak{X}$. By Lemma III.2, a closed densely defined operator T is affiliated with a von Neumann algebra $\mathcal{M}$ if and only if $\mathcal{A}(T) \subset \mathcal{M}$ (here and subsequently, we write $\mathcal{A}(T)$ instead of $\mathcal{A}(\{T\})$, where $\{T\}$ is the one-element set containing T).

Let $\mathfrak{X} \subset L(\mathfrak{H})$ be an involutive set of pairwise commuting operators. Then $\mathfrak{X} \subset \mathfrak{X}'$ and, hence, $\mathcal{A}(\mathfrak{X}) \subset \mathfrak{X}'$ because $\mathfrak{X}'$ is a von Neumann algebra. As $\mathfrak{X}' = \mathcal{A}(\mathfrak{X})'$, it follows that $\mathcal{A}(\mathfrak{X})$ is Abelian.

We say that two sets $\mathfrak{X}$ and $\mathfrak{Y}$ of closed densely defined operators in $\mathfrak{H}$ are equivalent if $\mathcal{A}(\mathfrak{X}) = \mathcal{A}(\mathfrak{Y})$. We say that $\mathfrak{X}$ is equivalent to a closed densely defined operator T if $\mathfrak{X}$ is equivalent to the one-element set $\{T\}$.

Remark III.3. If $\mathfrak{X} \subset L(\mathfrak{H})$ is an involutive set, then the smallest strongly closed algebra containing $\mathfrak{X}$ and the identity operator in $\mathfrak{H}$ is obviously a von Neumann algebra and, hence, coincides with $\mathcal{A}(\mathfrak{X})$. The above definition of $\mathcal{A}(\mathfrak{X})$ therefore complies with that used in Introduction.

Given a spectral measure $\mathcal{E}$ (see Sec. A 8), we denote by $\mathcal{P}_{\mathcal{E}}$ the set of all operators $\mathcal{E}(A)$, where A is an $\mathcal{E}$ -measurable set. As the elements of $\mathcal{P}_{\mathcal{E}}$ pairwise commute, the algebra $\mathcal{A}(\mathcal{P}_{\mathcal{E}})$ is Abelian. For an $\mathcal{E}$ -measurable complex function g , we let $J_g^{\mathcal{E}}$ denote the integral of g with respect to $\mathcal{E}$ (see Sec. A 8).

Lemma III.4. *Let $\mathfrak{H}$ be a separable Hilbert space and $\mathcal{E}$ be a spectral measure in $\mathfrak{H}$. Then the following statements hold:*

1. *The algebra $\mathcal{A}(\mathcal{P}_{\mathcal{E}})$ coincides with the set of all $J_g^{\mathcal{E}}$, where g is an $\mathcal{E}$ -measurable $\mathcal{E}$ -essentially bounded complex function.*
2. *A closed densely defined operator T in $\mathfrak{H}$ is affiliated with $\mathcal{A}(\mathcal{P}_{\mathcal{E}})$ if and only if $T = J_g^{\mathcal{E}}$ for an $\mathcal{E}$ -measurable complex function g .*

Proof. See Lemma 10 in Ref.4. □

A family of maps $\{g_{\iota}\}_{\iota \in I}$ is said to separate points of a set S if $S \subset D_{g_{\iota}}$ for all $\iota \in I$ and for any two distinct elements s_1 and s_2 of S , there is $\iota \in I$ such that $g_{\iota}(s_1) \neq g_{\iota}(s_2)$.

To cover both positive and spectral measures, the next definition is formulated in terms of a general $\mathfrak{A}$ -valued measure (see Sec. A 2).

Definition III.5. Let $\mathfrak{A}$ be a topological Abelian group and ν be a σ -finite $\mathfrak{A}$ -valued measure. A family $\{g_{\iota}\}_{\iota \in I}$ of maps is said to be ν -separating if I is countable and $\{g_{\iota}\}_{\iota \in I}$ separates points of $S_{\nu} \setminus N$ for some ν -null set N .²⁰

The next result gives a complete description of systems of generators for $\mathcal{A}(\mathcal{P}_{\mathcal{E}})$.

Proposition III.6. *Let $\mathfrak{H}$ be a separable Hilbert space, $\mathcal{E}$ be a standard²¹ spectral measure in $\mathfrak{H}$, and $\mathfrak{X}$ be a set of closed densely defined operators in $\mathfrak{H}$. Then $\mathcal{A}(\mathfrak{X}) = \mathcal{A}(\mathcal{P}_{\mathcal{E}})$ if and only if the following conditions hold*

1. $\mathcal{A}(\mathfrak{X}) \subset \mathcal{A}(\mathcal{P}_{\mathcal{E}})$
2. *There is an $\mathcal{E}$ -separating family $\{g_{\iota}\}_{\iota \in I}$ of $\mathcal{E}$ -measurable complex functions such that $J_{g_{\iota}}^{\mathcal{E}} \in \mathfrak{X}$ for all $\iota \in I$.*

Proof. By Lemma III.2 and statement 2 of Lemma III.4, condition 1 holds if and only if every element of $\mathfrak{X}$ is equal to $J_g^{\mathcal{E}}$ for some $\mathcal{E}$ -measurable complex function g . Hence the proposition follows from Theorem 3 in Ref. 4. □

Example III.7. Let T be a normal²² operator in a separable Hilbert space and $\mathcal{E}_T$ be its spectral measure (see Sec. A 8). By Lemma III.2 and statement 2 of Lemma III.4, condition 1 of Proposition III.6 is fulfilled for every set $\mathfrak{X}$ whose elements are functions of T . Let g be the identical function on $\mathbb{C}$: $g(z) = z$, $z \in \mathbb{C}$. Then the family containing the single function g separates points of $\mathbb{C}$, and Proposition III.6 implies that the operator $T = g(T)$ is equivalent to the set $\mathcal{P}_{\mathcal{E}_T}$ of its spectral projections. Let $\zeta \in \mathbb{C}$ and h_{ζ} be the function on $\mathbb{C} \setminus \{\zeta\}$ defined by the relation $h_{\zeta}(z) = (z - \zeta)^{-1}$. If ζ is not an eigenvalue of T , then $\mathcal{E}_T(\{\zeta\}) = 0$ and the family containing the single function h_{ζ} is $\mathcal{E}_T$ -separating. It follows from Proposition III.6 that the operator $(T - \zeta)^{-1} = h_{\zeta}(T)$ is equivalent to $\mathcal{P}_{\mathcal{E}_T}$ (and, hence, to T). Let $A \subset \mathbb{C}$ be a set having an accumulation point in $\mathbb{C}$ and let $f_{\zeta}(z) = e^{\zeta z}$ for $\zeta \in A$ and $z \in \mathbb{C}$. It is easy to show (see example 7 in Ref. 4 for details) that the family $\{f_{\zeta}\}_{\zeta \in A}$ contains a countable subfamily separating the points of $\mathbb{C}$. By Proposition III.6, we conclude that the set of all operators $e^{\zeta T}$ with $\zeta \in A$ is equivalent to $\mathcal{P}_{\mathcal{E}_T}$.

IV. DIAGONALIZATIONS

Given a positive σ -finite measure ν , we say that a ν -a.e. defined family $\mathfrak{S}$ of Hilbert spaces is ν -nondegenerate if $\mathfrak{S}(s) \neq \{0\}$ for ν -a.e. s .

Definition IV.1. Let $\mathfrak{X}$ be a set of closed densely defined operators in a Hilbert space $\mathfrak{H}$. A triple $(\nu, \mathfrak{S}, V)$, where ν is a positive σ -finite measure, $\mathfrak{S}$ is a ν -nondegenerate ν -measurable family of Hilbert spaces, and V is a unitary operator from $\mathfrak{H}$ to $\int^{\oplus} \mathfrak{S}(s) d\nu(s)$, is called a diagonalization for $\mathfrak{X}$ if every $T \in \mathfrak{X}$ is equal to $V^{-1} \mathcal{T}_g^{\nu, \mathfrak{S}} V$ for some complex ν -measurable function g . A diagonalization $(\nu, \mathfrak{S}, V)$ for $\mathfrak{X}$ is called exact if condition (5) holds for any complex ν -measurable ν -essentially bounded function g .

The next theorem gives an exactness criterion for diagonalizations, similar to condition (E) discussed in Introduction in the context of direct sum decompositions.

Theorem IV.2. *Let $\mathfrak{H}$ be a separable Hilbert space and $\mathfrak{X}$ be a set of closed densely defined operators in $\mathfrak{H}$. A diagonalization $(\nu, \mathfrak{S}, V)$ for $\mathfrak{X}$, where ν is standard, is exact if and only if there is a ν -separating family $\{g_\iota\}_{\iota \in I}$ of ν -measurable complex functions such that*

$$V^{-1} \mathcal{T}_{g_\iota}^{\nu, \mathfrak{S}} V \in \mathfrak{X}, \quad \iota \in I. \quad (38)$$

Sometimes the analysis of diagonalizations for $\mathfrak{X}$ simplifies if we replace $\mathfrak{X}$ with an equivalent set $\mathfrak{Y}$. The next result shows that passing to equivalent sets is always possible.

Proposition IV.3. *Let $\mathfrak{H}$ be a separable Hilbert space and $\mathfrak{X}$ and $\mathfrak{Y}$ be equivalent sets of closed densely defined operators in $\mathfrak{H}$. Then every (exact) diagonalization for $\mathfrak{X}$ is an (exact) diagonalization for $\mathfrak{Y}$.*

We say that normal operators T_1 and T_2 in $\mathfrak{H}$ commute if their spectral projections commute (if $T_2 \in L(\mathfrak{H})$, then this definition agrees with Definition I.1, see Ref. 23). If a set $\mathfrak{X}$ of closed densely defined operators admits a diagonalization, then $\mathfrak{X}$ consists of normal pairwise commuting operators because operators of multiplication by functions are normal and commute with each other. The converse statement is provided by the next proposition.

Proposition IV.4. *For every set of normal pairwise commuting operators in a separable Hilbert space, there exists a diagonalization $(\nu, \mathfrak{S}, V)$, where ν is standard.*

Before proceeding with the proofs of the above results, we give some simple examples of diagonalizations.

Example IV.5. Let Z be the set of all absolutely continuous square-integrable complex functions on $\mathbb{R}$ having square-integrable derivatives. Let P be the one-dimensional operator of momentum, i.e., the operator in $L_2(\mathbb{R})$ with the domain $D_P = \{[f]_\lambda : f \in Z\}$ satisfying the relation

$$(P[f]_\lambda)(x) = -if'(x), \quad f \in Z,$$

for λ -a.e. x (as in Sec. II, λ is the Lebesgue measure on $\mathbb{R}$). Let $\mathcal{F}: L_2(\mathbb{R}) \rightarrow L_2(\mathbb{R})$ be the operator of the Fourier transformation: $(\mathcal{F}f)(p) = (2\pi)^{-1/2} \int f(x) e^{-ipx} dx$. Then we have $\mathcal{F}P\mathcal{F}^{-1} = \mathcal{T}_g^\lambda$, where g is the identical function on $\mathbb{R}$: $g(p) = p$, $p \in \mathbb{R}$. By Theorem IV.2, we conclude that $(\lambda, \mathcal{I}_{\mathbb{C}, \lambda}, \mathcal{F})$ is an exact diagonalization for P (here, $\mathcal{I}_{\mathbb{C}, \lambda}$ is a constant family of Hilbert spaces, see Sec. B3; by (B1), we have $L_2(\mathbb{R}) = \int^\oplus \mathcal{I}_{\mathbb{C}, \lambda}(x) dx$). In view of Example III.7 and Proposition IV.3, $(\lambda, \mathcal{I}_{\mathbb{C}, \lambda}, \mathcal{F})$ is also an exact diagonalization for the set $\{e^{iaP}\}_{a \in \mathbb{R}}$ of translations in $L_2(\mathbb{R})$.

Example IV.6. Let P , $\mathcal{F}$, and g be as in Example IV.5 and let $H = P^2$ be the Hamiltonian of the one-dimensional free particle. Then $\mathcal{F}H\mathcal{F}^{-1} = \mathcal{T}_{g^2}^\lambda$ and, hence, $(\lambda, \mathcal{I}_{\mathbb{C}, \lambda}, \mathcal{F})$ is a diagonalization for H . Since the family containing the single function g^2 is not λ -separating, this diagonalization is not exact. Let $\mathbb{R}_+ = (0, \infty)$ and $V: L_2(\mathbb{R}) \rightarrow L_2(\mathbb{R}_+, \mathbb{C}^2, \lambda)$ be the unitary operator such that $(Vf)(p) = ((\mathcal{F}f)(p), (\mathcal{F}f)(-p))$, $f \in L_2(\mathbb{R})$, for λ -a.e. $p \in \mathbb{R}_+$. Then VHV^{-1} is the operator of multiplication by g^2 in $L_2(\mathbb{R}_+, \mathbb{C}^2, \lambda)$. In view of (B1), it follows that $VHV^{-1} = \mathcal{T}_{g^2}^{\lambda_+, \mathfrak{S}}$, where $\lambda_+ = \lambda|_{\mathbb{R}_+}$ and $\mathfrak{S} = \mathcal{I}_{\mathbb{C}^2, \lambda_+}$. Since g^2 separates points of $\mathbb{R}_+$, it follows from Theorem IV.2 that $(\lambda_+, \mathcal{I}_{\mathbb{C}^2, \lambda_+}, V)$ is an exact diagonalization for H .

We now turn to the proofs of Theorem IV.2 and Propositions IV.3 and IV.4. In the rest of this section, we assume that the Hilbert space $\mathfrak{H}$ is separable. For brevity, we say that $(\nu, \mathfrak{S}, V)$ is an $\mathfrak{H}$ -triple if ν is a positive σ -finite measure, $\mathfrak{S}$ is a ν -nondegenerate ν -measurable family of Hilbert spaces, and V is a unitary operator from $\mathfrak{H}$ to $\int^\oplus \mathfrak{S}(s) d\nu(s)$.

Given an $\mathfrak{H}$ -triple $t = (\nu, \mathfrak{S}, V)$, we define the map $\mathcal{E}^t$ on $\sigma(D_\nu)$ by the relation

$$\mathcal{E}^t(A) = V^{-1} \mathcal{T}_{\chi_A}^{\nu, \mathfrak{S}} V, \quad A \in \sigma(D_\nu),$$

where χ_A is equal to unity on A and vanishes on $S_\nu \setminus A$.

Lemma IV.7. *Let $t = (\nu, \mathfrak{S}, V)$ be an $\mathfrak{H}$ -triple. Then $\mathcal{E}^t$ is a spectral measure such that $\mathcal{E}^t$ -measurable and $\mathcal{E}^t$ -null sets coincide with ν -measurable and ν -null sets respectively. For any ν -measurable complex function g , we have $J_g^{\mathcal{E}^t} = V^{-1} \mathcal{T}_g^{\nu, \mathfrak{S}} V$.*

Proof. Clearly, $\mathcal{E}^t$ is an $L(\mathfrak{H})$ -valued σ -additive function satisfying condition (a) of Sec. A 2. Clearly, $\mathcal{N}_\nu \subset \mathcal{N}_{\mathcal{E}^t}$. If $N \in \mathcal{N}_{\mathcal{E}^t}$, then $\mathcal{T}_{\chi_N}^{\nu, \mathfrak{S}} = 0$ and, hence, $\nu(N) = 0$ because $\mathfrak{S}(s) \neq 0$ for ν -a.e. s . It follows that $\mathcal{N}_\nu = \mathcal{N}_{\mathcal{E}^t}$ and, therefore, condition (b) of Sec. A 2 is also fulfilled. Thus, $\mathcal{E}^t$ is a spectral measure having the same null sets as ν . Since $D_{\mathcal{E}^t} = \sigma(D_\nu)$, we have $\sigma(D_{\mathcal{E}^t}) = \sigma(D_\nu)$, i.e., ν -measurable sets coincide with $\mathcal{E}^t$ -measurable sets. For any $A \in D_{\mathcal{E}^t}$ and $\Psi \in \mathfrak{H}$, we have $\mathcal{E}_\Psi^t(A) = \int_A \|(V\Psi)(s)\|^2 d\nu(s)$ (see Sec. A 8). Hence, a ν -measurable function g is $\mathcal{E}_\Psi^t$ -integrable if and only if $s \rightarrow \|(V\Psi)(s)\|^2 f(s)$ is a ν -integrable function, in which case we have

$$\int g(s) d\mathcal{E}_\Psi^t(s) = \int \|(V\Psi)(s)\|^2 g(s) d\nu(s). \quad (39)$$

Let g be a ν -measurable function and $\Psi \in \mathfrak{H}$. By (A1), we have

$$\Psi \in D_{J_g^{\mathcal{E}^t}} \iff |g|^2 \text{ is } \mathcal{E}_\Psi^t\text{-integrable} \iff s \rightarrow \|g(s)(V\Psi)(s)\|^2 \text{ is } \nu\text{-integrable} \iff V\Psi \in D_{\mathcal{T}_g^{\nu, \mathfrak{S}}}$$

and, therefore, the domains of $J_g^{\mathcal{E}^t}$ and $V^{-1}\mathcal{T}_g^{\nu, \mathfrak{S}}V$ coincide. Now (A2) and (39) imply that

$$\langle \Psi, J_g^{\mathcal{E}^t} \Psi \rangle = \int g(s) d\mathcal{E}_\Psi^t(s) = \langle V\Psi, \mathcal{T}_g^{\nu, \mathfrak{S}} V\Psi \rangle$$

for any $\Psi \in D_{J_g^{\mathcal{E}^t}}$. Hence, $J_g^{\mathcal{E}^t} = V^{-1}\mathcal{T}_g^{\nu, \mathfrak{S}}V$. $\square$

Proposition IV.3 follows immediately from the next lemma.

Lemma IV.8. *Let $\mathfrak{X}$ be a set of closed densely defined operators in $\mathfrak{H}$. An $\mathfrak{H}$ -triple t is an (exact) diagonalization for $\mathfrak{X}$ if and only if $\mathcal{A}(\mathfrak{X}) \subset \mathcal{A}(\mathcal{P}_{\mathcal{E}^t})$ (resp., $\mathcal{A}(\mathfrak{X}) = \mathcal{A}(\mathcal{P}_{\mathcal{E}^t})$).*

Proof. Let t be an $\mathfrak{H}$ -triple. Lemma IV.7, statement 2 of Lemma III.4, and Lemma III.2 imply that

t is a diagonalization for $\mathfrak{X} \iff$

every element of $\mathfrak{X}$ is equal to $J_g^{\mathcal{E}^t}$ for some $\mathcal{E}^t$ -measurable $g \iff$

every element of $\mathfrak{X}$ is affiliated with $\mathcal{A}(\mathcal{P}_{\mathcal{E}^t}) \iff \mathcal{A}(\mathfrak{X}) \subset \mathcal{A}(\mathcal{P}_{\mathcal{E}^t})$.

In view of statement 1 of Lemma III.4, an $\mathfrak{H}$ -triple t is an exact diagonalization for $\mathfrak{X}$ if and only if it is a diagonalization for $\mathfrak{X}$ and $\mathcal{A}(\mathfrak{X}) \supset \mathcal{A}(\mathcal{P}_{\mathcal{E}^t})$. By the above, these two conditions are equivalent to the equality $\mathcal{A}(\mathfrak{X}) = \mathcal{A}(\mathcal{P}_{\mathcal{E}^t})$. $\square$

Proof of Theorem IV.2. Let $t = (\nu, \mathfrak{S}, V)$ be a diagonalization for $\mathfrak{X}$, where ν is standard. By Lemma IV.8 and Proposition III.6, t is exact if and only if there is an $\mathcal{E}^t$ -separating family $\{g_\iota\}_{\iota \in I}$ of $\mathcal{E}^t$ -measurable complex functions such that $J_{g_\iota}^{\mathcal{E}^t} \in \mathfrak{X}$ for all $\iota \in I$. Hence, the required statement follows from Lemma IV.7. $\square$

Proof of Proposition IV.4. Let $\mathfrak{X}$ be a set of pairwise commuting normal operators in $\mathfrak{H}$ and $\mathfrak{Y} = \bigcup_{T \in \mathfrak{X}} \mathcal{P}_{\mathcal{E}_T}$, where $\mathcal{E}_T$ is the spectral measure of T (see Sec. A 8). Then $\mathfrak{Y}$ is an involutive subset of $L(\mathfrak{H})$ whose elements pairwise commute and, therefore, $\mathcal{M} = A(\mathfrak{Y})$ is an Abelian von Neumann algebra. Since $\mathcal{A}(T) = \mathcal{A}(\mathcal{P}_{\mathcal{E}_T}) \subset \mathcal{M}$ for any $T \in \mathfrak{X}$ (see Example III.7), every element of $\mathfrak{X}$ is affiliated with $\mathcal{M}$. It follows from Lemma III.2 that $\mathcal{A}(\mathfrak{X}) \subset \mathcal{M}$. Hence, the algebra $\mathcal{A}(\mathfrak{X})$ is Abelian. By Théorème 2 of Sec. II.6.2 in Ref. 19, there are a finite Borel measure ν on a compact metrizable space, a ν -measurable family $\mathfrak{S}$ of Hilbert spaces, and a unitary operator $V: \mathfrak{H} \rightarrow \int^\oplus \mathfrak{S}(s) d\nu(s)$ such that $\mathcal{A}(\mathfrak{X})$ coincides with the set of all operators $V^{-1}\mathcal{T}_g^{\nu, \mathfrak{S}}V$, where g is a ν -measurable ν -essentially bounded complex function. This means that $(\nu, \mathfrak{S}, V)$ is an exact diagonalization for $\mathcal{A}(\mathfrak{X})$. Since $\mathfrak{X}$ is equivalent to $\mathcal{A}(\mathfrak{X})$, Proposition IV.3 implies that $(\nu, \mathfrak{S}, V)$ is also an exact diagonalization for $\mathfrak{X}$. $\square$

V. REDUCTION BY SYMMETRIES

In this section, we assume that

- (A) $\mathfrak{H}$ is a Hilbert space, $\mathfrak{X}$ is an involutive subset of $L(\mathfrak{H})$, and $(\nu, \mathfrak{S}, V)$ is an exact diagonalization for $\mathfrak{X}$.

Given a ν -measurable family $\mathcal{H}$ of closed operators in $\mathfrak{S}$, we define the operator $Q_{\mathcal{H}}$ in $\mathfrak{H}$ by setting

$$Q_{\mathcal{H}} = V^{-1} \int^{\oplus} \mathcal{H}(s) d\nu(s) V.$$

The structure of closed operators in $\mathfrak{H}$ commuting with operators in $\mathfrak{X}$ and of their closed (in particular, self-adjoint) symmetry preserving extensions is described by the next theorem.

Theorem V.1. *Let (A) be satisfied. Then the following statements hold:*

1. *H is a closed operator in $\mathfrak{H}$ commuting with all elements of $\mathfrak{X}$ if and only if $H = Q_{\mathcal{H}}$ for some ν -measurable family $\mathcal{H}$ of closed operators in $\mathfrak{S}$.*
2. *Let $\mathcal{H}$ and $\tilde{\mathcal{H}}$ be ν -measurable families of closed operators in $\mathfrak{S}$. Then $Q_{\tilde{\mathcal{H}}}$ is an extension of $Q_{\mathcal{H}}$ if and only if $\tilde{\mathcal{H}}(s)$ is an extension of $\mathcal{H}(s)$ for ν -a.e. s . In particular, $Q_{\tilde{\mathcal{H}}} = Q_{\mathcal{H}}$ if and only if $\tilde{\mathcal{H}}(s) = \mathcal{H}(s)$ for ν -a.e. s .*
3. *Let $\mathcal{H}$ be a ν -measurable family of closed operators in $\mathfrak{S}$. Then $Q_{\mathcal{H}}$ is self-adjoint if and only if $\mathcal{H}(s)$ is self-adjoint for ν -a.e. s .*

Proof. Let $\mathcal{H}$ be a ν -measurable family of closed operators in $\mathfrak{S}$. Since $\mathfrak{X} \subset L(\mathfrak{H})$, every element of $\mathfrak{X}$ is representable in the form $V^{-1} \mathcal{T}_g^{\nu, \mathfrak{S}} V$, where g is ν -essentially bounded. By Proposition B.22, $\int^{\oplus} \mathcal{H}(s) d\nu(s)$ is a closed operator commuting with all $\mathcal{T}_g^{\nu, \mathfrak{S}}$ and, therefore, $Q_{\mathcal{H}}$ is a closed operator commuting with all elements of $\mathfrak{X}$. Conversely, let H be a closed operator in $\mathfrak{H}$ commuting with all elements of $\mathfrak{X}$. Let $\mathcal{M}$ denote the subalgebra of $L(\mathfrak{H})$ consisting of all operators commuting with H . By Lemma 1 in Ref. 4, $\mathcal{M}$ is strongly closed. Since $\mathfrak{X}$ is involutive, $\mathcal{A}(\mathfrak{X})$ coincides with the smallest strongly closed subalgebra of $L(\mathfrak{H})$ containing $\mathfrak{X}$ and the identity operator in $\mathfrak{H}$. We hence have $\mathcal{A}(\mathfrak{X}) \subset \mathcal{M}$, i.e., H commutes with all operators in $\mathcal{A}(\mathfrak{X})$. As the diagonalization $(\nu, \mathfrak{S}, V)$ is exact, it follows from (5) that VHV^{-1} commutes with all operators $\mathcal{T}_g^{\nu, \mathfrak{S}}$, where g is a ν -measurable ν -essentially bounded function. By Proposition B.22, we have $VHV^{-1} = \int^{\oplus} \mathcal{H}(s) d\nu(s)$ for some ν -measurable family $\mathcal{H}$ of closed operators in $\mathfrak{S}$. This means that $H = Q_{\mathcal{H}}$ and statement 1 is proved. Statements 2 and 3 follow immediately from Proposition B.21 and Corollary B.24 respectively. $\square$

In particular, Statements 1 and 2 of Theorem V.1 imply the existence and uniqueness (up to ν -equivalence) of decomposition (4) for any closed operator H commuting with all elements of $\mathfrak{X}$.

Corollary V.2. *Let (A) be satisfied and $\mathcal{H}$ be a ν -measurable family of closed operators in $\mathfrak{S}$. Then the closed (resp., self-adjoint) extensions of $Q_{\mathcal{H}}$ commuting with all elements of $\mathfrak{X}$ are precisely the operators $Q_{\tilde{\mathcal{H}}}$, where $\tilde{\mathcal{H}}$ is a ν -measurable family of operators in $\mathfrak{S}$ such that $\tilde{\mathcal{H}}(s)$ is a closed (resp., self-adjoint) extension of $\mathcal{H}(s)$ for ν -a.e. s .*

In concrete examples, H usually comes as the closure of some non-closed operator $\check{H}$. It is often possible to establish by a direct computation that $Q_{\mathcal{H}}$ is an extension of $\check{H}$ for some ν -a.e. defined family $\mathcal{H}$ of closed operators. In this case, the next proposition may be used to prove that actually equality (4) holds.

Proposition V.3. *Let (A) be satisfied, $\check{H}$ be an operator in $\mathfrak{H}$, and $\mathcal{H}$ be a ν -a.e. defined family of closed operators in $\mathfrak{S}$ such that $Q_{\mathcal{H}}$ is an extension of $\check{H}$. Suppose $D_{\check{H}}$ is taken to itself by all operators in $\mathfrak{X}$ and there is a sequence $\xi_1, \xi_2, \dots$ of elements of $V(D_{\check{H}})$ such that the linear span of $(\xi_j(s), \mathcal{H}(s)\xi_j(s))$ is dense in the graph $G_{\mathcal{H}(s)}$ of $\mathcal{H}(s)$ for ν -a.e. s . Then $\mathcal{H}$ is ν -measurable, $\check{H}$ is closable, and $\overline{\check{H}} = Q_{\mathcal{H}}$.*

Proof. Since $\xi_j \in D_{V\check{H}V^{-1}}$ for all $j = 1, 2, \dots$ and $\int^{\oplus} \mathcal{H}(s) d\nu(s)$ is an extension of $V\check{H}V^{-1}$, we have $\mathcal{H}(s)\xi_j(s) = (V\check{H}V^{-1}\xi_j)(s)$ for ν -a.e. s and, hence, $s \rightarrow \mathcal{H}(s)\xi_j(s)$ are ν -measurable sections of $\mathfrak{S}$. It follows that the family $\mathcal{H}$ is ν -measurable. Statement 1 of Theorem V.1 implies that $Q_{\mathcal{H}}$ is a closed operator commuting with all elements of $\mathfrak{X}$. Since $Q_{\mathcal{H}}$ is an extension of $\check{H}$ and $D_{\check{H}}$ is taken to itself by all operators in $\mathfrak{X}$, it follows that $\check{H}$ is closable and commutes with all elements of $\mathfrak{X}$. By Lemma III.1, we conclude that $H = \overline{\check{H}}$ also commutes with all operators in $\mathfrak{X}$. By statement 1 of Theorem V.1, there is a ν -measurable family $\mathcal{H}_0$ of closed operators in $\mathfrak{S}$ such that $H = Q_{\mathcal{H}_0}$. As $Q_{\mathcal{H}}$ is a closed extension of $\check{H}$, it is also an extension of H . By statement 2 of Theorem V.1, it follows that $\mathcal{H}(s)$ is an extension of $\mathcal{H}_0(s)$ for ν -a.e. s . Since $V(D_{\check{H}}) \subset V(D_H)$ and $V(D_H)$ coincides with the domain of $\int^{\oplus} \mathcal{H}_0(s) d\nu(s)$, we have $\xi_j(s) \in D_{\mathcal{H}_0(s)}$ and $\mathcal{H}_0(s)\xi_j(s) = \mathcal{H}(s)\xi_j(s)$ for all j and ν -a.e. s . Thus, the linear span of $(\xi_j(s), \mathcal{H}(s)\xi_j(s))$ is contained in $G_{\mathcal{H}_0(s)}$ and, hence, $G_{\mathcal{H}_0(s)}$ is dense in $G_{\mathcal{H}(s)}$ for ν -a.e. s . In view of the closedness of $\mathcal{H}_0(s)$, this implies that $\mathcal{H}_0(s) = \mathcal{H}(s)$ for ν -a.e. s . $\square$

VI. ONE-DIMENSIONAL SCHRÖDINGER OPERATORS

In this section, we recall basic facts (see, e.g., Refs. 24–26) concerning one-dimensional Schrödinger operators and then briefly discuss self-adjoint realizations of differential expression (22) and their eigenfunction expansions. In Sec. VIII, we shall use the results of Sec. V to represent self-adjoint extensions of the Aharonov-Bohm Hamiltonian as direct integrals of one-dimensional Schrödinger operators of type (22). The proof of Theorem II.5 given in Sec. X is based on combining such a representation with the analysis of the one-dimensional problem given in this section.

Let $-\infty \leq a < b \leq \infty$ and $\lambda_{a,b}$ be the restriction to (a, b) of the Lebesgue measure λ on $\mathbb{R}$. We denote by $\mathcal{D}$ the space of all complex continuously differentiable functions on (a, b) whose derivative is absolutely continuous on (a, b) (i.e., absolutely continuous on every segment $[c, d]$ with $a < c \leq d < b$). Given a locally integrable real function q on (a, b) , we denote by l_q the linear operator from $\mathcal{D}$ to the space of complex $\lambda_{a,b}$ -equivalence classes such that

$$l_q f(r) = -f''(r) + q(r)f(r) \quad (40)$$

for λ -a.e. $r \in (a, b)$. For every $c \in (a, b)$ and complex numbers z_1 and z_2 , there is a unique solution f of the equation $l_q f = 0$ such that $f(c) = z_1$ and $f'(c) = z_2$. This implies that solutions of $l_q f = 0$ constitute a two-dimensional subspace of $\mathcal{D}$. For any functions $f, g \in \mathcal{D}$, their Wronskian $W_r(f, g)$ at point $r \in (a, b)$ is defined by the relation

$$W_r(f, g) = f(r)g'(r) - f'(r)g(r).$$

Clearly, $r \rightarrow W_r(f, g)$ is an absolutely continuous function on (a, b) . If f and g are such that $r \rightarrow W_r(f, g)$ is a constant function on (a, b) (this is the case, in particular, when f and g are solutions of $l_q f = l_q g = 0$), its value will be denoted by $W(f, g)$. Let

$$\mathcal{D}_q = \{f \in \mathcal{D} : f \text{ and } l_q f \text{ are both square-integrable on } (a, b)\}. \quad (41)$$

A $\lambda_{a,b}$ -measurable complex function f is said to be left (right) square-integrable on (a, b) if $\int_a^c |f(r)|^2 dr < \infty$ (resp., $\int_c^b |f(r)|^2 dx < \infty$) for any $c \in (a, b)$. The subspace of $\mathcal{D}$ consisting of left (right) square-integrable on (a, b) functions f such that $l_q f$ is also left (resp., right) square-integrable on (a, b) will be denoted by $\mathcal{D}_q^l$ (resp., $\mathcal{D}_q^r$). We obviously have $\mathcal{D}_q = \mathcal{D}_q^l \cap \mathcal{D}_q^r$. It follows from (40) by integrating by parts that

$$\int_c^d ((l_q f)(r)g(r) - f(r)(l_q g)(r)) dr = W_d(f, g) - W_c(f, g)$$

for every $f, g \in \mathcal{D}$ and $c, d \in (a, b)$. This implies the existence of limits $W_a(f, g) = \lim_{r \downarrow a} W_r(f, g)$ and $W_b(f, g) = \lim_{r \uparrow b} W_r(f, g)$ for $f, g \in \mathcal{D}_q^l$ and $f, g \in \mathcal{D}_q^r$ respectively. Moreover, it follows that²⁷

$$\langle l_q f, [g] \rangle - \langle [f], l_q g \rangle = W_b(\bar{f}, g) - W_a(\bar{f}, g) \quad (42)$$

for any $f, g \in \mathcal{D}_q$, where $\langle \cdot, \cdot \rangle$ is the scalar product in $L_2(a, b)$.

For any linear subspace Z of $\mathcal{D}_q$, let $L_q(Z)$ be the linear operator in $L_2(a, b)$ defined by the relations

$$\begin{aligned} D_{L_q(Z)} &= \{[f] : f \in Z\}, \\ L_q(Z)[f] &= l_q f, \quad f \in Z. \end{aligned}$$

We define the minimal operator L_q by setting

$$L_q = L_q(\mathcal{D}_q^0), \quad (43)$$

where

$$\mathcal{D}_q^0 = \{f \in \mathcal{D}_q : W_a(f, g) = W_b(f, g) = 0 \text{ for any } g \in \mathcal{D}_q\}. \quad (44)$$

By (42), the operator $L_q(Z)$ is symmetric if and only if $W_a(\bar{f}, g) = W_b(\bar{f}, g)$ for any $f, g \in Z$. In particular, L_q is a symmetric operator. Moreover, L_q is closed and densely defined and its adjoint L_q^* is given by

$$L_q^* = L_q(\mathcal{D}_q) \quad (45)$$

(see Ref. 25, Lemma 9.4). If T is a symmetric extension of L_q , then L_q^* is an extension of T^* and, hence, of T . In view of (45), this implies that T is of the form $L_q(Z)$ for some subspace Z of $\mathcal{D}_q$.

If $W_a(f, g) = 0$ for any $f, g \in \mathcal{D}_q^l$, then q is said to be in the limit point case (l.p.c.) at a . Otherwise q is said to be in the limit circle case (l.c.c.) at a . Similarly, q is said to be in the l.p.c. at b if $W_b(f, g) = 0$ for any $f, g \in \mathcal{D}_q^r$ and to be in l.c.c. at b otherwise. According to the well-known Weyl alternative (see, e.g., Ref 25, Theorem 9.9), q is in l.c.c. at a if and only if all solutions of $l_q f = 0$ are left square-integrable on (a, b) (and, hence, belong to $\mathcal{D}_q^l$).

If q is in l.p.c. at both a and b , then (45) implies that L_q^* is symmetric and, therefore, L_q is self-adjoint.

Suppose now that q is in l.c.c. at a . Let $f_1, f_2 \in \mathcal{D}$ be linearly independent functions such that $l_q f_1 = l_q f_2 = 0$. Then f_1 and f_2 are left square-integrable on (a, b) and, therefore, the function

$$\delta_q^g(r) = \frac{1}{W(f_1, f_2)} \left[f_1(r) \int_a^r (l_q g)(\rho) f_2(\rho) d\rho - f_2(r) \int_a^r (l_q g)(\rho) f_1(\rho) d\rho \right] \quad (46)$$

is well-defined and belongs to $\mathcal{D}$ for any $g \in \mathcal{D}_q$. It is straightforward to check that δ_q^g does not depend on the choice of the solutions f_1 and f_2 and $l_q \delta_q^g = l_q g$. Hence, the function

$$\gamma_q^g = g - \delta_q^g \quad (47)$$

satisfies the equation

$$l_q \gamma_q^g = 0. \quad (48)$$

Let q be in l.c.c. at a and in l.p.c. at b . In this case, L_q has deficiency indices $(1, 1)$ and the self-adjoint extensions of L_q are precisely the operators (see Ref. 18, Lemma 11)

$$L_q^f = L_q(Z_q^f), \quad (49)$$

where f is a nontrivial real solution of $l_q f = 0$ and the subspace Z_q^f of $\mathcal{D}_q$ is given by

$$Z_q^f = \{g \in \mathcal{D}_q : W_a(f, g) = 0\}. \quad (50)$$

The operator L_q^f determines f uniquely up to a nonzero real coefficient. Since the deficiency indices of L_q are both equal to 1, the orthogonal complement $G_{L_q^f} \ominus G_{L_q}$ of the graph G_{L_q} of L_q in the graph $G_{L_q^f}$ of L_q^f is one-dimensional.

Lemma VI.1. *Suppose q is in l.c.c. at a and in l.p.c. at b . Let T be a self-adjoint extension of L_q and g be a real function in $\mathcal{D}_q$ such that $[g] \in D_T \setminus D_{L_q}$. Then γ_q^g is a real nontrivial solution of (48) and $T = L_q^{\gamma_q^g}$.*

Proof. The reality of γ_q^g follows from (46) and (47) because we can choose f_1 and f_2 in (46) to be real. It follows easily from (46) that

$$W_r(\delta_q^g, h) = \frac{1}{W(f_1, f_2)} \left[W_r(f_1, h) \int_a^r (l_q g)(\rho) f_2(\rho) d\rho - W_r(f_2, h) \int_a^r (l_q g)(\rho) f_1(\rho) d\rho \right], \quad r \in (a, b),$$

for any $h \in \mathcal{D}$, where f_1, f_2 are linearly independent solutions of $l_q f_{1,2} = 0$. This implies that

$$W_a(\delta_q^g, h) = 0 \quad (51)$$

for any $h \in \mathcal{D}_q^l$ and, therefore, $W_a(g, h) = W_a(\gamma_q^g, h)$. If γ_q^g were trivial, we would have $W_a(g, h) = W_b(g, h) = 0$ for any $h \in \mathcal{D}_q$ (recall that q is in l.p.c. at b) and, hence, $[g] \in D_{L_q}$ by (44) and (43). Thus, γ_q^g is nontrivial. Let f be a nontrivial real solution of $l_q f = 0$ such that $T = L_q^f$. Then we have $W_a(f, g) = 0$, and it follows from (47) and (51) that $W(f, \gamma_q^g) = 0$. This means that $\gamma_q^g = Cf$ for some real $C \neq 0$ and, therefore, $T = L_q^{\gamma_q^g}$. $\square$

If q is locally square-integrable on (a, b) , then $\mathcal{D}_q^0 \supset C_0^\infty(a, b)$, where $C_0^\infty(a, b)$ is the space of smooth functions on (a, b) with compact support. In view of (43), this implies that $([f], l_q f) \in G_{L_q}$ for any $f \in C_0^\infty(a, b)$.

Lemma VI.2. *There exists a countable set $A \subset C_0^\infty(a, b)$ such that the elements $([f], l_q f)$ with $f \in A$ are dense in the graph G_{L_q} of L_q for any locally square-integrable real function q on (a, b) .*

Proof. Given $a < \alpha \leq \beta < b$, we denote by $Y_{\alpha,\beta}$ the linear subspace of the space $C_0^\infty(a, b)$ consisting of all its elements vanishing outside of $[\alpha, \beta]$. We make $Y_{\alpha,\beta}$ a normed space by setting

$$\|f\| = \sup_{\alpha \leq r \leq \beta} (|f(r)| + |f''(r)|), \quad f \in Y_{\alpha,\beta}.$$

Let $C[\alpha, \beta]$ be the space of all continuous functions on the segment $[\alpha, \beta]$. Then $f \rightarrow (f, f'')$ is an isometric embedding of $Y_{\alpha,\beta}$ into $C[\alpha, \beta]^2$ if the latter space is endowed with the norm

$$\|(f, g)\| = \sup_{\alpha \leq r \leq \beta} (|f(r)| + |g(r)|), \quad f, g \in C[\alpha, \beta].$$

Note that the space $C[\alpha, \beta]^2$ with this norm is separable because $C[\alpha, \beta]$ endowed with its ordinary supremum norm is separable. Since every subspace of a separable metric space is separable, it follows that $Y_{\alpha,\beta}$ is separable. We now pick sequences $\alpha_1, \alpha_2, \dots$ and $\beta_1, \beta_2, \dots$ such that $a < \alpha_j \leq \beta_j < b$ for all j and $\alpha_j \rightarrow a$ and $\beta_j \rightarrow b$ as $j \rightarrow \infty$. For each $j = 1, 2, \dots$, we choose a countable dense subset A_j of Y_{α_j, β_j} and define the set A by the relation $A = \bigcup_{j=1}^\infty A_j$. If q is a locally square-integrable real function on (a, b) , then L_q is the closure of $L_q(C_0^\infty(a, b))$ by Lemma 17 of Ref. 18. This means that elements of the form $([f], l_q f)$ with $f \in C_0^\infty(a, b)$ are dense in the graph of L_q , and it suffices to prove that every such element can be approximated by $([g], l_q g)$ with $g \in A$. For this, we choose a j such that $f \in Y_{\alpha_j, \beta_j}$ and find a sequence $g_1, g_2, \dots$ of elements of A_j converging to f in Y_{α_j, β_j} . As l_q obviously induces a continuous map from Y_{α_j, β_j} to $L_2(a, b)$ and $f \rightarrow [f]$ is a continuous embedding of Y_{α_j, β_j} into $L_2(a, b)$, we conclude that $[g_k] \rightarrow [f]$ and $l_q g_k \rightarrow l_q f$ in $L_2(a, b)$ as $k \rightarrow \infty$. $\square$

We now turn to the Schrödinger operators corresponding to differential expression (22) that arises as a result of the separation of variables in the Aharonov-Bohm model. We therefore assume $a = 0$ and $b = \infty$ and define the potential q_κ on $\mathbb{R}_+$ by the relation

$$q_\kappa(r) = \frac{\kappa^2 - 1/4}{r^2} \quad (52)$$

for every $\kappa \in \mathbb{R}$. Let

$$h_\kappa = L_{q_\kappa}. \quad (53)$$

Our aim is to describe all self-adjoint extensions of h_κ and their eigenfunction expansions.

The equation $l_{q_\kappa} f = 0$ has linearly independent solutions $r^{1/2 \pm \kappa}$ for $\kappa \neq 0$ and $r^{1/2}$ and $r^{1/2} \ln r$ for $\kappa = 0$. This implies that

- (i) q_κ is in l.p.c. at both 0 and ∞ for $|\kappa| \geq 1$,
- (ii) q_κ is in l.p.c. at ∞ and in l.c.c. at 0 for $|\kappa| < 1$.

For every $E \in \mathbb{R}$, let the function $u^\kappa(E)$ on $\mathbb{R}_+$ be given by (13). By (21), $u^\kappa(E)$ are generalized eigenfunctions for differential expression (22). In particular, for $E = 0$, we have

$$l_{q_\kappa} u^\kappa(0) = 0. \quad (54)$$

Let $L_2^c(\mathbb{R}_+)$ denote the subspace of $L_2(\mathbb{R}_+)$ consisting of all its elements vanishing λ -a.e. outside some compact subset of $\mathbb{R}_+$.

If $|\kappa| \geq 1$, then the operator h_κ is self-adjoint by (i). It is well known^{24,28-30} that this operator can be diagonalized by the Hankel transformation. In terms of the functions $u^\kappa(E)$, this result can be formulated as follows (see Theorem 1 and formula (36) in Ref. 18).

Proposition VI.3. *Let $|\kappa| \geq 1$ and the positive measure $\mathcal{V}_\kappa$ on $\mathbb{R}$ be defined by (23). Then there is a unique unitary operator $U_\kappa: L_2(\mathbb{R}_+) \rightarrow L_2(\mathbb{R}, \mathcal{V}_\kappa)$ such that*

$$(U_\kappa \psi)(E) = \int_0^\infty u^{|\kappa|}(E|r) \psi(r) dr, \quad \psi \in L_2^c(\mathbb{R}_+), \quad (55)$$

for $\mathcal{V}_\kappa$ -a.e. E . We have $U_\kappa h_\kappa U_\kappa^{-1} = \mathcal{T}_\iota^{\mathcal{V}_\kappa}$, where ι is the identity function on $\mathbb{R}$ (i.e., $\iota(E) = E$ for all $E \in \mathbb{R}$).

Suppose now that $-1 < \kappa < 1$. For every $\vartheta, E \in \mathbb{R}$, let the function $u_\vartheta^\kappa(E)$ on $\mathbb{R}_+$ be defined by (15) and (16). As was shown in Sec. II, equality (21) remains valid for $u_\vartheta^\kappa(E)$ in place of $u^{\pm\kappa}(E)$ and, hence, $u_\vartheta^\kappa(E)$ are generalized eigenfunctions for differential expression (22). For $E = 0$, this means that

$$l_{q_\kappa} u_\vartheta^\kappa(0) = 0. \quad (56)$$

It follows immediately from (15) and (16) that

$$u_\vartheta^\kappa(E) = u^\kappa(E) \cos(\vartheta - \vartheta_\kappa) + w^\kappa(E) \sin(\vartheta - \vartheta_\kappa) \quad (57)$$

for all $\vartheta, E \in \mathbb{R}$, where ϑ_κ is defined by (17) and

$$w^\kappa(E) = u_{\pi/2+\vartheta_\kappa}^\kappa(E), \quad E \in \mathbb{R}. \quad (58)$$

By (56) and (58), we have

$$l_{q_\kappa} w^\kappa(0) = 0. \quad (59)$$

It follows immediately from (13), (15), (16), and (58) that

$$W(u^\kappa(0), w^\kappa(0)) = \frac{2}{\pi} \quad (60)$$

and, therefore, $u^\kappa(0)$ and $w^\kappa(0)$ are real linearly independent elements of $\mathcal{D}$ for every $-1 < \kappa < 1$. By (54), (56), (57), and (59), we conclude that the set of all real nontrivial $f \in \mathcal{D}$ such that $l_{q_\kappa} f = 0$ coincides with the set of all elements of the form $c u_\vartheta^\kappa(0)$, where $c, \vartheta \in \mathbb{R}$ and $c \neq 0$. In view of (53) and (ii), this implies that the self-adjoint extensions of h_κ are precisely the operators

$$h_{\kappa, \vartheta} = L_{q_\kappa}^{u_\vartheta^\kappa(0)}, \quad (61)$$

where $\vartheta \in \mathbb{R}$. Given $\vartheta \in \mathbb{R}$, let the positive measure $\mathcal{V}_{\kappa, \vartheta}$ on $\mathbb{R}$ be defined by (24) and (27) for $0 < |\kappa| < 1$ and $\kappa = 0$ respectively. The next proposition gives eigenfunction expansions for $h_{\kappa, \vartheta}$.

Proposition VI.4. *For every $-1 < \kappa < 1$ and $\vartheta \in \mathbb{R}$, there is a unique unitary operator $U_{\kappa, \vartheta}: L_2(\mathbb{R}_+) \rightarrow L_2(\mathbb{R}, \mathcal{V}_{\kappa, \vartheta})$ such that*

$$(U_{\kappa, \vartheta} \psi)(E) = \int_0^\infty u_\vartheta^\kappa(E|r) \psi(r) dr, \quad \psi \in L_2^c(\mathbb{R}_+), \quad (62)$$

for $\mathcal{V}_{\kappa, \vartheta}$ -a.e. E . We have $U_{\kappa, \vartheta} h_{\kappa, \vartheta} U_{\kappa, \vartheta}^{-1} = \mathcal{T}_\iota^{\mathcal{V}_{\kappa, \vartheta}}$, where ι is the identity function on $\mathbb{R}$.

Proof. See Theorem 3 and formulas (36) and (57) in Ref. 18. $\square$

Remark VI.5. Another treatment of self-adjoint realizations of (22) and their eigenfunction expansions can be found in Ref. 29. Our consideration differs from that in Ref. 29 by the choice of parametrization of self-adjoint extensions. The advantage of our choice is that the generalized eigenfunctions $u_\vartheta^\kappa(E|r)$ and the measures $\mathcal{V}_{\kappa, \vartheta}$ are continuous in κ at $\kappa = 0$.

VII. MEASURABLE FAMILIES OF ONE-DIMENSIONAL SCHRÖDINGER OPERATORS

In this section, we derive a criterion for the measurability of families of one-dimensional Schrödinger operators. In Sec. VIII, we shall use this criterion together with Corollary V.2 to obtain direct integral representations for self-adjoint extensions of H^ϕ commuting with T_G for all $G \in \mathcal{G}$.

In what follows, we fix $-\infty \leq a < b \leq \infty$ and set $\mathfrak{h} = L_2(a, b)$.

Let ν be a σ -finite positive measure. A ν -a.e. defined map ξ is said to be a ν -measurable family of functions on (a, b) if $\xi(s)$ is a locally integrable complex function on (a, b) for ν -a.e. s and $s \rightarrow \int_\alpha^\beta \xi(s|r) dr$ is a ν -measurable complex function for any $a < \alpha \leq \beta < b$.

Lemma VII.1. *Let ν be a σ -finite positive measure. Then the following statements hold:*

1. *Let ξ and η be ν -measurable families of locally square-integrable functions on (a, b) . Then $s \rightarrow \xi(s)\eta(s)$ is a ν -measurable family of functions on (a, b) .*

2. ξ is an $\mathfrak{h}$ -valued ν -measurable map if and only if ξ is a ν -measurable family of functions on (a, b) such that $\xi(s) \in \mathfrak{h}$ for ν -a.e. s .
3. Let ξ be a ν -a.e. defined map such that $\xi(s)$ is a continuous function on (a, b) for ν -a.e. s . Then ξ is a ν -measurable family of functions on (a, b) if and only if $s \rightarrow \xi(s|r)$ is a ν -measurable complex function for any $r \in (a, b)$.
4. Let $r_0 \in (a, b)$, ξ be a ν -measurable family of functions on (a, b) , and η be a ν -a.e. defined map such that, for ν -a.e. s , $\eta(s)$ is a complex function on (a, b) satisfying the equality $\eta(s|r) = \int_{r_0}^r \xi(s|t) dt$ for all $r \in (a, b)$. Then η is a ν -measurable family of functions on (a, b) .
5. Let ξ be a ν -measurable family of functions on (a, b) such that $\xi(s)$ is absolutely continuous on (a, b) for ν -a.e. s . Then $s \rightarrow \xi(s)'$ is a ν -measurable family of functions on (a, b) .

Proof. 1. For any $a < \gamma \leq \delta < b$, let $\chi_{\gamma, \delta}$ be the function on (a, b) that is equal to unity on $[\gamma, \delta]$ and vanishes outside this segment. Then the linear span of all $[\chi_{\gamma, \delta}]$ with rational γ and δ is dense in $\mathfrak{h}$. Let $e_1, e_2, \dots$ be a basis in $\mathfrak{h}$ obtained by orthogonalization of this system. Given $a < \alpha \leq \beta < b$, we have³¹

$$\begin{aligned} \int_{\alpha}^{\beta} \xi(s|r) \eta(s|r) dr &= \langle [\chi_{\alpha, \beta} \overline{\xi(s)}], [\chi_{\alpha, \beta} \eta(s)] \rangle = \\ &= \sum_{i=1}^{\infty} \langle [\chi_{\alpha, \beta} \overline{\xi(s)}], e_i \rangle \langle e_i, [\chi_{\alpha, \beta} \eta(s)] \rangle = \sum_{i=1}^{\infty} \int_{\alpha}^{\beta} \xi(s|r) e_i(r) dr \int_{\alpha}^{\beta} \overline{e_i(r)} \eta(s|r) dr \end{aligned}$$

for ν -a.e. s , where $\langle \cdot, \cdot \rangle$ is the scalar product in $\mathfrak{h}$. Since each e_i is $\lambda_{a, b}$ -equivalent to a linear combination of $\chi_{\gamma, \delta}$, the right-hand side is a ν -measurable complex function of s .

2. Let $\chi_{\alpha, \beta}$ and e_i be as in the proof of 1. If ξ is an $\mathfrak{h}$ -valued ν -measurable map, then $s \rightarrow \langle \psi, \xi(s) \rangle$ is a ν -measurable function for any $\psi \in \mathfrak{h}$. In particular, $s \rightarrow \langle [\chi_{\alpha, \beta}], \xi(s) \rangle = \int_{\alpha}^{\beta} \xi(s|r) dr$ is a ν -measurable function for any $a < \alpha \leq \beta < b$, i.e., ξ is a ν -measurable family of functions on (a, b) . Conversely, if ξ is a ν -measurable family of functions on (a, b) , then $s \rightarrow \int_a^b e_i(r) \xi(s|r) dr$ is a ν -measurable function for every $i = 1, 2, \dots$. If, in addition, $\xi(s) \in \mathfrak{h}$ for ν -a.e. s , then $\int_a^b e_i(r) \xi(s|r) dr = \langle e_i, \xi(s) \rangle$ and, therefore, ξ is an $\mathfrak{h}$ -valued ν -measurable map.

3. Let ξ be a ν -measurable family of functions on (a, b) and $r \in (a, b)$. As $\xi(s)$ is continuous at r for ν -a.e. s , we have $\xi(s|r) = \lim_{n \rightarrow \infty} n \int_{r-1/2n}^{r+1/2n} \xi(s|t) dt$ for ν -a.e. s . This implies that $s \rightarrow \xi(s|r)$ is a ν -measurable function because $s \rightarrow n \int_{r-1/2n}^{r+1/2n} \xi(s|t) dt$ are ν -measurable functions. Conversely, let $s \rightarrow \xi(s|r)$ be a ν -measurable function for every $r \in (a, b)$ and let $a < \alpha < \beta < b$. Given $n = 1, 2, \dots$ and a function f on (a, b) , we denote by $S_n(f)$ the Riemann sum $\frac{\beta-\alpha}{n} \sum_{k=1}^n f(r_n^k)$ for f on $[\alpha, \beta]$, where $r_n^k = \alpha + (\beta - \alpha)k/n$. As $\xi(s)$ is continuous for ν -a.e. s , $\int_{\alpha}^{\beta} \xi(s|r) dr$ is the limit of $S_n(\xi(s))$ for ν -a.e. s . This means that $s \rightarrow \int_{\alpha}^{\beta} \xi(s|r) dr$ is a ν -measurable function because $s \rightarrow S_n(\xi(s))$ is a ν -measurable function for every n .

4. Clearly, $\eta(s)$ is a continuous function on (a, b) for ν -a.e. s and $s \rightarrow \eta(s|r)$ is a ν -measurable function for any $r \in (a, b)$. Hence, the statement follows from 3.

5. For ν -a.e. s , the function $\xi(s)'$ is locally integrable and³² $\int_{\alpha}^{\beta} \xi'(s|r) dr = \xi(s|\beta) - \xi(s|\alpha)$. Hence, the statement follows from 3. $\square$

Lemma VII.2. Let ν be a σ -finite positive measure and v be a ν -measurable family of real locally square-integrable functions on (a, b) . Then $s \rightarrow L_{v(s)}$ is a ν -measurable family of operators in $\mathfrak{h}$.

Proof. By Lemma VI.2, there is a sequence $\varphi_1, \varphi_2, \dots$ of functions in $C_0^{\infty}(a, b)$ such that the vectors $([\varphi_j], l_q \varphi_j)$ are dense in G_{L_q} for any locally square-integrable real function q on (a, b) . For each $j = 1, 2, \dots$, let ξ_j and η_j be ν -a.e. defined maps such that $\xi_j(s) = [\varphi_j]$ and $\eta_j(s) = l_{v(s)} \varphi_j$ for ν -a.e. s . Since $v(s)$ is locally square integrable for ν -a.e. s , the vectors $(\xi_j(s), \eta_j(s))$ are dense in $G_{L_{v(s)}}$ for ν -a.e. s . By statements 1 and 5 of Lemma VII.1, η_j is a ν -measurable family of functions on (a, b) , and statement 2 of Lemma VII.1 implies that η_j is an $\mathfrak{h}$ -valued ν -measurable map for any $j = 1, 2, \dots$. As ξ_j are obviously $\mathfrak{h}$ -valued ν -measurable maps for all j , we conclude that $s \rightarrow L_{v(s)}$ is a ν -measurable family of operators in $\mathfrak{h}$. $\square$

Lemma VII.3. Let ν and v be as in Lemma VII.2 and ξ be a $\mathcal{D}$ -valued ν -a.e. defined map such that

$$l_{v(s)} \xi(s) = 0 \tag{63}$$

for ν -a.e. s . Suppose there is $r_0 \in (a, b)$ such that the functions $s \rightarrow \xi(s|r_0)$ and $s \rightarrow \xi'(s|r_0)$ are ν -measurable. Then ξ is a ν -measurable family of functions on (a, b) .

Proof. Clearly, it suffices to show that ξ is a ν -measurable family of functions on (α, β) for any real numbers α and β such that $a < \alpha < r_0 < \beta < b$. We fix such α and β and set

$$A_N = \left\{ s \in S_\nu : \int_\alpha^\beta v(s|r)^2 dr < N^2/(\beta - \alpha) \right\}$$

for each $N = 1, 2, \dots$. By statement 1 of Lemma VII.1, A_N is a ν -measurable set. By the Cauchy–Bunyakovsky inequality, we have

$$\int_\alpha^\beta |v(s|r)| dr < N, \quad s \in A_N. \quad (64)$$

Let $\nu_N = \nu|_{A_N}$. To prove our statement, it suffices to show that the set

$$Q_N = \{r \in (\alpha, \beta) : s \rightarrow \xi(s|r) \text{ and } s \rightarrow \xi'(s|r) \text{ are } \nu_N\text{-measurable functions}\}$$

coincides with (α, β) for all $N = 1, 2, \dots$. Indeed, this condition implies that $s \rightarrow \xi(s|r)$ is a ν_N -measurable function for all N and $r \in (\alpha, \beta)$. Because $S_\nu \setminus \bigcup_{N=1}^\infty A_N$ is a ν -null set, this means that $s \rightarrow \xi(s|r)$ is a ν -measurable function for every $r \in (\alpha, \beta)$, and statement 3 of Lemma VII.1 ensures that ξ is a ν -measurable family of functions on (α, β) . Note that $r_0 \in Q_N$ for all N . Hence, to prove the equality $Q_N = (\alpha, \beta)$, it suffices to verify that the set $R_{N,r} = \{\rho \in (\alpha, \beta) : |\rho - r| < 1/2N\}$ is contained in Q_N for every $N = 1, 2, \dots$ and $r \in Q_N$. Fix N and $r \in Q_N$ and let $\xi_0, \xi_1, \dots$ be $\mathcal{D}$ -valued ν -a.e. defined maps such that, for ν -a.e. s , the relations

$$\begin{aligned} \xi_0(s|\rho) &= \xi(s|r) + \xi'(s|r)(\rho - r), \\ \xi_n(s|\rho) &= \xi_0(s|\rho) + \int_r^\rho d\rho' \int_r^{\rho'} v(s|t)\xi_{n-1}(s|t) dt, \quad n = 1, 2, \dots, \end{aligned} \quad (65)$$

hold for all $\rho \in (a, b)$. As $r \in Q_N$, ξ_0 is a ν_N -measurable family of functions on (a, b) , and it follows from statements 1 and 4 of Lemma VII.1 that ξ_n is a ν_N -measurable family of functions on (a, b) for every $n = 0, 1, \dots$. Since $\xi(s)$ satisfies (63) for ν -a.e. s , it follows that, for ν -a.e. s , the equality

$$\xi(s|\rho) = \xi_0(s|\rho) + \int_r^\rho d\rho' \int_r^{\rho'} v(s|t)\xi(s|t) dt \quad (66)$$

holds for any $\rho \in (a, b)$. Let $\rho \in R_{N,r}$. It follows from (64), (65) and (66) that

$$|\xi(s|\rho) - \xi_n(s|\rho)| \leq |\xi(s|\rho) - \xi_{n-1}(s|\rho)|/2$$

for ν_N -a.e. s and all $n = 0, 1, \dots$. We hence have

$$|\xi(s|\rho) - \xi_n(s|\rho)| \leq |\xi(s|\rho) - \xi_0(s|\rho)|/2^n$$

for ν_N -a.e. s . This means that, for any $\rho \in R_{N,r}$, the sequence $\xi_n(s|\rho)$ converges to $\xi(s|\rho)$ for ν_N -a.e. s and, therefore, $s \rightarrow \xi(s|\rho)$ is a ν_N -measurable function for every $\rho \in R_{N,r}$. As $\xi'(s|\rho)$ is the limit of $k(\xi(s|\rho + 1/k) - \xi(s|\rho))$ as $k \rightarrow \infty$ for ν -a.e. s , it also follows that $s \rightarrow \xi'(s|\rho)$ is a ν_N -measurable function for every $\rho \in R_{N,r}$. Hence $R_{N,r} \subset Q_N$ and the lemma is proved. $\square$

Corollary VII.4. *Let ν and v be as in Lemma VII.2. Then there are ν -measurable families ξ_1 and ξ_2 of functions on (a, b) such that $\xi_1(s)$ and $\xi_2(s)$ are linearly independent real elements of $\mathcal{D}$ satisfying equation (63) for ν -a.e. s .*

Proof. Choose $r_0 \in (a, b)$. Let ξ_1 and ξ_2 be ν -a.e. defined $\mathcal{D}$ -valued maps such that $\xi_1(s)$ and $\xi_2(s)$ are solutions of (63) satisfying the conditions

$$\xi_1(s|r_0) = \xi_2'(s|r_0) = 1, \quad \xi_1'(s|r_0) = \xi_2(s|r_0) = 0$$

for ν -a.e. s . Obviously, $\xi_1(s)$ and $\xi_2(s)$ are linearly independent for ν -a.e. s , and Lemma VII.3 implies that ξ_1 and ξ_2 are ν -measurable families of functions on (a, b) . $\square$

Proposition VII.5. *Let ν and v be as in Lemma VII.2. Suppose $v(s)$ is in l.c.c. at a and in l.p.c. at b for ν -a.e. s . If ξ is a ν -measurable family of elements of $\mathcal{D}$ such that $\xi(s)$ is a nontrivial real solution of (63) for ν -a.e. s , then $s \rightarrow L_{v(s)}^{\xi(s)}$ is a ν -measurable family of self-adjoint operators in $\mathfrak{h}$. If $\mathcal{R}$ is a ν -measurable family of operators in $\mathfrak{h}$ such that $\mathcal{R}(s)$ is a self-adjoint extension of $L_{v(s)}$ for ν -a.e. s , then there exists a ν -measurable family ξ of elements of $\mathcal{D}$ such that $\xi(s)$ is a nontrivial real solution of (63) and $\mathcal{R}(s) = L_{v(s)}^{\xi(s)}$ for ν -a.e. s .*

Proof. By Lemma VII.2, $s \rightarrow L_{v(s)}$ is a ν -measurable family of operators in $\mathfrak{h}$. This means that there is a sequence $\zeta_1, \zeta_2, \dots$ of $\mathfrak{h} \oplus \mathfrak{h}$ -valued ν -measurable maps such that the linear span of $\zeta_1(s), \zeta_2(s), \dots$ is dense in the graph $G_{L_{v(s)}}$ of $L_{v(s)}$ for ν -a.e. s .

Let ξ be a ν -measurable family of elements of $\mathcal{D}$ such that $\xi(s)$ is a nontrivial real solution of (63) for ν -a.e. s and let τ be a smooth function on (a, b) that is equal to unity in a neighborhood of a and vanishes in a neighborhood of b . Let ν -a.e. defined maps g and h be such that

$$g(s) = [\tau\xi(s)], \quad h(s) = -[\tau''\xi(s) + 2\tau'\xi'(s)]$$

for ν -a.e. s . We obviously have $g(s) \in \mathfrak{h}$ and $h(s) \in \mathfrak{h}$ for ν -a.e. s and, therefore, statements 1, 2, and 5 of Lemma VII.1 imply that g and h are $\mathfrak{h}$ -valued ν -measurable maps. Let ζ be an $\mathfrak{h} \oplus \mathfrak{h}$ -valued ν -measurable map such that $\zeta(s) = (g(s), h(s))$ for ν -a.e. s . Note that

$$l_{v(s)}(\tau\xi(s)) = h(s) \tag{67}$$

for ν -a.e. s and, therefore, $\tau\xi(s) \in \mathcal{D}_{v(s)}$ for ν -a.e. s (see (41)). Since $\xi(s)$ is a nontrivial real solution of (63) and $\tau\xi(s)$ coincides with $\xi(s)$ in a neighborhood of a , it follows from (44) and (50) that $\tau\xi(s) \in Z_{v(s)}^{\xi(s)}$ and $\tau\xi(s) \notin \mathcal{D}_{v(s)}^0$ for ν -a.e. s . In view of (43), (49), and (67), we conclude that $\zeta(s) \in G_{L_{v(s)}^{\xi(s)}} \setminus G_{L_{v(s)}}$ for ν -a.e. s . As $G_{L_{v(s)}^{\xi(s)}} \ominus G_{L_{v(s)}}$ is one-dimensional, this implies that the linear span of the sequence $\zeta(s), \zeta_1(s), \zeta_2(s), \dots$ is dense in the graph of $L_{v(s)}^{\xi(s)}$ for ν -a.e. s . This means that $s \rightarrow L_{v(s)}^{\xi(s)}$ is a ν -measurable family of operators in $\mathfrak{h}$.

Conversely, let $\mathcal{R}$ be a ν -measurable family of operators in $\mathfrak{h}$ such that $\mathcal{R}(s)$ is a self-adjoint extension of $L_{v(s)}$ for ν -a.e. s . Then both $s \rightarrow G_{\mathcal{R}(s)}$ and $s \rightarrow G_{L_{v(s)}}$ are ν -measurable families of subspaces of $\mathfrak{h} \oplus \mathfrak{h}$. By statement 1 of Lemma B.6, $s \rightarrow G_{\mathcal{R}(s)} \ominus G_{L_{v(s)}}$ is also a ν -measurable family of subspaces of $\mathfrak{h} \oplus \mathfrak{h}$. Since $G_{\mathcal{R}(s)} \ominus G_{L_{v(s)}}$ is nontrivial for ν -a.e. s , there is an $\mathfrak{h} \oplus \mathfrak{h}$ -valued ν -measurable map η such that $\eta(s)$ is a nonzero element of $G_{\mathcal{R}(s)} \ominus G_{L_{v(s)}}$ for ν -a.e. s . Let³³ $Q = \{s \in S_\nu : \eta(s) = -\overline{\eta(s)}\}$ and ζ be a ν -a.e. defined map such that $\zeta(s) = i\eta(s)$ for ν -a.e. $s \in Q$ and $\zeta(s) = \eta(s) + \overline{\eta(s)}$ for ν -a.e. $s \in S_\nu \setminus Q$. Then ζ is an $\mathfrak{h} \oplus \mathfrak{h}$ -valued ν -measurable map such that

$$\zeta(s) = \overline{\zeta(s)} \tag{68}$$

and $\zeta(s) \neq 0$ for ν -a.e. s . Moreover, since $G_{\mathcal{R}(s)}$ and $G_{L_{v(s)}}$ are both invariant under complex conjugation, we have $\zeta(s) \in G_{\mathcal{R}(s)} \ominus G_{L_{v(s)}}$ for ν -a.e. s . Let g and h be $\mathfrak{h}$ -valued ν -measurable maps such that $\zeta(s) = (g(s), h(s))$ for ν -a.e. s . For ν -a.e. s , we have

$$g(s) \in D_{\mathcal{R}(s)} \setminus D_{L_{v(s)}}. \tag{69}$$

As $g(s) \in D_{\mathcal{R}(s)}$ for ν -a.e. s , there exists a ν -a.e. defined map $\tilde{g}$ such that $\tilde{g}(s) \in \mathcal{D}_{v(s)}$ and $g(s) = [\tilde{g}(s)]$ for ν -a.e. s . It follows from (68) that $\tilde{g}(s)$ is real for ν -a.e. s . By statement 2 of Lemma VII.1, $\tilde{g}$ and h are ν -measurable families of functions on (a, b) . Let ξ be a ν -a.e. defined map such that $\xi(s) = \gamma_{v(s)}^{\tilde{g}(s)}$ for ν -a.e. s . In view of (69), it follows from Lemma VI.1 that $\xi(s)$ is a nontrivial real solution of (63) and $\mathcal{R}(s) = L_{v(s)}^{\xi(s)}$ for ν -a.e. s . Let ξ_1 and ξ_2 be as in Corollary VII.4. Since $h(s) = \mathcal{R}(s)g(s)$, we have $h(s) = l_{v(s)}\tilde{g}(s)$ for ν -a.e. s . Equations (46) and (47) imply that, for ν -a.e. s , equality

$$\xi(s|r) = \tilde{g}(s|r) - \frac{1}{W(\xi_1(s), \xi_2(s))} \left[\xi_1(s|r) \int_a^r h(s|\rho) \xi_2(s|\rho) d\rho - \xi_2(s|r) \int_a^r h(s|\rho) \xi_1(s|\rho) d\rho \right] \tag{70}$$

holds for all $r \in (a, b)$. By statements 3 and 5 of Lemma VII.1, $s \rightarrow W(\xi_1(s), \xi_2(s))$ is a ν -measurable function. It therefore follows from statements 1 and 4 of Lemma VII.1 that ξ is a ν -measurable family of functions on (a, b) . $\square$

VIII. SELF-ADJOINT EXTENSIONS OF THE THREE-DIMENSIONAL AHARONOV-BOHM HAMILTONIAN

In this section, we use the results of Secs. V and VII to represent the self-adjoint extensions of H^ϕ commuting with all operators T_G , $G \in \mathcal{G}$, as direct integrals of one-dimensional Schrödinger operators. Let $\mathfrak{h} = L^2(\mathbb{R}_+)$ and the positive Borel measure ν_0 on $S = \mathbb{Z} \times \mathbb{R}$ be as in Sec. II. We begin by constructing

a suitable unitary operator $V: L_2(\mathbb{R}^3) \rightarrow L_2(S, \mathfrak{h}, \nu_0)$ and then prove that $(\nu_0, \mathcal{I}_{\mathfrak{h}, \nu_0}, V)$ is actually an exact diagonalization for the set of the operators T_G . After that, we use Proposition V.3 to obtain a representation of form (4) for H^ϕ . Finally, we combine Corollary V.2 and Proposition VII.5 to get an explicit description of the self-adjoint extensions of H^ϕ commuting with T_G .

We denote by λ_+ the restriction to $\mathbb{R}_+$ of the Lebesgue measure λ on $\mathbb{R}$. For $\Phi \in C_0^\infty(\mathcal{O})$, let the map $\tilde{\Phi}$ from S to $C_0^\infty(\mathbb{R}_+)$ be defined by the relation

$$\tilde{\Phi}(s|r) = \frac{\sqrt{r}}{2\pi} \int_{-\infty}^{\infty} dx_3 \int_0^{2\pi} d\varphi \Phi(r \cos \varphi, r \sin \varphi, x_3) e^{-ipx_3 - im\varphi} \quad (71)$$

for any $s = (m, p) \in S$ and $r > 0$.

Lemma VIII.1. *There is a unique unitary operator $V: L_2(\mathbb{R}^3) \rightarrow L_2(S, \mathfrak{h}, \nu_0)$ such that the equality*

$$(V[\Phi]_\Lambda)(s) = [\tilde{\Phi}(s)]_{\lambda_+} \quad (72)$$

holds for ν_0 -a.e. s for any $\Phi \in C_0^\infty(\mathcal{O})$.

Proof. Let $\Phi \in C_0^\infty(\mathcal{O})$. Then $s \rightarrow \int_c^d \tilde{\Phi}(s|r) dr$ is a continuous and, hence, ν_0 -measurable function for any $c, d \in \mathbb{R}_+$. This means that $\tilde{\Phi}$ is a ν_0 -measurable family of functions on $(0, \infty)$, and statement 2 of Lemma VII.1 implies that $s \rightarrow [\tilde{\Phi}(s)]_{\lambda_+}$ is an $\mathfrak{h}$ -valued ν_0 -measurable map. Let f be the function on $\mathbb{R}^3$ defined by the formula

$$f(x_1, x_2, p) = \int_{-\infty}^{\infty} \Phi(x_1, x_2, x_3) e^{-ipx_3} dx_3.$$

Clearly, f is continuous and, hence, Lebesgue measurable. It follows from the Fubini theorem and the Parseval identity for the one-dimensional Fourier transformation that f is square-integrable and

$$\int |f(x_1, x_2, p)|^2 dx_1 dx_2 dp = 2\pi \int |\Phi(x_1, x_2, x_3)|^2 dx_1 dx_2 dx_3. \quad (73)$$

Let g be the continuous function on $\mathbb{R} \times \mathbb{R} \times \mathbb{R}_+$ defined by the relation $g(\varphi, p, r) = f(r \cos \varphi, r \sin \varphi, p)$. By the Fubini theorem, we have

$$\tilde{\Phi}(s|r) = \frac{\sqrt{r}}{2\pi} \int_0^{2\pi} g(\varphi, p, r) e^{-im\varphi} d\varphi$$

for any $s = (m, p) \in S$ and $r \in \mathbb{R}_+$. Further, by the Fubini theorem and Parseval identity for the Fourier series expansion, we have

$$\begin{aligned} \int d\nu_0(s) \int_0^\infty |\tilde{\Phi}(s|r)|^2 dr &= \int_0^\infty dr \int_{-\infty}^\infty dp \sum_{m \in \mathbb{Z}} |\tilde{\Phi}(m, p|r)|^2 = \\ &= \frac{1}{2\pi} \int_0^\infty r dr \int_{-\infty}^\infty dp \int_0^{2\pi} |g(\varphi, p, r)|^2 d\varphi = \frac{1}{2\pi} \int |f(x_1, x_2, p)|^2 dx_1 dx_2 dp \end{aligned}$$

and in view of (73), we conclude that $s \rightarrow [\tilde{\Phi}(s)]_{\lambda_+}$ is a square-integrable map and

$$\int \|[\tilde{\Phi}(s)]_{\lambda_+}\|^2 d\nu_0(s) = \int d\nu_0(s) \int_0^\infty |\tilde{\Phi}(s|r)|^2 dr = \int |\Phi(x_1, x_2, x_3)|^2 dx_1 dx_2 dx_3.$$

Since $[\Phi]_\Lambda$ with $\Phi \in C_0^\infty(\mathcal{O})$ are dense in $L_2(\mathbb{R}^3)$, it follows that there exists a unique isometric operator $V: L_2(\mathbb{R}^3) \rightarrow L_2(S, \mathfrak{h}, \nu_0)$ satisfying (72). It remains to check that the image of V is the entire space $L_2(S, \mathfrak{h}, \nu_0)$. For any $\xi \in L_2(S, \mathfrak{h}, \nu_0)$ and $h \in \mathfrak{h}$, $s \rightarrow \langle \xi(s), h \rangle$ is a ν_0 -square integrable function because $|\langle \xi(s), h \rangle| \leq \|\xi(s)\| \|h\|$ for ν_0 -a.e. s . Given $\psi \in C_0^\infty(\mathbb{R}_+)$, $\xi \in L_2(S, \mathfrak{h}, \nu_0)$, and $m \in \mathbb{Z}$, we denote by $F_{\xi, \psi, m}$ the element of $L_2(\mathbb{R})$ such that $F_{\xi, \psi, m}(p) = \langle \xi(m, p), [\psi]_{\lambda_+} \rangle$ for λ -a.e. p . For $\psi \in C_0^\infty(\mathbb{R}_+)$, $\chi \in C_0^\infty(\mathbb{R})$, and $m \in \mathbb{Z}$, let $\Phi_{\psi, \chi, m} \in C_0^\infty(\mathcal{O})$ be such that

$$\Phi_{\psi, \chi, m}(r \cos \varphi, r \sin \varphi, x_3) = \frac{1}{\sqrt{r}} \psi(r) \chi(x_3) e^{im\varphi} \quad (74)$$

for any $r > 0$ and $x_3, \varphi \in \mathbb{R}$. Then we have

$$\tilde{\Phi}_{\psi, \chi, m}(k, p|r) = \delta_{km} \hat{\chi}(p) \psi(r), \quad (75)$$

where $\delta_{km} = 0$ for $k \neq m$ and $\delta_{km} = 1$ for $k = m$ and $\hat{\chi}(p) = \int \chi(x_3) e^{-ipx_3} dx_3$ is the Fourier transform of χ . It follows that

$$\langle \xi, V[\Phi_{\psi, \chi, m}]_{\Lambda} \rangle_{L_2(S, \mathfrak{h}, \nu_0)} = \int \hat{\chi}(p) \langle \xi(m, p), [\psi]_{\lambda_+} \rangle_{\mathfrak{h}} dp = \langle \bar{F}_{\xi, \psi, m}, [\hat{\chi}]_{\lambda} \rangle_{L_2(\mathbb{R})} \quad (76)$$

for every $\xi \in L_2(S, \mathfrak{h}, \nu_0)$. Suppose now that ξ is orthogonal to every element of $\text{Im } V$. Since $[\hat{\chi}]_{\lambda}$ with $\chi \in C_0^\infty(\mathbb{R})$ are dense in $L_2(\mathbb{R})$, equation (76) implies that $F_{\xi, \psi, m} = 0$, i.e., $\langle \xi(s), [\psi]_{\lambda_+} \rangle = 0$ for ν_0 -a.e. s for any $\psi \in C_0^\infty(\mathbb{R}_+)$. As $[\psi]_{\lambda_+}$ with $\psi \in C_0^\infty(\mathbb{R}_+)$ are dense in $\mathfrak{h}$, it follows that $\xi = 0$ and, therefore, $\text{Im } V = L_2(S, \mathfrak{h}, \nu_0)$. $\square$

Lemma VIII.2. *Let V be as in Lemma VIII.1. Then $(\nu_0, \mathcal{I}_{\mathfrak{h}, \nu_0}, V)$ is an exact diagonalization for the set of the operators T_G with $G \in \mathcal{G}$.*

Proof. For $\alpha, \beta \in \mathbb{R}$, let $G_{\alpha\beta} \in \mathcal{G}$ be defined by (35). Clearly, each element of $\mathcal{G}$ is equal to $G_{\alpha\beta}$ for some $\alpha, \beta \in \mathbb{R}$. We have

$$\widetilde{\Phi \circ G_{\alpha\beta}^{-1}}(m, p) = e^{-i\alpha m - i\beta p} \tilde{\Phi}(m, p), \quad (m, p) \in S, \quad (77)$$

for any $\Phi \in C_0^\infty(\mathcal{O})$ and, therefore,

$$VT_{G_{\alpha\beta}}V^{-1} = \mathcal{T}_{\gamma_{\alpha\beta}}^{\nu_0, \mathfrak{S}},$$

where $\mathfrak{S} = \mathcal{I}_{\mathfrak{h}, \nu_0}$ and the function $\gamma_{\alpha\beta}$ on S is given by

$$\gamma_{\alpha\beta}(m, p) = e^{-i\alpha m - i\beta p}.$$

Suppose (m, p) and (m', p') are such that $\gamma_{\alpha\beta}(m, p) = \gamma_{\alpha\beta}(m', p')$ for all $(\alpha, \beta) \in \mathbb{Q}^2$, where $\mathbb{Q}$ is the set of rational numbers. Then we have

$$e^{i\alpha(m-m') + i\beta(p-p')} = 1$$

for all $(\alpha, \beta) \in \mathbb{Q}^2$. Since $\mathbb{Q}^2$ is dense in $\mathbb{R}^2$, it follows that $m = m'$ and $p = p'$. This means that the family $\{\gamma_{\alpha, \beta}\}_{(\alpha, \beta) \in \mathbb{Q}^2}$ separates the points of S . The desired statement now follows from Theorem IV.2. $\square$

The next lemma gives a representation of form (4) for H^ϕ .

Lemma VIII.3. *Let $\phi \in \mathbb{R}$, V be as in Lemma VIII.1, and $\mathcal{H}^\phi$ be the map on S such that*

$$\mathcal{H}^\phi(m, p) = h_{m+\phi} + p^2 1_{\mathfrak{h}}, \quad (m, p) \in S, \quad (78)$$

where $1_{\mathfrak{h}}$ is the identity operator in $\mathfrak{h}$ and the operator h_κ , $\kappa \in \mathbb{R}$, is given by (53). Then $\mathcal{H}^\phi$ is a ν_0 -measurable family of operators in $\mathfrak{h}$ and we have

$$H^\phi = V^{-1} \int^\oplus \mathcal{H}^\phi(s) d\nu_0(s) V.$$

Proof. It easily follows from (7) that

$$(\mathcal{H}^\phi \Phi)(r \cos \varphi, r \sin \varphi, x_3) = \left(-\partial_{x_3}^2 - \partial_r^2 - \frac{1}{r} \partial_r - \frac{1}{r^2} (\partial_\varphi^2 + 2i\phi \partial_\varphi - \phi^2) \right) F_\Phi(r, \varphi, x_3) \quad (79)$$

for any $\Phi \in C_0^\infty(\mathcal{O})$, where the operator $\mathcal{H}^\phi$ in $C_0^\infty(\mathcal{O})$ is given by (7) and F_Φ is the smooth function on $\mathbb{R} \times \mathbb{R} \times \mathbb{R}_+$ which represents Φ in the cylindrical coordinates,

$$F_\Phi(r, \varphi, x_3) = \Phi(r \cos \varphi, r \sin \varphi, x_3).$$

Substituting (79) in (71) and integrating by parts yields

$$\widetilde{\mathcal{H}^\phi \Phi}(s) = -\tilde{\Phi}(s)'' + q_{m+\phi} \tilde{\Phi}(s) + p^2 \tilde{\Phi}(s) \quad (80)$$

for every $s = (m, p) \in S$, where q_κ , $\kappa \in \mathbb{R}$, is given by (52). Since $\tilde{\Phi}(s) \in C_0^\infty(\mathbb{R}_+)$ for all $s \in S$, it follows from (72) and (53) that $(V[\Phi]_\Lambda)(s) \in D_{\mathcal{H}^\phi(s)}$ and

$$\mathcal{H}^\phi(s)(V[\Phi]_\Lambda)(s) = l_{q_{m+\phi}} \tilde{\Phi}(s) + p^2 [\tilde{\Phi}(s)]_{\lambda_+} = [\widetilde{\mathcal{H}^\phi \Phi}(s)]_{\lambda_+} = (V[\mathcal{H}^\phi \Phi]_\Lambda)(s) = (V\check{H}^\phi[\Phi]_\Lambda)(s) \quad (81)$$

for ν_0 -a.e. $s = (m, p)$. Since every element of $D_{\tilde{H}^\phi}$ is equal to $[\Phi]_\Lambda$ for some $\Phi \in C_0^\infty(\mathcal{O})$, this implies that $V^{-1} \int^\oplus \mathcal{H}^\phi(s) d\nu_0(s) V$ is an extension of $\tilde{H}^\phi$. In view of (53), Lemma VI.2 implies the existence of a sequence $f_1, f_2, \dots$ of elements of $C_0^\infty(\mathbb{R}_+)$ such that $([f_j]_{\lambda_+}, l_{q_\kappa} f_j)$ are dense in the graph of h_κ for any $\kappa \in \mathbb{R}$. Since $p^2 1_{\mathfrak{h}}$ is a bounded everywhere defined operator in $\mathfrak{h}$, it follows that the elements $([f_j]_{\lambda_+}, \mathcal{H}^\phi(s)[f_j]_{\lambda_+})$ are dense in the graph of $\mathcal{H}^\phi(s)$ for all $s \in S$. Let χ be a nonzero element of $C_0^\infty(\mathbb{R})$ and $\hat{\chi}(p) = \int_{-\infty}^\infty \chi(x) e^{-ipx} dx$ be its Fourier transform. For every $j = 1, 2, \dots$ and $m \in \mathbb{Z}$, we define ξ_{jm} to be the ν_0 -equivalence class such that

$$\xi_{jm}(k, p) = \delta_{km} \hat{\chi}(p) [f_j]_{\lambda_+}$$

for ν_0 -a.e. (k, p) , where $\delta_{km} = 0$ for $k \neq m$ and $\delta_{km} = 1$ for $k = m$. Let $\Phi_{jm} \in C_0^\infty(\mathcal{O})$ be defined by the relation $\Phi_{jm} = \Phi_{f_j, \chi, m}$, where $\Phi_{\psi, \chi, m}$ for $\psi \in C_0^\infty(\mathbb{R}_+)$ is given by (74). It follows from (72) and (75) that $(V[\Phi_{jm}]_\Lambda)(s) = \xi_{jm}(s)$ for ν_0 -a.e. s and, hence, $V[\Phi_{jm}]_\Lambda = \xi_{jm}$. Thus, $\xi_{jm} \in V(D_{\tilde{H}^\phi})$ for all $j = 1, 2, \dots$ and $m \in \mathbb{Z}$. Since $\hat{\chi}$ is the restriction to $\mathbb{R}$ of a nontrivial entire function, the set of its zeros is at most countable. This implies that the elements $(\xi_{jm}(s), \mathcal{H}^\phi(s)\xi_{jm}(s))$ are dense in the graph of $\mathcal{H}^\phi(s)$ for ν_0 -a.e. s . The statement of the lemma therefore follows from (8) and Proposition V.3. $\square$

Let $\phi \in \mathbb{R}$ and θ be a Borel real function on A^ϕ , where A^ϕ is given by (9). We denote by $\mathcal{H}_\theta^\phi$ the operator-valued map on S such that

$$\mathcal{H}_\theta^\phi(s) = \begin{cases} \mathcal{H}^\phi(s), & s \in S \setminus A^\phi, \\ h_{m+\phi, \theta(s)} + p^2 1_{\mathfrak{h}}, & s \in A^\phi, \end{cases} \quad (82)$$

for every $s = (m, p) \in S$, where $\mathcal{H}^\phi$ is defined by (78) and the operator $h_{\kappa, \vartheta}$ for $-1 < \kappa < 1$ and $\vartheta \in \mathbb{R}$ is given by (61).

Lemma VIII.4. *Let $\phi \in \mathbb{R}$. For any Borel real function θ on A^ϕ , $\mathcal{H}_\theta^\phi$ is a ν_0 -measurable family of self-adjoint operators in $\mathfrak{h}$ such that $\mathcal{H}_\theta^\phi(s)$ is an extension of $\mathcal{H}^\phi(s)$ for every $s \in S$. If $\tilde{\mathcal{H}}$ is a ν_0 -measurable family of operators in $\mathfrak{h}$ such that $\tilde{\mathcal{H}}(s)$ is a self-adjoint extension of $\mathcal{H}^\phi(s)$ for ν_0 -a.e. s , then $\tilde{\mathcal{H}}$ is ν_0 -equivalent to $\mathcal{H}_\theta^\phi$ for some Borel real function θ on A^ϕ .*

Proof. Let v be the map on S defined by the relation

$$v(m, p) = q_{m+\phi}, \quad (m, p) \in S, \quad (83)$$

where q_κ , $\kappa \in \mathbb{R}$, is given by (52). Then v is a ν_0 -measurable family of functions on $\mathbb{R}_+$, and (53) implies that

$$L_{v(s)} = h_{m+\phi} \quad (84)$$

for any $s = (m, p) \in S$. Clearly, $v(s)$ is in l.c.c. at 0 if and only if $s \in A^\phi$. Given a Borel real function θ on A^ϕ , we define the map ξ_θ on A^ϕ by setting

$$\xi_\theta(s) = u^{m+\phi}(0) \cos(\theta(s) - \vartheta_{m+\phi}) + w^{m+\phi}(0) \sin(\theta(s) - \vartheta_{m+\phi}) \quad (85)$$

for every $s = (m, p) \in A^\phi$, where u^κ , w^κ , and ϑ_κ are given by (13), (58), and (17) respectively. By (54), (59), and (83), we have $l_{v(s)} \xi_\theta(s) = 0$ for every $s \in A^\phi$. It follows from (57), (61), (83), and (85) that

$$h_{m+\phi, \theta(s)} = L_{v(s)}^{\xi_\theta(s)}, \quad s = (m, p) \in A^\phi. \quad (86)$$

Since ξ_θ is obviously a $\nu_0|_{A^\phi}$ -measurable family of functions on $\mathbb{R}_+$, it follows from Proposition VII.5 that $s \rightarrow h_{m+\phi, \theta(s)}$ is a ν_0 -measurable family of operators in $\mathfrak{h}$ on A^ϕ . Since $(m, p) \rightarrow p^2 1_{\mathfrak{h}}$ is a ν_0 -measurable family of operators in $\mathfrak{h}$ by Lemma B.3, the ν_0 -measurability of $\mathcal{H}_\theta^\phi$ on A^ϕ follows from (82) and Lemma B.9. As $\mathcal{H}_\theta^\phi$ is ν_0 -measurable on $S \setminus A^\phi$ by (82) and Lemma VIII.3, we see that $\mathcal{H}_\theta^\phi$ is a ν_0 -measurable family of operators in $\mathfrak{h}$. It follows immediately from (78) and (82) that $\mathcal{H}_\theta^\phi(s)$ is a self-adjoint extension of $\mathcal{H}^\phi(s)$ for every $s \in S$.

Let $\tilde{\mathcal{H}}$ be a ν_0 -measurable family of operators in $\mathfrak{h}$ such that $\tilde{\mathcal{H}}(s)$ is a self-adjoint extension of $\mathcal{H}^\phi(s)$ for ν_0 -a.e. s . Let $\mathcal{R}$ be a ν_0 -a.e. defined family of operators in $\mathfrak{h}$ such that $\mathcal{R}(s) = \tilde{\mathcal{H}}(s) - p^2 1_{\mathfrak{h}}$ for ν_0 -a.e. $s = (m, p)$. By Lemma B.3, $\mathcal{R}$ is ν_0 -measurable. In view of (78) and (84), $\mathcal{R}(s)$ is a self-adjoint extension of $L_{v(s)}$ for ν_0 -a.e. s . By Proposition VII.5, there exists a ν_0 -a.e. defined map ξ on A^ϕ such that ξ is a $\nu_0|_{A^\phi}$ -measurable family of functions on $\mathbb{R}_+$, $\xi(s)$ is a nontrivial real element of $\mathcal{D}$ satisfying $l_{v(s)} \xi(s) = 0$

for ν_0 -a.e. $s \in A^\phi$, and $\mathcal{R}(s) = L_{v(s)}^{\xi(s)}$ for ν_0 -a.e. $s \in A^\phi$. By (54), (59), and (60), $u^{m+\phi}(0)$ and $w^{m+\phi}(0)$ are real linearly independent solutions of $l_{v(s)}u^{m+\phi}(0) = l_{v(s)}w^{m+\phi}(0) = 0$ for every $s = (m, p) \in A^\phi$. Hence, there exist ν_0 -a.e. defined real functions C_1 and C_2 on A^ϕ such that

$$\xi(s) = C_1(s)u^{m+\phi}(0) + C_2(s)w^{m+\phi}(0)$$

for ν_0 -a.e. $s = (m, p) \in A^\phi$. In view of (60), we have

$$C_1(s) = \frac{\pi}{2}W(\xi(s), w^{m+\phi}(0)), \quad C_2(s) = -\frac{\pi}{2}W(\xi(s), u^{m+\phi}(0))$$

for ν_0 -a.e. $s = (m, p) \in A^\phi$. It follows from statements 3 and 5 of Lemma VII.1 that C_1 and C_2 are ν_0 -measurable functions on A^ϕ . Since $\xi(s) \neq 0$, we have $C_1(s)^2 + C_2(s)^2 \neq 0$ for ν_0 -a.e. $s \in A^\phi$. Let θ be a Borel function on A^ϕ such that

$$\theta(s) = \vartheta_{m+\phi} + \tau(C_2(s)) \arccos \frac{C_1(s)}{C_1(s)^2 + C_2(s)^2}$$

for ν_0 -a.e. $s = (m, p) \in A^\phi$, where $\tau(y) = 1$ for $y \geq 0$ and $\tau(y) = -1$ for $y < 0$. We then have

$$\cos(\theta(s) - \vartheta_{m+\phi}) = \frac{C_1(s)}{C_1(s)^2 + C_2(s)^2}, \quad \sin(\theta(s) - \vartheta_{m+\phi}) = \frac{C_2(s)}{C_1(s)^2 + C_2(s)^2}$$

for ν_0 -a.e. $s = (m, p) \in A^\phi$. This means that $\xi(s)$ is proportional to the function $\xi_\theta(s)$ given by (85) and it follows from (86) that $\mathcal{R}(s) = h_{m+\phi, \theta(s)}$ for ν_0 -a.e. $s = (m, p) \in A^\phi$. Hence, $\tilde{\mathcal{H}}(s) = \mathcal{H}_\theta^\phi(s)$ for ν_0 -a.e. $s \in A^\phi$. As $\mathcal{H}^\phi(s)$ is self-adjoint for all $s \in S \setminus A^\phi$, we have $\tilde{\mathcal{H}}(s) = \mathcal{H}^\phi(s) = \mathcal{H}_\theta^\phi(s)$ for ν_0 -a.e. $s \in S \setminus A^\phi$. We therefore have $\tilde{\mathcal{H}}(s) = \mathcal{H}_\theta^\phi(s)$ for ν_0 -a.e. s . $\square$

Given $\phi \in \mathbb{R}$ and a Borel real function θ on A^ϕ , we define the operator R_θ^ϕ in $L_2(\mathbb{R}^3)$ by the relation

$$R_\theta^\phi = V^{-1} \int^{\oplus} \mathcal{H}_\theta^\phi(s) d\nu_0(s) V, \quad (87)$$

where the unitary operator $V: L_2(\mathbb{R}^3) \rightarrow L_2(S, \mathfrak{h}, \nu_0)$ is as in Lemma VIII.1.

Proposition VIII.5. *Let $\phi \in \mathbb{R}$. For any Borel real function θ on A^ϕ , R_θ^ϕ is a self-adjoint extension of H^ϕ commuting with T_G for all $G \in \mathcal{G}$. Every self-adjoint extension of H^ϕ commuting with T_G for all $G \in \mathcal{G}$ is equal to R_θ^ϕ for some Borel real function θ on A^ϕ . Given Borel real functions θ and $\tilde{\theta}$ on A^ϕ , we have $R_\theta^\phi = R_{\tilde{\theta}}^\phi$ if and only if $\theta(s) - \tilde{\theta}(s) \in \pi\mathbb{Z}$ for ν_0 -a.e. $s \in A^\phi$.*

Proof. If θ is a Borel real function on A^ϕ , then it follows immediately from Corollary V.2 and Lemmas VIII.2, VIII.3, and VIII.4 that R_θ^ϕ is a self-adjoint extension of H^ϕ commuting with T_G for all $G \in \mathcal{G}$.

Conversely, let $\tilde{\mathcal{H}}$ be a self-adjoint extension of H^ϕ commuting with T_G for all $G \in \mathcal{G}$. By Corollary V.2 and Lemmas VIII.2 and VIII.3, there is a ν_0 -measurable family $\tilde{\mathcal{H}}$ of operators in $\mathfrak{h}$ such that $\tilde{\mathcal{H}}(s)$ is a self-adjoint extension of $\mathcal{H}^\phi(s)$ for ν_0 -a.e. s and

$$\tilde{H} = V^{-1} \int^{\oplus} \tilde{\mathcal{H}}(s) d\nu_0(s) V. \quad (88)$$

By Lemma VIII.4, there is a Borel real function θ on A^ϕ such that $\tilde{\mathcal{H}}(s) = \mathcal{H}_\theta^\phi(s)$ for ν_0 -a.e. s . In view of (87) and (88), it follows that $\tilde{H} = R_\theta^\phi$.

Let θ and $\tilde{\theta}$ be Borel real functions on A^ϕ . By Proposition B.21, the equality $R_\theta^\phi = R_{\tilde{\theta}}^\phi$ holds if and only if $\mathcal{H}_\theta^\phi(s) = \mathcal{H}_{\tilde{\theta}}^\phi(s)$ for ν_0 -a.e. s . By (61) and (82), the latter condition is fulfilled if and only if $u_{\theta(s)}^{m+\phi}(0)$ is proportional to $u_{\tilde{\theta}(s)}^{m+\phi}(0)$ for ν_0 -a.e. $s = (m, p) \in A^\phi$. In view of (57), this is true if and only if $\theta(s) - \tilde{\theta}(s) \in \pi\mathbb{Z}$ for ν_0 -a.e. $s \in A^\phi$. $\square$

IX. DIRECT INTEGRALS OF MEASURES

Let ν be a σ -finite positive measure and μ be a map such that $\mu(s)$ is a σ -finite positive measure for ν -a.e. s . Suppose Σ is a σ -algebra whose every element is a $\mu(s)$ -measurable set for ν -a.e. s . Under suitable measurability conditions on μ , we shall define a new measure M , called the direct integral of μ and denoted by $(\Sigma)\text{-}\int^\oplus \mu(s) d\nu(s)$, satisfying $S_M = S_\nu \times G$, where G is the largest element of Σ . This measure has a Fubini-type property (Proposition IX.10) and can be viewed as a generalization of the direct product of measures. The main result of this section (Proposition IX.14) states that the space $L_2(S_M, M)$ can be identified with $\int^\oplus L_2(G, \mu(s)) d\nu(s)$. We shall see in Sec. X that the measure $(\mathcal{B}_\mathbb{R})\text{-}\int^\oplus \mu_\theta^\phi(s) d\nu_0(s)$, where μ_θ^ϕ is given by (30) and $\mathcal{B}_\mathbb{R}$ is the Borel σ -algebra on $\mathbb{R}$, satisfies the conditions of Proposition II.4 and, therefore, coincides with M_θ^ϕ . This will allow us to obtain a direct integral representation of the space $L_2(S \times \mathbb{R}, M_\theta^\phi)$ entering Proposition II.1 and relate it to the results of Sec. VIII.

Definition IX.1. Let ν be a σ -finite positive measure, Σ be a σ -algebra, and a map μ be such that $\mu(s)$ is a σ -finite positive measure for ν -a.e. s . Let the δ -ring $\mathcal{Q}_{\mu,\nu}^\Sigma$ be defined by the relation

$$\mathcal{Q}_{\mu,\nu}^\Sigma = \{A \in \Sigma : A \in D_{\mu(s)} \text{ for } \nu\text{-a.e. } s\}$$

We say that μ is a (ν, Σ) -measurable family of measures if $\Sigma = \sigma(\mathcal{Q}_{\mu,\nu}^\Sigma)$ and $s \rightarrow \mu(s|A)$ is a ν -measurable function for any $A \in \mathcal{Q}_{\mu,\nu}^\Sigma$.

Lemma IX.2. Let ν , Σ , μ , and $\mathcal{Q}_{\mu,\nu}^\Sigma$ be as in Definition IX.1. Let $\mathcal{K} \subset \mathcal{Q}_{\mu,\nu}^\Sigma$ be closed under finite intersections and satisfy $\sigma(\mathcal{K}) = \Sigma$. Let a map ξ be such that, for every $A \in \mathcal{Q}_{\mu,\nu}^\Sigma$, $\xi(s)$ is a $\mu(s)$ -integrable complex function on A for ν -a.e. s . If $s \rightarrow \int_A \xi(s|E) d\mu(s|E)$ is a ν -measurable function for any $A \in \mathcal{K}$, then the same is true for every $A \in \mathcal{Q}_{\mu,\nu}^\Sigma$.

If $\xi(s)$ is $\mu(s)$ -equivalent to unity for ν -a.e. s , this lemma reduces to the next result.

Corollary IX.3. Let ν , Σ , μ , and $\mathcal{K}$ be as in Lemma IX.2. If $s \rightarrow \mu(s|A)$ is a ν -measurable function for any $A \in \mathcal{K}$, then μ is a (ν, Σ) -measurable family of measures.

For the proof of Lemma IX.2, we need the next definition.

Definition IX.4. We say that a nonempty set of sets $\mathcal{Q}$ is an α -class if it satisfies the following conditions

- (1) If $A, B \in \mathcal{Q}$ and $A \cap B = \emptyset$, then $A \cup B \in \mathcal{Q}$.
- (2) If $A, B \in \mathcal{Q}$ and $B \subset A$, then $A \setminus B \in \mathcal{Q}$.
- (3) If $A_1 \supset A_2 \supset \dots$ is a nonincreasing sequence of elements of $\mathcal{Q}$, then $\bigcap_{i=1}^\infty A_i \in \mathcal{Q}$.

Remark IX.5. In the context of δ -rings, the role of α -classes is essentially the same as that of Dynkin systems (see, e.g., Sec. 2 in Ref. 34) with respect to σ -algebras. In particular, Lemma IX.6 below is an analogue of Theorem 2.4 in Ref. 34.

Lemma IX.6. Let $\mathcal{Q}$ be an α -class. If there exists a set $\mathcal{K} \subset \mathcal{Q}$ that is closed under finite intersections and satisfies $\mathcal{Q} \subset \sigma(\mathcal{K})$, then $\mathcal{Q}$ is a δ -ring.

Proof of Lemma IX.2. It is straightforward to check that the set

$$\mathcal{L} = \{A \in \mathcal{Q}_{\mu,\nu}^\Sigma : s \rightarrow \int_A \xi(s|E) d\mu(s|E) \text{ is a } \nu\text{-measurable function}\}.$$

is an α -class containing $\mathcal{K}$. As $\sigma(\mathcal{K}) = \Sigma$, we have $\mathcal{L} \subset \sigma(\mathcal{K})$, and Lemma IX.6 implies that $\mathcal{L}$ is a δ -ring. As $\sigma(\mathcal{L}) = \Sigma$, every $A \in \mathcal{Q}_{\mu,\nu}^\Sigma$ is representable as the union of a sequence $A_1 \subset A_2 \subset \dots$ of elements of $\mathcal{L}$. Hence,

$$\int_A \xi(s|E) d\mu(s|E) = \lim_{k \rightarrow \infty} \int_{A_k} \xi(s|E) d\mu(s|E)$$

for ν -a.e. s and, therefore, $s \rightarrow \int_A \xi(s|E) d\mu(s|E)$ is ν -measurable. Thus, $\mathcal{Q}_{\mu,\nu}^\Sigma = \mathcal{L}$ and the lemma is proved. $\square$

Proof of Lemma IX.6. Let $\mathcal{L}$ be the smallest α -class containing $\mathcal{K}$.³⁵ We first prove that $\mathcal{L}$ is a δ -ring. To this end, it suffices to show that $\mathcal{L}$ is closed under finite intersections because every α -class with this property is a δ -ring. For $A \in \mathcal{L}$, we set

$$\mathcal{L}^A = \{B \in \mathcal{L} : A \cap B \in \mathcal{L}\}.$$

Let $A \in \mathcal{L}$ and $B, C \in \mathcal{L}^A$ be such that $B \cap C = \emptyset$. As $(A \cap B) \cap (A \cap C) = \emptyset$ and both sets $A \cap B$ and $A \cap C$ belong to $\mathcal{L}$, the set $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ also belongs to $\mathcal{L}$ by condition 1 of Definition IX.4. This means that $B \cup C \in \mathcal{L}^A$.

If $B, C \in \mathcal{L}^A$ are such that $C \subset B$, then $A \cap (B \setminus C) = (A \cap B) \setminus (A \cap C)$. Since both $A \cap B$ and $A \cap C$ belong to $\mathcal{L}$ and $A \cap C \subset A \cap B$, we have $A \cap (B \setminus C) \in \mathcal{L}$ by condition 2 of Definition IX.4 and, hence, $B \setminus C \in \mathcal{L}^A$.

Now let $B_1 \supset B_2 \supset \dots$ be a nonincreasing sequence of elements of $\mathcal{L}^A$ and let $B = \bigcap_{i=1}^{\infty} B_i$. Since $A \cap B = \bigcap_{i=1}^{\infty} (A \cap B_i)$ and the sets $A \cap B_1, A \cap B_2, \dots$ constitute a nonincreasing sequence of elements of $\mathcal{L}$, we have $A \cap B \in \mathcal{L}$ by condition 3 of Definition IX.4 and, therefore, $B \in \mathcal{L}^A$.

It follows from the above that $\mathcal{L}^A$ is an α -class for any $A \in \mathcal{L}$. If $B \in \mathcal{K}$, then we obviously have $\mathcal{K} \subset \mathcal{L}^B$ and, hence, $\mathcal{L}^B = \mathcal{L}$. It follows that $A \cap B \in \mathcal{L}$ for arbitrary $A \in \mathcal{L}$ and $B \in \mathcal{K}$ and, therefore, $B \in \mathcal{L}^A$. This means that $\mathcal{K} \subset \mathcal{L}^A$ for every $A \in \mathcal{L}$. Hence, $\mathcal{L}^A = \mathcal{L}$ for every $A \in \mathcal{L}$ and, consequently, $A \cap B \in \mathcal{L}$ for every $A, B \in \mathcal{L}$. Thus, $\mathcal{L}$ is a δ -ring.

We now show that $\mathcal{Q}$ is a δ -ring. As for $\mathcal{L}$, it suffices to show that $A \cap B \in \mathcal{Q}$ for any $A, B \in \mathcal{Q}$. As $\mathcal{L}$ is a δ -ring, every element of $\sigma(\mathcal{K}) = \sigma(\mathcal{L})$ is a countable union of elements of $\mathcal{L}$. Let $A_1 \subset A_2 \subset \dots$ and $B_1 \subset B_2 \subset \dots$ be nondecreasing sequences of elements of $\mathcal{L}$ such that $A = \bigcup_{i=1}^{\infty} A_i$ and $B = \bigcup_{i=1}^{\infty} B_i$. As $\mathcal{L}$ is a ring, $C_i = A_i \cap B_i$ belongs to $\mathcal{L}$ (and, hence, to $\mathcal{Q}$) for every $i = 1, 2, \dots$. It follows from condition 2 of Definition IX.4 that $C'_i = A_i \setminus C_i$ belongs to $\mathcal{Q}$ for every $i = 1, 2, \dots$. As $C'_1 \supset C'_2 \supset \dots$, condition 3 of Definition IX.4 implies that $A \setminus B = \bigcap_{i=1}^{\infty} C'_i$ belongs to $\mathcal{Q}$. Applying condition 2 of Definition IX.4 again, we conclude that $A \cap B = A \setminus (A \setminus B)$ belongs to $\mathcal{Q}$. $\square$

Given sets of sets $\mathcal{Q}_1$ and $\mathcal{Q}_2$, we denote by $\mathcal{Q}_1 \boxtimes \mathcal{Q}_2$ the set of all sets $A_1 \times A_2$, where $A_1 \in \mathcal{Q}_1$ and $A_2 \in \mathcal{Q}_2$. Let A be a set. For any s , we define its section A_s by the relation $A_s = \{E : (s, E) \in A\}$.

Let ν be a σ -finite positive measure, Σ be a σ -algebra, and μ be a (ν, Σ) -measurable family of measures. We set $\Xi_{\mu, \nu}^{\Sigma} = \sigma(D_{\nu} \boxtimes \Sigma)$ and denote by $\Delta_{\mu, \nu}^{\Sigma}$ the set of all $A \in \Xi_{\mu, \nu}^{\Sigma}$ such that $A_s \in D_{\mu(s)}$ for ν -a.e. s and $s \rightarrow \mu(s|A_s)$ is a ν -integrable function.

Proposition IX.7. *Let ν be a σ -finite positive measure, Σ be a σ -algebra, and μ be a (ν, Σ) -measurable family of measures. Then there is a unique $\Xi_{\mu, \nu}^{\Sigma}$ -compatible measure M such that $\Delta_{\mu, \nu}^{\Sigma} \subset D_M$ and*

$$M(A) = \int \mu(s|A_s) d\nu(s) \quad (89)$$

for any $A \in \Delta_{\mu, \nu}^{\Sigma}$. Moreover, the equality $\Xi_{\mu, \nu}^{\Sigma} \cap D_M = \Delta_{\mu, \nu}^{\Sigma}$ is fulfilled. If ν is $\tilde{\Sigma}$ -compatible for some σ -algebra $\tilde{\Sigma}$, then M is $\sigma(\tilde{\Sigma} \boxtimes \Sigma)$ -compatible.

Proof. Let $\mathcal{Q} = \mathcal{Q}_{\mu, \nu}^{\Sigma}$, $\Xi = \Xi_{\mu, \nu}^{\Sigma}$, $\Delta = \Delta_{\mu, \nu}^{\Sigma}$, and $\mathcal{K} = \Delta \cap (D_{\nu} \boxtimes \mathcal{Q})$. Since $\mathcal{Q}$ is a δ -ring, $\mathcal{K}$ is closed under finite intersections. Given $B \in D_{\nu}$, $C \in \mathcal{Q}$, and $n = 1, 2, \dots$, we set $B_n = \{s \in B : \mu(s|C) \leq n\}$. Clearly, $B = N \cup \bigcup_n B_n$, where N is a ν -null set. Since $B_n \times C \in \mathcal{K}$ for all n and $N \times C \in \mathcal{K}$, we conclude that $B \times C \in \sigma(\mathcal{K})$. Thus, $D_{\nu} \boxtimes \mathcal{Q} \subset \sigma(\mathcal{K})$. As $\sigma(\mathcal{Q}) = \Sigma$, it follows that $\sigma(\mathcal{K}) = \Xi$. As Δ is obviously an α -class, Lemma IX.6 implies that Δ is a δ -ring. Let the function $m: \Delta \rightarrow \mathbb{R}$ be such that $m(A)$ is equal to the right-hand side of (89) for any $A \in \Delta$. It follows from the monotone convergence theorem that m is a σ -additive function satisfying condition (a) of Sec. A 2. Hence, the existence and uniqueness of the measure M , as well as the equality $\Xi \cap D_M = \Delta$, are ensured by Lemma A.3. Now suppose ν is $\tilde{\Sigma}$ -compatible and let $\mathcal{L}$ denote the set of all $A \in \Xi$ such that there exist $B_1, B_2 \in \sigma(\tilde{\Sigma} \boxtimes \Sigma)$ satisfying the conditions $B_1 \subset A \subset B_2$ and $M(B_2 \setminus B_1) = 0$. Then $\mathcal{L}$ is a σ -algebra that contains $D_{\nu} \boxtimes \Sigma$ and, hence, coincides with Ξ . In view of the Ξ -compatibility of M , this implies that M is $\sigma(\tilde{\Sigma} \boxtimes \Sigma)$ -compatible. $\square$

Definition IX.8. Under the conditions of Proposition IX.7, we call the measure M the direct integral of μ with respect to (ν, Σ) and denote it by $(\Sigma)\text{-}\int^{\oplus} \mu(s) d\nu(s)$.

Lemma IX.9. *Let ν be a σ -finite positive measure, Σ be a σ -algebra, μ be a (ν, Σ) -measurable family of measures, and $M = (\Sigma)\text{-}\int^{\oplus} \mu(s) d\nu(s)$. An M -measurable set N is an M -null set if and only if N_s is a $\mu(s)$ -null set for ν -a.e. s .*

Proof. Let $\Xi = \Xi_{\mu, \nu}^{\Sigma}$ and N be an M -measurable set. By the Ξ -compatibility of M , there is a set $N' \in \Xi$ such that $N' \supset N$ and $N' \setminus N$ is an M -null set. If N is an M -null set, then N' is also an M -null set, and it follows from Proposition IX.7 that $N'_s \in D_{\mu(s)}$ for ν -a.e. s and $\int \mu(s|N'_s) d\nu(s) = 0$, i.e., N'_s is a $\mu(s)$ -null set for ν -a.e. s . Since $N_s \subset N'_s$, this implies that N_s is a $\mu(s)$ -null set for ν -a.e. s . Conversely, suppose N_s is a $\mu(s)$ -null set for ν -a.e. s . By the above $(N' \setminus N)_s$ is a $\mu(s)$ -null set and, hence, N'_s is a $\mu(s)$ -null set for ν -a.e. s . Proposition IX.7 now implies that $M(N') = 0$ and, therefore, N is an M -null set. $\square$

Proposition IX.10. *Let ν , Σ , μ , and M be as in Lemma IX.9. For any M -integrable complex function f , the function $E \rightarrow f(s, E)$ is $\mu(s)$ -integrable for ν -a.e. s and we have*

$$\int f(s, E) dM(s, E) = \int d\nu(s) \int f(s, E) d\mu(s|E). \quad (90)$$

Proof. Let $\Xi = \Xi_{\mu, \nu}^{\Sigma}$. As M is Ξ -compatible, every $A \in D_M$ is representable in the form $A = B \cup N$, where $B \in D_M \cap \Xi$ and N is an M -null set. By Proposition IX.7, we have $B \in \Delta_{\mu, \nu}^{\Sigma}$. Hence, $B_s \in D_{\mu(s)}$ for ν -a.e. s , the function $s \rightarrow \mu(s|B_s)$ is ν -integrable, and $M(B) = \int \mu(s|B_s) d\nu(s)$. Since $A_s = B_s \cup N_s$, it follows from Lemma IX.9 that $A_s \in D_{\mu(s)}$ and $\mu(s|A_s) = \mu(s|B_s)$ for ν -a.e. s . Hence, the function $s \rightarrow \mu(s|A_s)$ is ν -integrable, and $M(A) = \int \mu(s|A_s) d\nu(s)$. This proves the proposition for $f = X_A$, where X_A is the function on S_M that is equal to unity on A and vanishes on $S_M \setminus A$. We say that a function f on S_M is simple if it can be represented in the form $f = \sum_{i=1}^n c_i X_{A_i}$ with $c_i \in \mathbb{C}$ and $A_i \in D_M$. Clearly, the proposition is true if f is a simple function. If f is a nonnegative M -integrable function, then there is a nondecreasing sequence f_k of nonnegative simple functions such that $f_k(s, E) \leq f(s, E)$ and $f_k(s, E) \rightarrow f(s, E)$ as $k \rightarrow \infty$ for M -a.e. (s, E) . Let g_k be a ν -integrable function such that $g_k(s) = \int f_k(s, E) d\mu(s|E)$ for ν -a.e. s . The dominated convergence theorem implies that

$$\int f(s, E) dM(s, E) = \lim_{k \rightarrow \infty} \int f_k(s, E) dM(s, E) = \lim_{k \rightarrow \infty} \int g_k(s) d\nu(s).$$

By the monotone convergence theorem, there exists a ν -integrable function g such that $g(s) = \lim_{k \rightarrow \infty} g_k(s)$ for ν -a.e. s , and we have

$$\int f(s, E) dM(s, E) = \int g(s) d\nu(s). \quad (91)$$

By Lemma IX.9, for ν -a.e. s , the relations $f_k(s, E) \leq f(s, E)$ and $f_k(s, E) \rightarrow f(s, E)$ hold for $\mu(s)$ -a.e. E . In view of the existence of $\lim_{k \rightarrow \infty} g_k(s)$, the monotone convergence theorem implies that the function $E \rightarrow f(s, E)$ is $\mu(s)$ -integrable and $\int f(s, E) d\mu(s|E) = g(s)$ for ν -a.e. s . Substituting this equality in (91) yields (90). To complete the proof, it remains to note that every M -integrable function is a linear combination of nonnegative M -integrable functions. $\square$

Corollary IX.11. *Let ν , Σ , μ , and M be as in Lemma IX.9 and f be an M -measurable complex function. Then $E \rightarrow f(s, E)$ is a $\mu(s)$ -measurable complex function for ν -a.e. s . If $E \rightarrow f(s, E)$ is $\mu(s)$ -integrable for ν -a.e. s , then $s \rightarrow \int f(s, E) d\mu(s|E)$ is a ν -measurable function. If, in addition, f is nonnegative and $s \rightarrow \int f(s, E) d\mu(s|E)$ is a ν -integrable function, then f is M -integrable.*

Proof. Let $A_1 \subset A_2 \subset \dots$ be a sequence of elements of D_M such that $S_M = \bigcup_{k=1}^{\infty} A_k$. Let $B_k = A_k \cap \{(s, E) \in S_M : |f(s, E)| \leq k\}$ and $f_k = X_{B_k} f$, where X_{B_k} is as in the proof of Proposition IX.10. Then S_M coincides with $\bigcup_{k=1}^{\infty} B_k$ up to an M -null set and, therefore, $f_k(s, E) \rightarrow f(s, E)$ as $k \rightarrow \infty$ for M -a.e. (s, E) . Since f_k are M -integrable for all k , Lemma IX.9 and Proposition IX.10 imply that, for ν -a.e. s , the functions $E \rightarrow f_k(s, E)$ are $\mu(s)$ -integrable and converge $\mu(s)$ -a.e. to the function $E \rightarrow f(s, E)$. Hence, the latter is $\mu(s)$ -measurable for ν -a.e. s . If it is also $\mu(s)$ -integrable for ν -a.e. s , then it follows from the dominated convergence theorem that $\int f(s, E) d\mu(s|E) = \lim_{k \rightarrow \infty} \int f_k(s, E) d\mu(s|E)$ for ν -a.e. s . Since $s \rightarrow \int f_k(s, E) d\mu(s|E)$ are ν -measurable functions by Proposition IX.10, we conclude that $s \rightarrow \int f(s, E) d\mu(s|E)$ is ν -measurable. If f is nonnegative and the function $s \rightarrow \int f(s, E) d\mu(s|E)$ is ν -integrable, then Proposition IX.10 implies that

$$\int f_k(s, E) dM(s, E) \leq \int d\nu(s) \int f(s, E) d\mu(s|E)$$

for all k . The M -integrability of f therefore follows from the monotone convergence theorem. $\square$

Lemma IX.12. *Let $\mathcal{K}$ be a set of sets closed under finite intersections and ν be a positive σ -finite $\sigma(\mathcal{K})$ -compatible measure. Suppose f is a ν -measurable complex function that is ν -integrable on every set in $\mathcal{K}$. If $\int_A f(s) d\nu(s) = 0$ for every $A \in \mathcal{K}$, then $f(s) = 0$ for ν -a.e. s .*

Proof. If f satisfies the conditions of the lemma, then the same is true for its real and imaginary parts. So we can assume that f is real. Let

$$\mathcal{Q} = \{A \in \sigma(\mathcal{K}) : f \text{ is } \nu\text{-integrable on } A \text{ and } \int_A f(s) d\nu(s) = 0\}.$$

Clearly, $\mathcal{Q}$ is an α -class satisfying $\mathcal{K} \subset \mathcal{Q} \subset \sigma(\mathcal{K})$ and, therefore, is a δ -ring by Lemma IX.6. As ν is $\sigma(\mathcal{K})$ -compatible and f is real, every $A \in \mathcal{Q}$ can be represented in the form $A = A_+ \cup A_-$, where $A_{\pm} \in \sigma(\mathcal{K})$ are such that $f(s) \geq 0$ for ν -a.e. $s \in A_+$ and $f(s) \leq 0$ for ν -a.e. $s \in A_-$. Since $A_{\pm}$ are elements of $\sigma(\mathcal{Q})$ contained in an element of $\mathcal{Q}$, we conclude that $A_{\pm} \in \mathcal{Q}$ and, therefore, $\int_{A_{\pm}} f(s) d\nu(s) = 0$. This implies that $f(s) = 0$ for ν -a.e. $s \in A$. Since S_{ν} is, up to a ν -null set, a countable union of elements of $\mathcal{Q}$, we have $f(s) = 0$ for ν -a.e. s . $\square$

Proposition IX.13. *Let ν be a σ -finite positive measure, Σ be a countably generated σ -algebra, and μ be a (ν, Σ) -measurable family of measures such that $\mu(s)$ is Σ -compatible for ν -a.e. s . Then there is a unique (up to ν -identity) ν -measurable family $\mathfrak{S}$ of Hilbert spaces satisfying the conditions:*

- (1) $\mathfrak{S}(s) = L_2(G, \mu(s))$ for ν -a.e. s , where G is the largest set in Σ .
- (2) A ν -a.e. defined section ξ of $\mathfrak{S}$ is ν -measurable if and only if $s \rightarrow \int_A \xi(s|E) d\mu(s|E)$ is a ν -measurable function for any $A \in \mathcal{Q}_{\mu, \nu}^{\Sigma}$.

Let $\mathcal{K} \subset \mathcal{Q}_{\mu, \nu}^{\Sigma}$ be closed under finite intersections and satisfy $\sigma(\mathcal{K}) = \Sigma$. If ξ is a ν -a.e. defined section of $\mathfrak{S}$ such that $s \rightarrow \int_A \xi(s|E) d\mu(s|E)$ is a ν -measurable function for any $A \in \mathcal{K}$, then ξ is ν -measurable.

Proof. Let $\mathcal{L}_0$ be a countable set generating Σ . Without loss of generality, we can assume that $\mathcal{L}_0$ is closed under finite intersections. Let $B_1 \subset B_2 \subset \dots$ be a sequence of elements of $\mathcal{Q}_{\mu, \nu}^{\Sigma}$ such that $G = \bigcup_{i=1}^{\infty} B_i$ and let $\mathcal{L}$ be the set of all sets of the form $A \cap B_i$ for some $A \in \mathcal{L}_0$ and $i = 1, 2, \dots$. Then $\mathcal{L}$ is a countable subset of $\mathcal{Q}_{\mu, \nu}^{\Sigma}$ that is closed under finite intersections and satisfies $\Sigma = \sigma(\mathcal{L})$. Let $\mathfrak{S}$ be ν -a.e. defined family of Hilbert spaces satisfying (1). We choose a numbering $A_1, A_2, \dots$ of the set $\mathcal{L}$ and endow $\mathfrak{S}$ with a ν -measurable structure by setting

$$\xi_i^{\mathfrak{S}}(s) = [\chi_{A_i}]_{\mu(s)}, \quad i = 1, 2, \dots,$$

for ν -a.e. s , where the function χ_{A_i} is equal to unity on A_i and vanishes on $G \setminus A_i$. Since $\langle \xi_i^{\mathfrak{S}}(s), \xi_j^{\mathfrak{S}}(s) \rangle = \mu(s|A_i \cap A_j)$ and $\mathcal{Q}_{\mu, \nu}^{\Sigma}$ is a ring, $s \rightarrow \langle \xi_i^{\mathfrak{S}}(s), \xi_j^{\mathfrak{S}}(s) \rangle$ is a ν -measurable function for any $i, j = 1, 2, \dots$. If $\psi \in \mathfrak{S}(s)$ is orthogonal to $\xi_i^{\mathfrak{S}}(s)$ for all i , then Lemma IX.12 implies that $[\psi]_{\mu(s)} = 0$. This means that the linear span of the sequence $\xi_1^{\mathfrak{S}}(s), \xi_2^{\mathfrak{S}}(s), \dots$ is dense in $\mathfrak{S}(s)$ for ν -a.e. s . Thus, $\mathfrak{S}$ endowed with the sequence $\xi_1^{\mathfrak{S}}, \xi_2^{\mathfrak{S}}, \dots$ is indeed a ν -measurable family of Hilbert spaces. If ξ is a ν -measurable section of $\mathfrak{S}$, then $s \rightarrow \langle \xi(s), [\chi_A]_{\mu(s)} \rangle = \int_A \xi(s|E) d\mu(s|E)$ is a ν -measurable function for any $A \in \mathcal{L}$, and Lemma IX.2 implies that this is also true for every $A \in \mathcal{Q}_{\mu, \nu}^{\Sigma}$. Thus, $\mathfrak{S}$ satisfies (2). Suppose now that $\mathcal{K} \subset \mathcal{Q}_{\mu, \nu}^{\Sigma}$ is closed under finite intersections and satisfies $\sigma(\mathcal{K}) = \Sigma$. By Lemma IX.2, if ξ is a ν -a.e. defined section of $\mathfrak{S}$ such that $s \rightarrow \int_A \xi(s|E) d\mu(s|E)$ is a ν -measurable function for any $A \in \mathcal{K}$, then this is also true for any $A \in \mathcal{Q}_{\mu, \nu}^{\Sigma}$ and it follows from (2) that ξ is ν -measurable. $\square$

Proposition IX.14. *Let ν , μ , Σ , and $\mathfrak{S}$ be as in Proposition IX.13. Let $\mathfrak{H} = \int^{\oplus} \mathfrak{S}(s) d\nu(s)$ and $M = (\Sigma)\text{-}\int^{\oplus} \mu(s) d\nu(s)$. Given an M -measurable function f , let $\hat{f}$ denote the ν -equivalence class such that $\hat{f}(s)$ is the $\mu(s)$ -equivalence class of the map $E \rightarrow f(s, E)$ for ν -a.e. s . Then the following statements hold:*

1. M -measurable functions f_1 and f_2 are M -equivalent if and only if $\hat{f}_1 = \hat{f}_2$.
2. If $f \in L_2(S_M, M)$, then $\hat{f} \in \mathfrak{H}$ and the operator $Q: L_2(S_M, M) \rightarrow \mathfrak{H}$ taking f to $\hat{f}$ is unitary.
3. If g is an M -measurable complex function, then $\hat{g}(s)$ is $\mu(s)$ -measurable for ν -a.e. s , $s \rightarrow \mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ is a ν -measurable family of operators in $\mathfrak{S}$, and

$$Q\mathcal{T}_g^M Q^{-1} = \int^{\oplus} \mathcal{T}_{\hat{g}(s)}^{\mu(s)} d\nu(s).$$

Proof. 1. Since f_1 and f_2 are M -measurable, the set $N = \{(s, E) \in S_M : f_1(s, E) \neq f_2(s, E)\}$ is M -measurable. By Lemma IX.9, f_1 and f_2 are M -equivalent if and only if N_s is a $\mu(s)$ -null set for ν -a.e. s . Clearly, this condition holds if and only if $\hat{f}_1(s) = \hat{f}_2(s)$ for ν -a.e. s and, hence, $\hat{f}_1 = \hat{f}_2$.

2. Let $f \in L_2(S_M, M)$. By Corollary IX.11, $\hat{f}(s)$ is a $\mu(s)$ -measurable function for ν -a.e. s . Since $|f|^2$ is M -integrable, Proposition IX.10 implies that $|\hat{f}(s)|^2$ is $\mu(s)$ -integrable and, therefore, $\hat{f}(s) \in \mathfrak{S}(s)$ for ν -a.e. s . For every $A \in \mathcal{Q}_{\mu, \nu}^\Sigma$, the function $\chi_A \hat{f}(s)$, where χ_A is as in the proof of Proposition IX.13, is a product of two $\mu(s)$ -square-integrable functions and, hence, is $\mu(s)$ -integrable for ν -a.e. s . Applying Corollary IX.11 to the function $(s, E) \rightarrow \chi_A(E)f(s, E)$, we conclude that $s \rightarrow \int_A \hat{f}(s|E) d\mu(s|E)$ is a ν -measurable function for any $A \in \mathcal{Q}_{\mu, \nu}^\Sigma$. In view of Proposition IX.13, this means that $\hat{f}$ is a ν -measurable section of $\mathfrak{S}$. By Proposition IX.10, $s \rightarrow \|\hat{f}(s)\|^2$ is a ν -integrable function and we have $\|f\|^2 = \int \|\hat{f}(s)\|^2 d\nu(s)$. This means that $\hat{f} \in \mathfrak{H}$ and the operator Q is isometric. We now prove that Q is unitary. For this, it suffices to show that its image is dense in $\mathfrak{H}$. In other words, we have to show that every $\xi \in \mathfrak{H}$ that is orthogonal to $\hat{f}$ for every $f \in L_2(S_M, M)$ is equal to zero. Given $A \in \mathcal{Q}_{\mu, \nu}^\Sigma$, let $\mathcal{K}_A$ denote the set of all $B \in D_\nu$ such that $B \times A \in D_M$. It is clear that $\mathcal{K}_A$ is closed under finite intersections and $\sigma(\mathcal{K}_A) = \sigma(D_\nu)$. For a set $C \subset S_M$, let X_C denote the function on S_M that is equal to unity on C and vanishes on $S_M \setminus C$. For $A \in \mathcal{Q}_{\mu, \nu}^\Sigma$ and $B \in \mathcal{K}_A$, the function $X_{B \times A}$ is obviously M -square-integrable and, therefore, we have

$$\int_B \langle [\chi_A]_{\mu(s)}, \xi(s) \rangle d\nu(s) = \int \langle \hat{X}_{B \times A}(s), \xi(s) \rangle d\nu(s) = \langle \hat{X}_{B \times A}, \xi \rangle = 0.$$

Applying Lemma IX.12 to $\mathcal{K} = \mathcal{K}_A$, we conclude that $\langle [\chi_A]_{\mu(s)}, \xi(s) \rangle = 0$ for ν -a.e. s for every $A \in \mathcal{Q}_{\mu, \nu}^\Sigma$. Acting as in the proof of Proposition IX.13, we choose a sequence $A_1, A_2, \dots$ of elements of $\mathcal{Q}_{\mu, \nu}^\Sigma$ such that the linear span of $[\chi_{A_i}]_{\mu(s)}$ is dense in $\mathfrak{S}(s)$ for ν -a.e. s . Then, for ν -a.e. s , $\xi(s)$ is orthogonal to all vectors $[\chi_{A_i}]_{\mu(s)}$ and, hence, is equal to zero. Thus, $\xi = 0$ and the unitarity of Q is proved.

3. By Corollary IX.11, $\hat{g}(s)$ is $\mu(s)$ -measurable for ν -a.e. s . Let $C_1 \subset C_2 \subset \dots$ be a sequence of elements of D_M such that $S_M = \bigcup_{j=1}^\infty C_j$ and g is M -essentially bounded on C_j for every $j = 1, 2, \dots$. For $i, j = 1, 2, \dots$, we define the function h_{ij} on S_M by setting $h_{ij}(s, E) = \chi_{A_i}(E)X_{C_j}(s, E)$, where A_i are as in the proof of (2). It is clear that h_{ij} is M -square-integrable and, therefore, $\hat{h}_{ij}$ is a ν -measurable section of $\mathfrak{S}$ for all i, j . To prove the ν -measurability of the family $s \rightarrow \mathcal{T}_{\hat{g}(s)}^{\mu(s)}$, it suffices to show that $\hat{h}_{ij}(s)$ belongs to the domain of $\mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ and the linear span of the vectors $(\hat{h}_{ij}(s), \mathcal{T}_{\hat{g}(s)}^{\mu(s)} \hat{h}_{ij}(s))$ is dense in the graph of $\mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ for ν -a.e. s . For $j = 1, 2, \dots$, let P_j be a ν -a.e. defined map such that $P_j(s) = \mathcal{T}_{\chi_{C_j, s}}^{\mu(s)}$ for ν -a.e. s , where $C_{j, s} = \{E : (s, E) \in C_j\}$. Then, for ν -a.e. s , $P_j(s)$ is an orthogonal projection commuting with $\mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ and satisfying the equality

$$\hat{h}_{ij}(s) = P_j(s)[\chi_{A_i}(s)]_{\mu(s)} \quad (92)$$

for all i, j . In view of Lemma IX.9, $\hat{g}(s)$ is $\mu(s)$ -essentially bounded on $C_{j, s}$ for ν -a.e. s . Hence $\text{Im } P_j(s)$ is contained in the domain of $\mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ and $\mathcal{T}_{\hat{g}(s)}^{\mu(s)} P_j(s)$ is a bounded operator for ν -a.e. s . In particular, it follows from (92) that $\hat{h}_{ij}(s)$ is in the domain of $\mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ for ν -a.e. s . Let $\mathcal{G}_s$ be the subset of the graph of $\mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ consisting of all its elements $(\psi, \tilde{\psi})$ such that $\psi, \tilde{\psi} \in \text{Im } P_j(s)$ for some $j = 1, 2, \dots$. For ν -a.e. s , we have $\lim_{j \rightarrow \infty} P_j(s)\psi = \psi$ for every $\psi \in \mathfrak{S}(s)$. As $P_j(s)$ commute with $\mathcal{T}_{\hat{g}(s)}^{\mu(s)}$, this implies that $\mathcal{G}_s$ is dense in the graph of $\mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ for ν -a.e. s . Given $(\psi, \tilde{\psi}) \in \mathcal{G}_s$ and $\varepsilon > 0$, we can find a finite linear combination τ of vectors $[\chi_{A_i}]_{\mu(s)}$ such that $\|\psi - \tau\| < \varepsilon$. Then we have

$$\begin{aligned} \|\psi - P_j(s)\tau\| &= \|P_j(s)(\psi - \tau)\| < \varepsilon, \\ \|\tilde{\psi} - \mathcal{T}_{\hat{g}(s)}^{\mu(s)} P_j(s)\tau\| &= \|\mathcal{T}_{\hat{g}(s)}^{\mu(s)} P_j(s)(\psi - \tau)\| < \|\mathcal{T}_{\hat{g}(s)}^{\mu(s)} P_j(s)\| \varepsilon, \end{aligned}$$

where j is such that $\psi, \tilde{\psi} \in \text{Im } P_j(s)$. In view of (92), $P_j(s)\tau$ is a linear combination of $\hat{h}_{ij}(s)$, and it follows from the above inequalities that the linear span of the vectors $(\hat{h}_{ij}(s), \mathcal{T}_{\hat{g}(s)}^{\mu(s)} \hat{h}_{ij}(s))$ is dense in $\mathcal{G}_s$ and, hence, in the graph of $\mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ for ν -a.e. s . The ν -measurability of the family $s \rightarrow \mathcal{T}_{\hat{g}(s)}^{\mu(s)}$ is thus proved. Set $T = \int^\oplus \mathcal{T}_{\hat{g}(s)}^{\mu(s)} d\nu(s)$. Let f belong to the domain of $\mathcal{T}_g^M$ and $F = \mathcal{T}_g^M f$. In view of Lemma IX.9, $\hat{F}(s)$ is $\mu(s)$ -equivalent to $\hat{g}(s)\hat{f}(s)$ for ν -a.e. s . This means that $Qf = \hat{F} \in D_T$ and $Q\mathcal{T}_g^M f = TQf$. It follows that T is an extension of $Q\mathcal{T}_g^M Q^{-1}$. To finish the proof, we have to show that $Q^{-1}\xi$ belongs to the domain of $\mathcal{T}_g^M$ for any $\xi \in D_T$. Let $\varphi = gQ^{-1}\xi$. For ν -a.e. s , we have $(T\xi)(s|E) = \hat{g}(s|E)\xi(s|E) = \varphi(s, E)$ for $\mu(s)$ -a.e. E . As $T\xi \in \mathfrak{H}$, it follows that the function $E \rightarrow \varphi(s, E)$ is $\mu(s)$ -square-integrable for ν -a.e. s and the function $s \rightarrow \int |\varphi(s, E)|^2 d\mu(s|E)$ is ν -integrable. In view of Corollary IX.11, this implies that φ is M -square-integrable, i.e., $Q^{-1}\xi$ is in the domain of $\mathcal{T}_g^M$. $\square$

X. EIGENFUNCTION EXPANSIONS

In this section, we prove the results formulated in Sec. II.

Let $\mathcal{B}_{\mathbb{R}}$ denote the Borel σ -algebra on $\mathbb{R}$. Proposition II.4 follows immediately from the next lemma.

Lemma X.1. *Let $\phi \in \mathbb{R}$ and θ be a Borel real function on A^ϕ . Then μ_θ^ϕ is a $(\nu_0, \mathcal{B}_{\mathbb{R}})$ -measurable family of measures and $M = (\mathcal{B}_{\mathbb{R}})\text{-}\int^\oplus \mu_\theta^\phi(s) d\nu_0(s)$ is the unique Borel measure on $S \times \mathbb{R}$ satisfying the conditions of Proposition II.4.*

Proof. Let $\mathcal{K}$ be the set of compact subsets of $\mathbb{R}$. Clearly, $\mathcal{K}$ is contained in the domain of $\mu_\theta^\phi(s)$ for all $s \in S$. Hence, the $(\nu_0, \mathcal{B}_{\mathbb{R}})$ -measurability of μ_θ^ϕ is ensured by Lemma II.3 and Corollary IX.3 (note that $\sigma(\mathcal{K}) = \mathcal{B}_{\mathbb{R}}$).

Let $\mathcal{B}_S$ and $\mathcal{B}_{S \times \mathbb{R}}$ denote the Borel σ -algebras of S and $S \times \mathbb{R}$ respectively. By Definition IX.8, M satisfies the conditions of Proposition IX.7 for $\mu = \mu_\theta^\phi$, $\nu = \nu_0$, $\Sigma = \mathcal{B}_{\mathbb{R}}$, and $\tilde{\Sigma} = \mathcal{B}_S$. By Lemma II.3, we have $K' \times K \in \Delta_{\mu, \nu}^\Sigma$ for any compact sets $K' \subset S$ and $K \subset \mathbb{R}$. Since $\mathcal{B}_{S \times \mathbb{R}} = \sigma(\mathcal{B}_S \boxtimes \mathcal{B}_{\mathbb{R}})$, Proposition IX.7 implies that M is a Borel measure on $S \times \mathbb{R}$ satisfying (31). If f is an M -measurable function, then equality (32) is ensured by Proposition IX.10. Thus, M satisfies the conditions of Proposition II.4.

Suppose $\tilde{M}$ is another Borel measure on S satisfying (31). Let $\mathcal{L}$ be the set of all sets $A \in \mathcal{B}_{S \times \mathbb{R}} \cap D_M \cap D_{\tilde{M}}$ such that $M(A) = \tilde{M}(A)$. Let $\tilde{\mathcal{K}}$ be the set of all sets $K' \times K$, where $K' \subset S$ and $K \subset \mathbb{R}$ are compact sets. Then $\tilde{\mathcal{K}}$ is closed under finite intersections and $\mathcal{L}$ is an α -class containing $\tilde{\mathcal{K}}$. Since $\mathcal{L} \subset \sigma(\tilde{\mathcal{K}}) = \mathcal{B}_{S \times \mathbb{R}}$, Lemma IX.6 implies that $\mathcal{L}$ is a δ -ring. We hence have $M = \tilde{M}$ by Lemma A.3. $\square$

Let $\phi \in \mathbb{R}$ and θ be a Borel real function on A^ϕ . We define the operator-valued map $\mathcal{U}_\theta^\phi$ on S by setting

$$\mathcal{U}_\theta^\phi(s) = \begin{cases} U_{m+\phi}, & s \in S \setminus A^\phi, \\ U_{m+\phi, \theta(s)}, & s \in A^\phi, \end{cases} \quad (93)$$

for all $s = (m, p) \in S$, where the operators U_κ and $U_{\kappa, \vartheta}$ satisfy the conditions of Proposition VI.3 and Proposition VI.4 respectively. By Lemma X.1, the assumptions of Proposition IX.13 are fulfilled for $\Sigma = \mathcal{B}_{\mathbb{R}}$, $\mu = \mu_\theta^\phi$, and $\nu = \nu_0$. A ν_0 -measurable family $\mathfrak{S}$ of Hilbert spaces satisfying the conditions of Proposition IX.13 for such Σ , μ , and ν will be denoted by $\mathfrak{S}_\theta^\phi$.

Lemma X.2. *Let $\phi \in \mathbb{R}$, θ be a Borel real function on A^ϕ , and $\psi \in L_2^c(\mathbb{R}_+)$. Then $s \rightarrow \mathcal{U}_\theta^\phi(s) \psi$ is a ν_0 -measurable section of $\mathfrak{S}_\theta^\phi$.*

Proof. As follows from Proposition IX.13 applied to the set $\mathcal{K}$ of all compact subsets of $\mathbb{R}$, it suffices to show that

$$I_K(s) = \int_K (\mathcal{U}_\theta^\phi(s) \psi)(E) d\mu_\theta^\phi(s|E)$$

is a Borel function on S for any compact set $K \subset \mathbb{R}$. Given $-1 < \kappa < 1$, let the function f_K^κ on $\mathbb{R}$ be defined by the relation

$$f_K^\kappa(\vartheta) = \cos(\vartheta - \vartheta_\kappa) \int_K F_\kappa^{(1)}(E) d\mathcal{V}_{\kappa, \vartheta}(E) + \sin(\vartheta - \vartheta_\kappa) \int_K F_\kappa^{(2)}(E) d\mathcal{V}_{\kappa, \vartheta}(E),$$

where ϑ_κ is defined by (17) and the continuous functions $F_\kappa^{(1)}$ and $F_\kappa^{(2)}$ on $\mathbb{R}$ are given by

$$F_\kappa^{(1)}(E) = \int_0^\infty u^\kappa(E|r) \psi(r) dr, \quad F_\kappa^{(2)}(E) = \int_0^\infty w^\kappa(E|r) \psi(r) dr.$$

It follows from (30), (57), (62), and (93) that $I_K(s) = f_K^{m+\phi}(\theta(s))$ for every $s = (m, p) \in A^\phi$. On the other hand, $p \rightarrow I_K(m, p)$ is a constant function on $\mathbb{R}$ for every m satisfying $|m + \phi| \geq 1$. Since f_K^κ is a Borel function on $\mathbb{R}$ by Lemma II.2, we conclude that I_K is a Borel function on S . $\square$

In what follows, we set $\mathfrak{h} = L_2(\mathbb{R}_+)$.

Corollary X.3. *Let $\phi \in \mathbb{R}$ and θ be a Borel real function on A^ϕ . Then $\mathcal{U}_\theta^\phi$ is a ν_0 -measurable family of operators from $\mathcal{I}_{\mathfrak{h}, \nu_0}$ to $\mathfrak{S}_\theta^\phi$.*

Proof. The statement follows immediately from Lemma X.2 and Lemma B.3. $\square$

Given $\phi \in \mathbb{R}$ and a Borel real function θ on A^ϕ , we define the linear operator U_θ^ϕ from $L_2(S, \mathfrak{h}, \nu_0)$ to $\int^\oplus \mathfrak{S}_\theta^\phi(s) d\nu_0(s)$ by setting

$$U_\theta^\phi = \int^\oplus \mathcal{U}_\theta^\phi(s) d\nu_0(s). \quad (94)$$

Since $\mathcal{U}_\theta^\phi(s)$ is a unitary operator from $\mathfrak{h}$ to $\mathfrak{S}_\theta^\phi(s)$ for ν_0 -a.e. s , it follows from Corollary X.3 and Proposition B.23 that U_θ^ϕ is a unitary operator. If f is an M_θ^ϕ -measurable complex function, we denote by $f|_\theta^\phi$ the ν_0 -equivalence class such that $f|_\theta^\phi(s)$ is the $\mu_\theta^\phi(s)$ -equivalence class of the map $E \rightarrow f(s, E)$ for ν_0 -a.e. s . In view of Lemma X.1, all conditions of Proposition IX.14 are fulfilled for $\Sigma = \mathcal{B}_\mathbb{R}$, $\nu = \nu_0$, $\mu = \mu_\theta^\phi$, $M = M_\theta^\phi$, $\mathfrak{S} = \mathfrak{S}_\theta^\phi$, and $\hat{f} = f|_\theta^\phi$. It follows from statement 2 of Proposition IX.14 that the correspondence $f \rightarrow f|_\theta^\phi$ induces a unitary operator Q_θ^ϕ from $L_2(S \times \mathbb{R}, M_\theta^\phi)$ to $\int^\oplus \mathfrak{S}_\theta^\phi(s) d\nu_0(s)$.

Note that the uniqueness of W in Proposition II.1 is ensured by the density of $L_2^c(\mathbb{R}^3)$ in $L_2(\mathbb{R}^3)$. Hence, Proposition II.1 follows from the next lemma.

Lemma X.4. *Let $\phi \in \mathbb{R}$, θ be a Borel real function on A^ϕ , and V be as in Lemma VIII.1. Then the operator $W = (Q_\theta^\phi)^{-1}U_\theta^\phi V$ satisfies the conditions of Proposition II.1.*

Proof. Let $\Phi \in C_0^\infty(\mathcal{O})$ and $\Psi = [\Phi]_\Lambda$. For $s \in S$ and $E \in \mathbb{R}$, let $h_\Psi(s, E)$ denote the right-hand side of (10). For $-1 < \kappa < 1$ and $m \in \mathbb{Z}$, we define the continuous function F_m^κ on $\mathbb{R} \times \mathbb{R} \times (\mathcal{O})$ by the relation

$$F_m^\kappa(E, \vartheta, x) = \frac{e^{-ipx_3}}{2\pi\sqrt{r_x}} \left(\frac{x_1 - ix_2}{r_x} \right)^m u_\vartheta^\kappa(E|r_x)$$

for all $E, \vartheta \in \mathbb{R}$ and $x = (x_1, x_2, x_3) \in \mathcal{O}$, where $r_x = \sqrt{x_1^2 + x_2^2}$. Set $f_{m, \Psi}^\kappa(E, \vartheta) = \int F_m^\kappa(E, \vartheta, x) \Psi(x) dx$. By (11), (19), and (10), we have $h_\Psi(s, E) = f_{m, \Psi}^{m+\phi}(E, \theta(s))$ for all $s = (m, p) \in A^\phi$ and $E \in \mathbb{R}$. If $|m + \phi| > 1$, then (11), (19), and (10) imply that $(p, E) \rightarrow h_\Psi(m, p; E)$ is a continuous function on $\mathbb{R}^2$. Since $f_{m, \Psi}^\kappa$ is a continuous function on $\mathbb{R}^2$ for all $m \in \mathbb{Z}$ and $-1 < \kappa < 1$, it follows that $(p, E) \rightarrow h_\Psi(m, p; E)$ is a Borel function on $\mathbb{R}^2$ for every $m \in \mathbb{Z}$. Thus, h_Ψ is a Borel (and, hence, M_θ^ϕ -measurable) function on $S \times \mathbb{R}$.

In view of (11), passing to the polar coordinates in the (x_1, x_2) -plane in the integral in (10) yields

$$h_\Psi(s, E) = \int_0^\infty \mathcal{J}_\theta^\phi(s, E|r) \tilde{\Phi}(s|r) dr \quad (95)$$

for all $s \in S$ and $E \in \mathbb{R}$, where the $C_0^\infty(\mathbb{R}_+)$ -valued map $\tilde{\Phi}$ on S is given by (71). It follows from (19), (30), (93), Proposition VI.3, and Proposition VI.4 that, for every $s \in S$ and $\psi \in L_2^c(\mathbb{R}_+)$, the equality

$$(\mathcal{U}_\theta^\phi(s)\psi)(E) = \int_0^\infty \mathcal{J}_\theta^\phi(s, E|r) \psi(r) dr \quad (96)$$

holds for $\mu_\theta^\phi(s)$ -a.e. E . In view of (72), equalities (94), (95) and (96) imply that the function $E \rightarrow h_\Psi(s, E)$ is $\mu_\theta^\phi(s)$ -equivalent to $(U_\theta^\phi V \Psi)(s)$ for ν_0 -a.e. s , whence $U_\theta^\phi V \Psi = h_\Psi|_\theta^\phi$. Since $U_\theta^\phi V \Psi = Q_\theta^\phi W \Psi = (W \Psi)|_\theta^\phi$, it follows from statement 1 of Proposition IX.14 that $W \Psi$ is M_θ^ϕ -equivalent to h_Ψ . Thus, (10) holds for $\Psi = [\Phi]_\Lambda$ with $\Phi \in C_0^\infty(\mathcal{O})$.

For a general $\Psi \in L_2^c(\mathbb{R}^3)$, we choose a sequence $\Phi_n \in C_0^\infty(\mathcal{O})$ such that $\Psi_n = [\Phi_n]_\Lambda$ converge to Ψ in $L_2(\mathbb{R}^3)$ and the supports of all Φ_n are contained in a fixed compact set for all n . Since $\mathcal{W}_\theta^\phi(s, E)$ is locally square-integrable, we have $h_{\Psi_n}(s, E) \rightarrow h_\Psi(s, E)$ for all $s \in S$ and $E \in \mathbb{R}$. As $W \Psi_n \rightarrow W \Psi$ in $L_2(S \times \mathbb{R}, M_\theta^\phi)$ by continuity of W , it follows that $W \Psi$ is M_θ^ϕ -equivalent to h_Ψ . $\square$

Lemma X.4 and the uniqueness statement of Proposition II.1 imply that

$$W_\theta^\phi = (Q_\theta^\phi)^{-1}U_\theta^\phi V. \quad (97)$$

Theorem II.5 follows from Proposition VIII.5 and the next statement.

Proposition X.5. *Let $\phi \in \mathbb{R}$, θ be a Borel real function on A^ϕ , and R_θ^ϕ be defined by (87). Then we have $H_\theta^\phi = R_\theta^\phi$.*

Proof. For brevity, we set $\mu = \mu_\theta^\phi$, $\mathcal{U} = \mathcal{U}_\theta^\phi$, $U = U_\theta^\phi$, $Q = Q_\theta^\phi$, $M = M_\theta^\phi$, and $\mathfrak{g} = \mathfrak{f}|_\theta^\phi$. It follows from (30), (33), (78), (82), (93), and Propositions VI.3 and VI.4 that

$$\mathcal{U}(s)\mathcal{H}_\theta^\phi(s)\mathcal{U}(s)^{-1} = \mathcal{T}_{\mathfrak{g}(s)}^{\mu(s)}$$

for ν_0 -a.e. s . By Lemma VIII.4, $\mathcal{H}_\theta^\phi$ is a ν_0 -measurable family of operators in $\mathfrak{h}$. Hence, it follows from (87), (94), Corollary X.3, statement 1 of Proposition B.23, and Proposition B.25 that $s \rightarrow \mathcal{T}_{\mathfrak{g}(s)}^{\mu(s)}$ is a ν_0 -measurable family of operators in $\mathfrak{S}_\theta^\phi$ and

$$UV R_\theta^\phi (UV)^{-1} = \int^\oplus \mathcal{T}_{\mathfrak{g}(s)}^{\mu(s)} d\nu_0(s).$$

By statement 3 of Proposition IX.14, the operator in the right-hand side is equal to $Q\mathcal{T}_f^M Q^{-1}$. Equalities (34) and (97) therefore imply that $H_\theta^\phi = R_\theta^\phi$. $\square$

ACKNOWLEDGMENTS

The author is grateful to I.V. Tyutin and B.L. Voronov for useful discussions.

Appendix A: Measure theory

In this Appendix, we briefly describe the measure-theoretic concepts used in this paper. Our aim here is not only to fix the notation but also to formulate definitions in such a way as to provide a unified treatment of positive and spectral measures. In particular, in contrast to the standard approach (see, e.g., Refs. 34 and 36), we define positive measures as functions on δ -rings rather than σ -algebras and do not allow them to take infinite values (see Sec. A 2 below).

1. Rings and algebras of sets

Recall that a nonempty set of sets $\mathcal{Q}$ is called a ring of sets if $A \cup B \in \mathcal{Q}$ and $A \setminus B \in \mathcal{Q}$ for any $A, B \in \mathcal{Q}$. As $A \cap B = A \setminus (A \setminus B)$, every ring is closed under finite intersections. A ring $\mathcal{Q}$ is called a σ -ring (a δ -ring) if it is closed under countable unions (resp., under countable intersections). For a set of sets $\mathcal{Q}$, we denote by $\sigma(\mathcal{Q})$ ($\delta(\mathcal{Q})$) the σ -ring (resp., δ -ring) generated by $\mathcal{Q}$, i.e., the smallest σ -ring (resp., δ -ring) containing $\mathcal{Q}$. If $\mathcal{Q}$ is a δ -ring, then $\sigma(\mathcal{Q})$ is just the set of all countable unions of elements of $\mathcal{Q}$. A σ -ring $\mathcal{Q}$ is said to be countably generated if there is a countable set $\tilde{\mathcal{Q}} \subset \mathcal{Q}$ such that $\mathcal{Q} = \sigma(\tilde{\mathcal{Q}})$. A σ -ring $\mathcal{Q}$ is called a σ -algebra if it has the largest element with respect to inclusion.

The σ -ring generated by all open sets of a topological space X is obviously a σ -algebra. It is called the Borel σ -algebra of X and its elements are called Borel subsets of X . A map f is called a Borel map from X to a topological space Y if X is contained in the domain D_f of f , $f(X) \subset Y$, and $f^{-1}(A) \cap X$ is a Borel subset of X for any Borel subset A of Y .

2. Measures

Let $\mathfrak{A}$ be a topological Abelian group and ν be an $\mathfrak{A}$ -valued map. Let $\mathcal{N}_\nu$ denote the set of all $N \in D_\nu$ with the property: if $N' \subset N$ and $N' \in D_\nu$, then $\nu(N') = 0$. We say that ν is an $\mathfrak{A}$ -valued σ -additive function if $\nu(A) = \sum_{i \in I} \nu(A_i)$ for any $A \in D_\nu$ and any countable partition $A = \bigcup_{i \in I} A_i$ with $A_i \in D_\nu$. An $\mathfrak{A}$ -valued σ -additive function is called an $\mathfrak{A}$ -valued measure if its domain D_ν is a δ -ring and the following completeness conditions are satisfied:

- (a) If $A \in \sigma(D_\nu)$ and the family $\{\nu(A_i)\}_{i \in I}$ is summable in $\mathfrak{A}$ for any countable partition $A = \bigcup_{i \in I} A_i$ with $A_i \in D_\nu$, then $A \in D_\nu$.
- (b) If $N \in \mathcal{N}_\nu$ and $N' \subset N$, then $N' \in D_\nu$ (and, hence, $N' \in \mathcal{N}_\nu$).

Elements of $\mathcal{N}_\nu$ are called ν -null sets and elements of $\sigma(D_\nu)$ are called ν -measurable sets. A measure ν is called σ -finite if $\sigma(D_\nu)$ is a σ -algebra. In this case, we denote by S_ν the largest element of $\sigma(D_\nu)$ with respect to inclusion. A measure ν is called finite if it is σ -finite and $S_\nu \in D_\nu$. A measure ν is called positive if it is $\mathbb{R}$ -valued and $\nu(A) \geq 0$ for any $A \in D_\nu$.

Remark A.1. In most expositions of measure theory (see, e.g., Refs 34 and 36), a positive measure is defined as a σ -additive function on a σ -algebra taking values in the extended real semi-axis $\bar{\mathbb{R}}_+ = [0, \infty]$. Such an $\bar{\mathbb{R}}_+$ -valued measure ν is called σ -finite if the largest set in D_ν is a countable union of elements of D_ν with a finite measure. It is called complete if every subset of an element of D_ν with zero measure also belongs to D_ν . Complete σ -finite $\bar{\mathbb{R}}_+$ -valued measures can be naturally identified with positive σ -finite measures in our sense (note that a complete σ -finite $\bar{\mathbb{R}}_+$ -valued measure ν restricted to the set of all elements of D_ν with a finite measure is a positive σ -finite measure in our sense). The advantage of our definition is that it does not involve the extended real axis and, therefore, makes it possible to treat positive measures in the same way as vector-valued measures.

Given a σ -ring $\mathcal{Q}$, a measure ν is called $\mathcal{Q}$ -compatible if $\mathcal{Q} \subset \sigma(D_\nu)$ and for every $A \in D_\nu$, there is an $N \in \mathcal{N}_\nu$ such that $A \cup N \in \mathcal{Q}$. In this case, D_ν consists of all sets of the form $B \cup N$, where $B \in \mathcal{Q} \cap D_\nu$ and $N \in \mathcal{N}_\nu$. If X is a topological space and $\mathcal{B}$ is its Borel σ -algebra, then a $\mathcal{B}$ -compatible measure is called a Borel measure on X .

Remark A.2. Positive Borel measures on a topological space X are usually defined as $\bar{\mathbb{R}}_+$ -valued σ -additive functions on the Borel σ -algebra of X . If such a measure is σ -finite, then its completion corresponds (as described in Remark A.1) to a positive Borel measure on X in our sense. Since we consider only complete measures (see condition (b) above), we introduce the notion of $\mathcal{Q}$ -compatibility to relate measure and topology.

Lemma A.3. *Let $\mathfrak{A}$ be a sequentially complete topological Abelian group and ν be an $\mathfrak{A}$ -valued σ -additive function such that D_ν is a δ -ring. Then there is a unique $\sigma(D_\nu)$ -compatible $\mathfrak{A}$ -valued measure $\hat{\nu}$ such that $D_\nu \subset D_{\hat{\nu}}$ and $\hat{\nu}$ coincides with ν on D_ν . If ν satisfies (a), then $D_\nu = D_{\hat{\nu}} \cap \sigma(D_\nu)$.*

The measure $\hat{\nu}$ is called the completion of the σ -additive function ν .

Proof. We give the proof only for positive ν because the lemma is used in this paper only in this case. The existence and uniqueness of $\hat{\nu}$ is then guaranteed by the well-known results on the extension of positive measures. Let ν satisfy (a) and $A \in D_{\hat{\nu}} \cap \sigma(D_\nu)$. Let $\{A_i\}_{i \in I}$ be a countable partition of A such that $A_i \in D_\nu$ for all $i \in I$. Since $\hat{\nu}(A_i) = \nu(A_i)$ for all $i \in I$, the σ -additivity of $\hat{\nu}$ implies that the family $\{\nu(A_i)\}_{i \in I}$ is summable. Condition (a) hence ensures that $A \in D_\nu$ and, therefore, $D_\nu \supset D_{\hat{\nu}} \cap \sigma(D_\nu)$. As the opposite inclusion also obviously holds, the lemma is proved. $\square$

Let ν be an $\mathfrak{A}$ -valued measure and A be a ν -measurable set. The restriction $\nu|_A$ of ν to A is, by definition, the restriction of the map ν to the domain $D_{\nu|_A}$ consisting of all elements of D_ν that are contained in A . Clearly, $\nu|_A$ is a σ -finite measure for any ν -measurable set A and $S_{\nu|_A} = A$.

Let ν_1 and ν_2 be positive measures and $\mathcal{Q}$ be the σ -ring generated by all sets of the form $A_1 \times A_2$, where $A_1 \in D_{\nu_1}$ and $A_2 \in D_{\nu_2}$. Then there is a unique positive $\mathcal{Q}$ -compatible measure ν such that $\nu(A_1 \times A_2) = \nu_1(A_1)\nu_2(A_2)$ for every $A_1 \in D_{\nu_1}$ and $A_2 \in D_{\nu_2}$. This measure is called the product of ν_1 and ν_2 and is denoted by $\nu_1 \times \nu_2$.

3. Standard measures

Let $\mathfrak{A}$ be a topological Abelian group. An $\mathfrak{A}$ -valued measure ν is called standard if it is σ -finite and there exists a complete separable metric space X such that $X \subset S_\nu$, $S_\nu \setminus X$ is a ν -null set, and $\nu|_X$ is a Borel measure on X . If ν is a standard measure and A is a ν -measurable set, then $\nu|_A$ is also a standard measure.

Remark A.4. Standard measures were first introduced in Ref. 37. Unlike Ref. 37, we consider only complete measures, and standard measures in our sense are actually the completions of those in the sense of Ref. 37. In the probabilistic context, such complete measures were introduced in Ref. 38 under the name of Lebesgue measure spaces. The class of standard measures is broad enough to include most measures encountered in applications and, at the same time, is narrow enough to exclude measures with a pathological behavior.

4. Measurable maps

Let ν be a σ -finite $\mathfrak{A}$ -valued measure. A map f is said to be defined ν -a.e. if $S_\nu \setminus D_f$ is a ν -null set. Given a set X , a map f is said to be an X -valued ν -a.e. defined map if $S_\nu \setminus f^{-1}(X)$ is a ν -null set. If X is a topological space, then a map f is called an X -valued ν -measurable map if f is an X -valued ν -a.e.

defined map and $f^{-1}(B) \cap S_\nu$ is a ν -measurable set for any Borel subset B of X . We say that some property $P(s)$ holds for ν -a.e. s , if there is a ν -null set N such that $P(s)$ holds for all $s \in S_\nu \setminus N$.

Given a ν -measurable set A , we say that f is an X -valued ν -measurable map on A if f is an X -valued $\nu|_A$ -measurable map. A property $P(s)$ is said to hold for ν -a.e. $s \in A$ if it holds for $\nu|_A$ -a.e. s .

A complex ν -a.e. defined function f is said to be ν -essentially bounded if there is $C > 0$ such that $|f(s)| \leq C$ for ν -a.e. s .

Remark A.5. Note that the domain D_f of a ν -a.e. defined or ν -measurable map f is not assumed to be contained in S_ν . Moreover, X -valued ν -a.e. defined (and, in particular, ν -measurable) maps are allowed to take values outside X on some ν -null set contained in $S_\nu \cap D_f$. This implies in particular, that using “the set of all X -valued ν -measurable maps” would lead to the same kind of set-theoretic problems as the use of “the set of all sets”.

Let X be a complete separable metric space, f be an X -valued ν -a.e. defined map, and $f_1, f_2, \dots$ be a sequence of X -valued ν -measurable maps such that $f_n(s)$ converge to $f(s)$ in X as $n \rightarrow \infty$ for ν -a.e. s . Then f is ν -measurable.

5. Equivalence classes

Let ν be a σ -finite $\mathfrak{A}$ -valued measure. Two ν -a.e. defined maps f and g are called ν -equivalent if $f(s) = g(s)$ for ν -a.e. s . All ν -a.e. defined maps fall into disjoint classes of ν -equivalent maps, which are called ν -equivalence classes. In every ν -equivalence class, we choose an arbitrary fixed element whose domain of definition is contained in S_ν . Given a ν -a.e. defined map f , we denote by $[f]_\nu$ such a chosen element belonging to the ν -equivalence class containing f . Thus, the map $[f]_\nu$ is ν -equivalent to f for any ν -a.e. defined map f , and we have $[f]_\nu = [g]_\nu$ for every pair of ν -equivalent maps f and g . If X is a topological space, then f is an X -valued ν -measurable map if and only if so is $[f]_\nu$.

In this paper, ν -equivalence classes *per se* are not used. Whenever we speak of ν -equivalence classes, we always refer to representatives of the form $[f]_\nu$, where f is a ν -a.e. defined map. There are two reasons for such a redefinition of the notion of a ν -equivalence class. First, the same arguments as in Remark A.5 show that the ν -equivalence classes in the true sense cannot be considered as well-defined sets. Hence, using them as elements of sets is not satisfactory from the viewpoint of foundations of mathematics. Choosing a fixed representative in each class allows us to circumvent this difficulty. Second, since ν -equivalence classes in our sense are just some maps, all definitions and notations introduced for maps become directly applicable to ν -equivalence classes.

6. Integrable functions

Let ν be a σ -finite positive measure. A ν -measurable complex function f is called ν -integrable if

$$I_\nu(f) = \sup_{A_i \in D_\nu, c_i \in \mathbb{R}} \sum_{i=1}^n c_i \nu(A_i) < \infty,$$

where the supremum is taken over all finite sets $A_1, \dots, A_n$ of disjoint elements of D_ν and $c_1, \dots, c_n \in \mathbb{R}$ such that $c_i \leq |f(s)|$ for ν -a.e. $s \in A_i$. The integral $\int f(s) d\nu(s)$ of a ν -integrable function f is a complex number that is uniquely determined by the conditions that $\int f(s) d\nu(s)$ be linear in f and coincide with $I_\nu(f)$ if $f(s) \geq 0$ for ν -a.e. s .

Clearly, passing to a ν -equivalent function does not affect its ν -integrability. In particular, a ν -a.e. defined function f is ν -integrable if and only if $[f]_\nu$ is ν -integrable. If A is a ν -measurable set and f is a $\nu|_A$ -integrable function, then we say that f is a ν -integrable function on A . In this case, we write $\int_A f(s) d\nu(s)$ in place of $\int f(s) d\nu|_A(s)$.

Remark A.6. As for ν -measurable functions, the domain of a ν -integrable function is not assumed to be contained in S_ν .

7. L_2 -spaces

Let ν be a σ -finite positive measure and $\mathfrak{h}$ be a separable Hilbert space.³⁹ We denote by $\mathcal{M}(\mathfrak{h}, \nu)$ the set of all elements $[f]_\nu$, where f is an $\mathfrak{h}$ -valued ν -measurable map. The set $\mathcal{M}(\mathfrak{h}, \nu)$ has a natural structure of a vector space over $\mathbb{C}$ (for any $\mathfrak{h}$ -valued ν -a.e. defined maps f and g and any $k \in \mathbb{C}$, we set

$[f]_\nu + [g]_\nu = [f + g]_\nu$ and $k[f]_\nu = [kf]_\nu$. Given a ν -measurable set A , we denote by $L_2(A, \mathfrak{h}, \nu)$ the linear subspace of $\mathcal{M}(\mathfrak{h}, \nu|_A)$ consisting of all its elements f such that $s \rightarrow \|f(s)\|^2$ is a ν -integrable function. Defining the scalar product of $f, g \in L_2(A, \mathfrak{h}, \nu)$ by the formula

$$\langle f, g \rangle = \int_A \langle f(s), g(s) \rangle d\nu(s),$$

we make $L_2(A, \mathfrak{h}, \nu)$ into a Hilbert space.

For $\mathfrak{h} = \mathbb{C}$, we denote the space $L_2(A, \mathfrak{h}, \nu)$ by $L_2(A, \nu)$. If $A \subset \mathbb{R}^n$ and ν is the Lebesgue measure on $\mathbb{R}^n$, the space $L_2(A, \nu)$ is denoted by $L_2(A)$.

Given a ν -measurable complex function g , we denote by $\mathcal{T}_g^\nu$ the operator of multiplication by g in $L_2(S_\nu, \nu)$. By definition, the graph of $\mathcal{T}_g^\nu$ consists of all pairs (f_1, f_2) of elements of $L_2(S_\nu, \nu)$ such that $f_2(s) = g(s)f_1(s)$ for ν -a.e. s . The operator $\mathcal{T}_g^\nu$ is closed and densely defined and its adjoint is equal to $\mathcal{T}_{\bar{g}}^\nu$, where $\bar{g}$ is the complex conjugate function of g . In particular, if g is real, then $\mathcal{T}_g^\nu$ is self-adjoint.

8. Spectral measures

We refer the reader to Chapter 5 of Ref. 40 for a detailed exposition of the theory of spectral measures. Here, we only give a brief summary of the facts needed in this paper. Let $\mathfrak{H}$ be a separable Hilbert space and $L(\mathfrak{H})$ be the space of bounded everywhere defined linear operators in $\mathfrak{H}$ endowed with the strong operator topology. A finite $L(\mathfrak{H})$ -valued measure $\mathcal{E}$ is called a spectral measure in $\mathfrak{H}$ if $\mathcal{E}(A)$ is an orthogonal projection in $\mathfrak{H}$ for any $A \in D_{\mathcal{E}}$ and $\mathcal{E}(S_{\mathcal{E}})$ is the identity operator in $\mathfrak{H}$. For any $\Psi \in \mathfrak{H}$, we define the positive measure $\mathcal{E}_\Psi$ as the completion of the positive σ -additive function $A \rightarrow \langle \mathcal{E}(A)\Psi, \Psi \rangle$ on $D_{\mathcal{E}}$, where $\langle \cdot, \cdot \rangle$ is the scalar product on $\mathfrak{H}$.

Let $\mathcal{E}$ be a spectral measure. Given an $\mathcal{E}$ -measurable complex function f , the integral $J_f^\mathcal{E}$ of f with respect to $\mathcal{E}$ is defined as the unique linear operator in $\mathfrak{H}$ such that

$$D_{J_f^\mathcal{E}} = \left\{ \Psi \in \mathfrak{H} : \int |f(s)|^2 d\mathcal{E}_\Psi(s) < \infty \right\} \quad (\text{A1})$$

$$\langle \Psi, J_f^\mathcal{E} \Psi \rangle = \int f(s) d\mathcal{E}_\Psi(s), \quad \Psi \in D_{J_f^\mathcal{E}}. \quad (\text{A2})$$

For any $\mathcal{E}$ -measurable complex function f , the operator $J_f^\mathcal{E}$ is closed and densely defined and its adjoint is equal to $J_{\bar{f}}^\mathcal{E}$.

For every normal operator T , there is a unique Borel spectral measure $\mathcal{E}_T$ on $\mathbb{C}$ such that $J_g^{\mathcal{E}_T} = T$, where g is the identity function on $\mathbb{C}$. The operators $\mathcal{E}_T(A)$, where A is a Borel subset of $\mathbb{C}$, are called the spectral projections of T . If f is an $\mathcal{E}_T$ -measurable complex function, then the operator $J_f^{\mathcal{E}_T}$ is also denoted as $f(T)$.

Appendix B: Direct integral decompositions of operators in Hilbert space

One of the main mathematical tools used in this paper are the direct integral decompositions of operators in Hilbert space. In its original form⁶, the theory of such decompositions (also known as von Neumann's reduction theory) is applicable only to bounded operators⁴¹ and, therefore, is insufficient for quantum-mechanical applications, where unbounded operators play a prominent role. An extension of the reduction theory to unbounded operators was given in Ref. 7 (most results of Ref. 7 were earlier formulated without proofs in Ref. 8). The approach proposed in Ref. 7 is based on the observation⁴² that the properties of a closed operator in a Hilbert space can be encoded in a set of four bounded operators forming its so called characteristic matrix. This allows one to reformulate problems concerning the direct integrals of arbitrary closed operators in terms of direct integrals of bounded operators. Although this approach makes it possible to extend the results of the original von Neumann's theory to unbounded operators, it seems to be somewhat inconvenient for applications to concrete problems, where the computation of the characteristic matrices may turn out to be a difficult task. In particular, the definition of the measurability of families of operators given in Ref. 7 may be difficult to verify for concrete examples. From the viewpoint of applications, the definition of measurability proposed in Ref. 8 and formulated directly in terms of the graphs of operators seems to be more suitable (both definitions are actually equivalent, see Remark B.12 below).

In this appendix, we give a self-contained exposition of the theory of direct integral decompositions of (generally, unbounded) operators in Hilbert space. Our treatment is based on the formulation of

measurability proposed in Ref. 8 and does not involve characteristic matrices. Such an approach allows us to derive all results in a straightforward manner: unlike Ref. 7, we do not need the theory for bounded operators as a prerequisite. Moreover, it becomes possible to give a concise treatment of direct integral decompositions of sums and products of operators (see Propositions B.25 and B.26 below), whose original analysis in Ref. 43 involves lengthy computations of characteristic matrices.

In what follows, we use the measure-theoretic framework described in Appendix A. Throughout this appendix, ν denotes a σ -finite positive measure.

1. Measurable families of Hilbert spaces

A ν -a.e. defined map $\mathfrak{S}$ is called a ν -a.e. defined family of Hilbert spaces if $\mathfrak{S}(s)$ is a separable Hilbert space for ν -a.e. s . A map ξ is called a ν -a.e. defined section of $\mathfrak{S}$ if $\xi(s) \in \mathfrak{S}(s)$ for ν -a.e. s .

A ν -a.e. defined family of Hilbert spaces $\mathfrak{S}$ is called ν -measurable if it is equipped with a sequence $\{\xi_i^\mathfrak{S}\}_{i=1}^\infty$ of ν -a.e. defined sections of $\mathfrak{S}$ such that the linear span of the vectors $\xi_1^\mathfrak{S}(s), \xi_2^\mathfrak{S}(s), \dots$ is dense in $\mathfrak{S}(s)$ for ν -a.e. s , and $s \rightarrow \langle \xi_i^\mathfrak{S}(s), \xi_j^\mathfrak{S}(s) \rangle$ is a ν -measurable complex function for any $i, j = 1, 2, \dots$. A ν -a.e. defined section ξ of $\mathfrak{S}$ is called ν -measurable if $s \rightarrow \langle \xi_i^\mathfrak{S}(s), \xi(s) \rangle$ is a ν -measurable complex function for any $i = 1, 2, \dots$.

We say that ν -measurable families $\mathfrak{S}_1$ and $\mathfrak{S}_2$ of Hilbert spaces are ν -identical if they are ν -equivalent and have the same ν -measurable sections.

A sequence $e_1, e_2, \dots$ of elements of a Hilbert space $\mathfrak{H}$ is said to be a generalized orthonormal system in $\mathfrak{H}$ if $\langle e_i, e_j \rangle = 0$ for $i \neq j$ and $\|e_i\|$ is equal to either 1 or 0 for any $i = 1, 2, \dots$.

Let $\mathfrak{S}$ be a ν -a.e. defined family of Hilbert spaces and $\xi_1, \xi_2, \dots$ be a sequence of ν -a.e. defined sections of $\mathfrak{S}$. We say that ν -a.e. defined sections $\tilde{\xi}_1, \tilde{\xi}_2, \dots$ of $\mathfrak{S}$ constitute an orthonormal sequence associated with $\xi_1, \xi_2, \dots$ if the following conditions hold

- (a) For ν -a.e. s , the sequence $\tilde{\xi}_1(s), \tilde{\xi}_2(s), \dots$ is a generalized orthonormal system in $\mathfrak{S}(s)$ whose linear span coincides with that of $\xi_1(s), \xi_2(s), \dots$.
- (b) There exist ν -measurable complex functions f_{ij} defined for $i \leq j$ such that $\tilde{\xi}_j(s) = \sum_{i=1}^j f_{ij}(s) \xi_i(s)$ for all $j = 1, 2, \dots$ and ν -a.e. s .

Lemma B.1. *Let $\mathfrak{S}$ be a ν -a.e. defined family of Hilbert spaces and $\xi_1, \xi_2, \dots$ be a sequence of ν -a.e. defined sections of $\mathfrak{S}$ such that $s \rightarrow \langle \xi_i(s), \xi_j(s) \rangle$ is a ν -measurable complex function for any $i, j = 1, 2, \dots$. Then there exists an orthonormal sequence associated with $\xi_1, \xi_2, \dots$.*

Proof. The required sequence is constructed by applying the standard orthogonalization procedure to $\xi_1(s), \xi_2(s), \dots$, see the proof of Lemma 1 in Sec. II.1.2 of Ref. 19 for details. $\square$

If ξ and η are ν -measurable sections of a ν -measurable family $\mathfrak{S}$ of Hilbert spaces, then $s \rightarrow \langle \xi(s), \eta(s) \rangle$ is a ν -measurable complex function. Indeed, by Lemma B.1, there exists an orthonormal sequence $\tilde{\xi}_1, \tilde{\xi}_2, \dots$ associated with $\xi_1^\mathfrak{S}, \xi_2^\mathfrak{S}, \dots$. By (a), the linear span of $\tilde{\xi}_1(s), \tilde{\xi}_2(s), \dots$ is dense in $\mathfrak{S}(s)$ for ν -a.e. s . This implies that $\langle \xi(s), \eta(s) \rangle = \sum_{i=1}^\infty \langle \xi(s), \tilde{\xi}_i(s) \rangle \langle \tilde{\xi}_i(s), \eta(s) \rangle$ for ν -a.e. s . Hence the required statement follows because $s \rightarrow \langle \xi(s), \tilde{\xi}_i(s) \rangle$ and $s \rightarrow \langle \tilde{\xi}_i(s), \eta(s) \rangle$ are ν -measurable complex functions by (b) and the ν -measurability of ξ and η .

Lemma B.2. *Let $\mathfrak{S}$ be a ν -measurable family of Hilbert spaces and $\xi_1, \xi_2, \dots$ be ν -measurable sections of $\mathfrak{S}$. Let ξ be a ν -a.e. defined map such that $\xi(s)$ belongs to the closed linear span of $\xi_1(s), \xi_2(s), \dots$ for ν -a.e. s . If $s \rightarrow \langle \xi_j(s), \xi(s) \rangle$ is a ν -measurable function for all $j = 1, 2, \dots$, then ξ is a ν -measurable section of $\mathfrak{S}$.*

Proof. By Lemma B.1, there exists an orthonormal sequence $\tilde{\xi}_1, \tilde{\xi}_2, \dots$ associated with $\xi_1, \xi_2, \dots$. By condition (b), $\tilde{\xi}_j$ is a ν -measurable section of $\mathfrak{S}$ for every $j = 1, 2, \dots$. By condition (a), $\xi_1(s), \xi_2(s), \dots$ is a generalized orthonormal system whose closed linear span coincides with that of $\tilde{\xi}_1(s), \tilde{\xi}_2(s), \dots$ and, hence, contains $\xi(s)$ for ν -a.e. s . This implies that $\xi(s) = \sum_{j=1}^\infty \langle \tilde{\xi}_j(s), \xi(s) \rangle \tilde{\xi}_j(s)$ for ν -a.e. s and, therefore, ξ is a ν -measurable section of $\mathfrak{S}$. $\square$

2. Direct sums of measurable families

Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces. Then there is a unique (up to ν -identity) ν -measurable family $\mathfrak{S}$ of Hilbert spaces satisfying the conditions

(1) $\mathfrak{S}(s) = \mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s)$ for ν -a.e. s .

(2) A ν -a.e. defined map ζ is a ν -measurable section of $\mathfrak{S}$ if and only if there are ν -measurable sections ξ and η of $\mathfrak{S}_1$ and $\mathfrak{S}_2$ respectively such that $\zeta(s) = (\xi(s), \eta(s))$ for ν -a.e. s .

Indeed, every ν -a.e. defined map $\mathfrak{S}$ satisfying (1) becomes a ν -measurable family of Hilbert spaces satisfying (2) if we choose $\xi_1^\mathfrak{S}, \xi_2^\mathfrak{S}, \dots$ in such a way that $\xi_{2k}^\mathfrak{S}(s) = (\xi_k^{\mathfrak{S}_1}(s), 0)$ and $\xi_{2k-1}^\mathfrak{S}(s) = (0, \xi_k^{\mathfrak{S}_2}(s))$ for ν -a.e. s and every $k = 1, 2, \dots$. The ν -measurable family $\mathfrak{S}$ of Hilbert spaces satisfying (1) and (2) is called the direct sum of $\mathfrak{S}_1$ and $\mathfrak{S}_2$ and is denoted by $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$.

3. Constant and discrete families

Let $\mathfrak{h}$ be a separable Hilbert space. Then there exists a unique (up to ν -identity) ν -measurable family $\mathfrak{S}$ of Hilbert spaces satisfying the conditions

(1) $\mathfrak{S}(s) = \mathfrak{h}$ for ν -a.e. s .

(2) A ν -a.e. defined map ξ is a ν -measurable section of $\mathfrak{S}$ if and only if ξ is an $\mathfrak{h}$ -valued ν -measurable map.

Indeed, let $e_1, e_2, \dots$ be an orthonormal basis in $\mathfrak{h}$ and $\mathfrak{S}$ be a ν -a.e. defined map satisfying (1). To make $\mathfrak{S}$ into a ν -measurable family, we require $\xi_1^\mathfrak{S}, \xi_2^\mathfrak{S}, \dots$ to be ν -a.e. defined maps such that $\xi_i^\mathfrak{S}(s) = e_i$ for ν -a.e. s . If ξ is a ν -measurable section of $\mathfrak{S}$, then ξ is an $\mathfrak{h}$ -valued ν -measurable map because $\xi(s) = \sum_i \langle \xi_i^\mathfrak{S}(s), \xi(s) \rangle e_i$ for ν -a.e. s and, therefore, $\mathfrak{S}$ satisfies (2). The ν -measurable family $\mathfrak{S}$ of Hilbert spaces satisfying (1) and (2) is denoted by $\mathcal{I}_{\mathfrak{h}, \nu}$.

Suppose now that $\mathcal{S}$ is a countable set, ν is a counting measure on $\mathcal{S}$ and $\mathfrak{S}$ is a ν -a.e. defined family of Hilbert spaces (i.e., $\mathcal{S} \subset D_\mathfrak{S}$ and $\mathfrak{S}(s)$ is a separable Hilbert space for every $s \in \mathcal{S}$). Since every ν -a.e. defined complex function is ν -measurable, there is a unique (up to ν -identity) ν -measurable family $\mathfrak{S}_\mathcal{S}$ of Hilbert spaces such that $\mathfrak{S}_\mathcal{S}(s) = \mathfrak{S}(s)$ for every $s \in \mathcal{S}$. Every ν -a.e. defined section of $\mathfrak{S}_\mathcal{S}$ is ν -measurable.

4. Measurable families of operators

Let $\mathfrak{S}$ be a ν -measurable family of Hilbert spaces. A ν -a.e. defined map $\mathfrak{S}'$ is said to be a ν -a.e. defined family of subspaces of $\mathfrak{S}$ if $\mathfrak{S}'(s)$ is a linear (not necessarily closed) subspace of $\mathfrak{S}(s)$ for ν -a.e. s . A ν -a.e. defined family $\mathfrak{S}'$ of subspaces of $\mathfrak{S}$ is called ν -measurable if there is a sequence $\xi_1, \xi_2, \dots$ of ν -measurable sections of $\mathfrak{S}$ such that the linear span of $\xi_1(s), \xi_2(s), \dots$ is dense in $\mathfrak{S}'(s)$ for ν -a.e. s (such a sequence will be called a ν -measurable basis in $\mathfrak{S}'$).

Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces. A ν -a.e. defined map $\mathcal{R}$ is called a ν -a.e. defined family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ if $\mathcal{R}(s)$ is an operator (possibly not everywhere defined and unbounded) from $\mathfrak{S}_1(s)$ to $\mathfrak{S}_2(s)$ for ν -a.e. s . A ν -a.e. defined family $\mathcal{R}$ of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ is called ν -measurable if there are sequences $\xi_1, \xi_2, \dots$ and $\eta_1, \eta_2, \dots$ of ν -measurable sections of $\mathfrak{S}_1$ and $\mathfrak{S}_2$ respectively such that the linear span of the vectors $(\xi_j(s), \eta_j(s))$ is dense in the graph $G_{\mathcal{R}(s)}$ of the operator $\mathcal{R}(s)$ for ν -a.e. s . In other words, $\mathcal{R}$ is ν -measurable if $s \rightarrow G_{\mathcal{R}(s)}$ is a ν -measurable family of subspaces of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$.

Given a ν -measurable family $\mathfrak{S}$ of Hilbert spaces, we say that $\mathcal{R}$ is a ν -a.e. defined (ν -measurable) family of operators in $\mathfrak{S}$ if it is a ν -a.e. defined (resp., ν -measurable) family of operators from $\mathfrak{S}$ to $\mathfrak{S}$. If $\mathfrak{h}$ is a separable Hilbert space, then ν -a.e. defined (ν -measurable) families of operators in $\mathcal{I}_{\mathfrak{h}, \nu}$ are called ν -a.e. defined (resp., ν -measurable) families of operators in $\mathfrak{h}$.

Lemma B.3. *Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces and $\mathcal{R}$ be a ν -a.e. defined family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ such that $\mathcal{R}(s)$ is an everywhere defined bounded operator for ν -a.e. s . Then the following statements hold*

1. *If $\mathcal{R}$ is ν -measurable then $s \rightarrow \mathcal{R}(s)\xi(s)$ is a ν -measurable section of $\mathfrak{S}_2$ for any ν -measurable section ξ of $\mathfrak{S}_1$.*
2. *Suppose there exists a ν -measurable basis $\xi_1, \xi_2, \dots$ in $\mathfrak{S}_1$ such that $s \rightarrow \mathcal{R}(s)\xi_j(s)$ is a ν -measurable section of $\mathfrak{S}_2$ for all $j = 1, 2, \dots$. Then $\mathcal{R}$ is ν -measurable.*

Proof. 1. Let ξ be a ν -measurable section of $\mathfrak{S}_1$ and $\xi_1, \xi_2, \dots$ and $\eta_1, \eta_2, \dots$ be ν -measurable sections of $\mathfrak{S}_1$ and $\mathfrak{S}_2$ respectively such that the linear span of the vectors $(\xi_j(s), \eta_j(s))$ is dense in $G_{\mathcal{R}(s)}$ for ν -a.e. s . By Lemma B.1, there exists an orthonormal sequence $\tilde{\xi}_1, \tilde{\xi}_2, \dots$ associated with $\xi_1, \xi_2, \dots$. By condition (b) of Sec. B 1, we have $\mathcal{R}(s)\tilde{\xi}_j(s) = \sum_{i=1}^j f_{ij}(s)\eta_i(s)$ with some ν -measurable functions f_{ij} for any $j = 1, 2, \dots$ and ν -a.e. s . Hence, $s \rightarrow \mathcal{R}(s)\tilde{\xi}_j(s)$ is a ν -measurable section of $\mathfrak{S}_2$ for any $j = 1, 2, \dots$. By condition (a) of Sec. B 1, the linear span of $\xi_1(s), \tilde{\xi}_2(s), \dots$ coincides with that of $\xi_1(s), \xi_2(s), \dots$ and, therefore, is dense in $\mathfrak{S}_1(s)$ for ν -a.e. s . This implies that $\xi(s) = \sum_{j=1}^{\infty} \langle \tilde{\xi}_j(s), \xi(s) \rangle \tilde{\xi}_j(s)$ for ν -a.e. s . By the continuity of $\mathcal{R}(s)$, it follows that $\mathcal{R}(s)\xi(s) = \sum_{j=1}^{\infty} \langle \tilde{\xi}_j(s), \xi(s) \rangle \mathcal{R}(s)\tilde{\xi}_j(s)$ for ν -a.e. s and, therefore, $s \rightarrow \mathcal{R}(s)\xi(s)$ is a ν -measurable section of $\mathfrak{S}_2$.

2. Let ν -measurable sections $\eta_1, \eta_2, \dots$ of $\mathfrak{S}_2$ be such that $\eta_j(s) = \mathcal{R}(s)\xi_j(s)$ for all $j = 1, 2, \dots$ and ν -a.e. s . As the linear span of $\xi_1(s), \xi_2(s), \dots$ is dense in $\mathfrak{S}_1(s)$ and $\mathcal{R}(s)$ is everywhere defined and continuous, the linear span of $(\xi_j(s), \eta_j(s))$ is dense in $G_{\mathcal{R}(s)}$ for ν -a.e. s . This means that $\mathcal{R}$ is ν -measurable. $\square$

Lemma B.4. *Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces and $\mathcal{R}$ be a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ such that $\mathcal{R}(s)$ is an everywhere defined bounded operator for ν -a.e. s . Let $\mathfrak{S}'_1$ be a ν -measurable family of subspaces of $\mathfrak{S}_1$. Then the images of $\mathfrak{S}'_1(s)$ under $\mathcal{R}(s)$ constitute a ν -measurable family of subspaces of $\mathfrak{S}_2$.*

Proof. Let $\xi_1, \xi_2, \dots$ be a ν -measurable basis in $\mathfrak{S}'_1$. By the continuity of $\mathcal{R}(s)$, the linear span of $\mathcal{R}(s)\xi_1(s), \mathcal{R}(s)\xi_2(s), \dots$ is dense in the image of $\mathfrak{S}'_1(s)$ under $\mathcal{R}(s)$ for ν -a.e. s . Hence the statement follows because $s \rightarrow \mathcal{R}(s)\xi_j(s)$ is a ν -measurable section of $\mathfrak{S}_2$ for any $j = 1, 2, \dots$ by Lemma B.3. $\square$

Lemma B.5. *Let $\mathfrak{S}$ be a ν -measurable family of Hilbert spaces and $\mathfrak{S}'$ and $\mathcal{P}$ be ν -a.e. defined maps such that $\mathfrak{S}'(s)$ is a closed subspace of $\mathfrak{S}(s)$ and $\mathcal{P}(s)$ is the orthogonal projection of $\mathfrak{S}(s)$ onto $\mathfrak{S}'(s)$ for ν -a.e. s . Then $\mathcal{P}$ is a ν -measurable family of operators in $\mathfrak{S}$ if and only if $\mathfrak{S}'$ is a ν -measurable family of subspaces of $\mathfrak{S}$.*

Proof. If $\mathcal{P}$ is ν -measurable, then so is $\mathfrak{S}'$ by Lemma B.4. Let $\mathfrak{S}'$ be ν -measurable and $\xi_1, \xi_2, \dots$ be a ν -measurable basis in $\mathfrak{S}'$. Let ξ be a ν -measurable section of $\mathfrak{S}$. For ν -a.e. s , we have $\langle \mathcal{P}(s)\xi(s), \xi_j(s) \rangle = \langle \xi(s), \xi_j(s) \rangle$. Hence, $s \rightarrow \langle \mathcal{P}(s)\xi(s), \xi_j(s) \rangle$ is a ν -measurable section of $\mathfrak{S}$ for all $j = 1, 2, \dots$. In view of Lemma B.2, this implies that $s \rightarrow \mathcal{P}(s)\xi(s)$ is a ν -measurable section of $\mathfrak{S}$ and, therefore, $\mathcal{P}$ is ν -measurable by Lemma B.3. $\square$

Let $\mathfrak{H}'$ and $\mathfrak{H}''$ be closed subspaces of a Hilbert space $\mathfrak{H}$ and P' and P'' be the orthogonal projections of $\mathfrak{H}$ onto $\mathfrak{H}'$ and $\mathfrak{H}''$ respectively. Then we have $\mathfrak{H}' \cap \mathfrak{H}'' = \text{Ker } R$, where $R = 1_{\mathfrak{H}} - (P' + P'')/2$ and $1_{\mathfrak{H}}$ denotes the identity operator in $\mathfrak{H}$. Indeed, suppose $\psi \in \text{Ker } R$ and $\psi \notin \mathfrak{H}' \cap \mathfrak{H}''$. Assume, for definiteness, that $\psi \notin \mathfrak{H}'$. Then $\|P'\psi\| < \|\psi\|$. Since $\psi = (P'\psi + P''\psi)/2$ and $\|P''\psi\| \leq \|\psi\|$, this implies that $\|\psi\| < \|\psi\|$. We thus obtain a contradiction and the statement is proved.

Lemma B.6. *Let $\mathfrak{S}$ be a ν -measurable family of Hilbert spaces and $\mathfrak{S}'$ and $\mathfrak{S}''$ be ν -measurable families of closed subspaces of $\mathfrak{S}$. Then*

1. *If $\mathfrak{S}''(s) \subset \mathfrak{S}'(s)$ for ν -a.e. s , then $s \rightarrow \mathfrak{S}'(s) \ominus \mathfrak{S}''(s)$, where $\mathfrak{S}'(s) \ominus \mathfrak{S}''(s)$ is the orthogonal complement of $\mathfrak{S}''(s)$ in $\mathfrak{S}'(s)$, is a ν -measurable family of subspaces of $\mathfrak{S}$.*
2. *$s \rightarrow \mathfrak{S}'(s) \cap \mathfrak{S}''(s)$ is a ν -measurable family of subspaces of $\mathfrak{S}$.*

Proof. 1. Let $\mathcal{P}'$ and $\mathcal{P}''$ be ν -a.e. defined maps such that $\mathcal{P}'(s)$ and $\mathcal{P}''(s)$ are orthogonal projections of $\mathfrak{S}(s)$ onto $\mathfrak{S}'(s)$ and $\mathfrak{S}''(s)$ respectively for ν -a.e. s . By Lemma B.5, $\mathcal{P}'$ and $\mathcal{P}''$ are ν -measurable families of operators in $\mathfrak{S}$. It follows from Lemma B.3 that $s \rightarrow \mathcal{P}'(s) - \mathcal{P}''(s)$ is also a ν -measurable family of operators in $\mathfrak{S}$. As $\mathcal{P}'(s) - \mathcal{P}''(s)$ is the orthogonal projection of $\mathfrak{S}(s)$ onto $\mathfrak{S}'(s) \ominus \mathfrak{S}''(s)$ for ν -a.e. s , the desired statement follows from Lemma B.5.

2. Let ν -measurable families $\mathcal{P}'$ and $\mathcal{P}''$ of operators in $\mathfrak{S}$ be as in the proof of (1) and $\mathcal{R}$ be a ν -a.e. defined map such that $\mathcal{R}(s) = 1_{\mathfrak{S}(s)} - (\mathcal{P}'(s) + \mathcal{P}''(s))/2$ for ν -a.e. s . It follows from Lemma B.3 that $\mathcal{R}$ is a ν -measurable family of operators in $\mathfrak{S}$. Since $\mathfrak{S}'(s) \cap \mathfrak{S}''(s) = \text{Ker } \mathcal{R}(s) = (\text{Im } \mathcal{R}(s))^{\perp}$ for ν -a.e. s , the statement follows from Lemma B.4 and (1). $\square$

Lemma B.7. *Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces and $\mathcal{R}$ be a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$. Then*

1. *If $\mathcal{R}(s)$ is invertible for ν -a.e. s , then $s \rightarrow \mathcal{R}(s)^{-1}$ is a ν -measurable family of operators from $\mathfrak{S}_2$ to $\mathfrak{S}_1$.*

2. If $\mathcal{R}(s)$ is closed for ν -a.e. s , then $s \rightarrow \text{Ker } \mathcal{R}(s)$ is a ν -measurable family of subspaces of $\mathfrak{S}_1$.

Proof. For ν -a.e. s , let $\mathcal{S}(s)$ be everywhere defined bounded operator from $\mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s)$ to $\mathfrak{S}_2(s) \oplus \mathfrak{S}_1(s)$ taking (ψ_1, ψ_2) to (ψ_2, ψ_1) . Obviously, $\mathcal{S}$ is a ν -measurable family of operators from $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$ to $\mathfrak{S}_2 \oplus_\nu \mathfrak{S}_1$. If $\mathcal{R}(s)$ is invertible for ν -a.e. s , then $G_{\mathcal{R}(s)^{-1}}$ coincides with the image of $G_{\mathcal{R}(s)}$ under $\mathcal{S}(s)$ for ν -a.e. s and, therefore, (1) follows from Lemma B.4. Suppose now that $\mathcal{R}(s)$ is closed for ν -a.e. s . Since $s \rightarrow \mathfrak{S}_1(s) \times \{0\}$ is a ν -measurable family of closed subspaces of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$ and $G_{\mathcal{R}(s)} \cap (\mathfrak{S}_1(s) \times \{0\}) = \text{Ker } \mathcal{R}(s) \times \{0\}$ for ν -a.e. s , statement 2 of Lemma B.6 implies that $s \rightarrow \text{Ker } \mathcal{R}(s) \times \{0\}$ is a ν -measurable family of closed subspaces of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$, whence (2) obviously follows. $\square$

Lemma B.8. Let $\mathfrak{S}_1$, $\mathfrak{S}_2$, and $\mathfrak{S}_3$ be ν -measurable families of Hilbert spaces and $\mathcal{R}_1$ and $\mathcal{R}_2$ be ν -measurable families of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ and from $\mathfrak{S}_2$ to $\mathfrak{S}_3$ respectively. Then $s \rightarrow \mathcal{R}_2(s)\mathcal{R}_1(s)$ is a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_3$.

Proof. Let $\mathfrak{S}'$ and $\mathfrak{S}''$ be ν -a.e. defined maps such that, for ν -a.e. s , $\mathfrak{S}'(s)$ and $\mathfrak{S}''(s)$ are subspaces of $\mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s) \oplus \mathfrak{S}_3(s)$ consisting of all (ψ_1, ψ_2, ψ_3) with $(\psi_1, \psi_2) \in G_{\mathcal{R}_1(s)}$ and $(\psi_2, \psi_3) \in G_{\mathcal{R}_2(s)}$ respectively. Clearly, both $\mathfrak{S}'$ and $\mathfrak{S}''$ are ν -measurable families of closed subspaces of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2 \oplus_\nu \mathfrak{S}_3$. Let $\mathcal{S}$ be a ν -a.e. defined map such that, for ν -a.e. s , $\mathcal{S}(s)$ is the everywhere defined bounded operator from $\mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s) \oplus \mathfrak{S}_3(s)$ to $\mathfrak{S}_1(s) \oplus \mathfrak{S}_3(s)$ taking (ψ_1, ψ_2, ψ_3) to (ψ_1, ψ_3) . Then $G_{\mathcal{R}_2(s)\mathcal{R}_1(s)}$ is the image of $\mathfrak{S}'(s) \cap \mathfrak{S}''(s)$ under $\mathcal{S}(s)$ for ν -a.e. s . Hence, the result follows from Lemma B.4 and statement 2 of Lemma B.6. $\square$

Lemma B.9. Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces and $\mathcal{R}_1$ and $\mathcal{R}_2$ be ν -measurable families of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$. Then $s \rightarrow \mathcal{R}_2(s) + \mathcal{R}_1(s)$ is a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$.

Proof. Let $\mathfrak{S}'$ and $\mathfrak{S}''$ be ν -a.e. defined maps such that, for ν -a.e. s , $\mathfrak{S}'(s)$ and $\mathfrak{S}''(s)$ are subspaces of $\mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s) \oplus \mathfrak{S}_2(s)$ consisting of all (ψ_1, ψ_2, ψ_3) with $(\psi_1, \psi_2) \in G_{\mathcal{R}_1(s)}$ and $(\psi_1, \psi_3) \in G_{\mathcal{R}_2(s)}$ respectively. Clearly, both $\mathfrak{S}'$ and $\mathfrak{S}''$ are ν -measurable families of closed subspaces of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2 \oplus_\nu \mathfrak{S}_2$. Let $\mathcal{S}$ be a ν -a.e. defined map such that, for ν -a.e. s , $\mathcal{S}(s)$ is the everywhere defined bounded operator from $\mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s) \oplus \mathfrak{S}_2(s)$ to $\mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s)$ taking (ψ_1, ψ_2, ψ_3) to $(\psi_1, \psi_2 + \psi_3)$. Then $G_{\mathcal{R}_2(s) + \mathcal{R}_1(s)}$ is the image of $\mathfrak{S}'(s) \cap \mathfrak{S}''(s)$ under $\mathcal{S}(s)$ for ν -a.e. s . Hence, the result follows from Lemma B.4 and statement 2 of Lemma B.6. $\square$

Lemma B.10. Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces and $\mathcal{R}$ be a ν -measurable family of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$. Let ξ be a ν -measurable section of $\mathfrak{S}_1$ such that $\xi(s) \in D_{\mathcal{R}(s)}$ for ν -a.e. s . Then $s \rightarrow \mathcal{R}(s)\xi(s)$ is a ν -measurable section of $\mathfrak{S}_2$.

Proof. Let ζ be a ν -a.e. defined section of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$ such that $\zeta(s) = (\xi(s), \mathcal{R}(s)\xi(s))$ for ν -a.e. s . It suffices to show that ζ is ν -measurable. Let $\xi_1, \xi_2, \dots$ be a ν -measurable basis in $\mathfrak{S}_1$. Let $\mathcal{P}$ be a ν -a.e. defined map such that $\mathcal{P}(s)$ is the orthogonal projection of $\mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s)$ onto $G_{\mathcal{R}(s)}$ for ν -a.e. s . For $j = 1, 2, \dots$, let ζ_j be a ν -a.e. defined map such that $\zeta_j(s) = \mathcal{P}(s)(\xi_j(s), 0)$ for ν -a.e. s . By Lemma B.5, $\mathcal{P}$ is a ν -measurable family of operators in $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$, and Lemma B.3 implies that ζ_j is a ν -measurable section of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$ for all $j = 1, 2, \dots$. Moreover, the linear span of $\zeta_1(s), \zeta_2(s), \dots$ is dense in $G_{\mathcal{R}(s)}$ for ν -a.e. s .⁴⁴ As $\langle \zeta_j(s), \zeta(s) \rangle = \langle \xi_j(s), \xi(s) \rangle$ for ν -a.e. s , we conclude that $s \rightarrow \langle \zeta_j(s), \zeta(s) \rangle$ is a ν -measurable function for all $j = 1, 2, \dots$. Hence, ζ is ν -measurable by Lemma B.2. $\square$

Remark B.11. The definition of measurability of a family of operators given in this section is mainly the same as in Ref. 8. The only difference is that, in contrast to Ref. 8, we do not require operators to be closed and densely defined. This is essential for Propositions B.8 and B.9 because the properties of being closed and densely defined are not inherited by sums and products of operators.

Remark B.12. Let R be a closed operator in a Hilbert space $\mathfrak{H}$, $P(R)$ be the orthogonal projection of $\mathfrak{H} \oplus \mathfrak{H}$ onto G_R , and $\pi_{1,2}(\mathfrak{H}): \mathfrak{H} \oplus \mathfrak{H} \rightarrow \mathfrak{H}$ and $j_{1,2}(\mathfrak{H}): \mathfrak{H} \rightarrow \mathfrak{H} \oplus \mathfrak{H}$ be the canonical projections and embeddings respectively. For $i, k = 1, 2$, we set $P_{ik}(R) = \pi_i(\mathfrak{H})P(R)j_k(\mathfrak{H})$. The 2×2 -matrix composed of bounded operators $P_{ik}(R)$ is called the characteristic matrix of (generally, unbounded) operator R . In Ref. 7, a ν -a.e. defined family $\mathcal{R}$ of closed operators in a ν -measurable family $\mathfrak{S}$ of Hilbert spaces was called ν -measurable if $s \rightarrow P_{ik}(\mathcal{R}(s))\xi(s)$ is a ν -measurable section of $\mathfrak{S}$ for all $i, k = 1, 2$ and every ν -measurable section ξ of $\mathfrak{S}$. It follows from Lemmas B.3 and B.5 that this definition is equivalent to that given in this section because

$$P(\mathcal{R}(s)) = \sum_{i,k=1}^2 j_i(\mathfrak{S}(s))P_{ik}(\mathcal{R}(s))\pi_k(\mathfrak{S}(s))$$

for ν -a.e. s and $s \rightarrow \pi_{1,2}(\mathfrak{S}(s))$ and $s \rightarrow j_{1,2}(\mathfrak{S}(s))$ are obviously ν -measurable families (in our sense) of operators from $\mathfrak{S} \oplus_\nu \mathfrak{S}$ to $\mathfrak{S}$ and from $\mathfrak{S}$ to $\mathfrak{S} \oplus_\nu \mathfrak{S}$ respectively.

Remark B.13. Let $\mathcal{R}$ be a ν -a.e. defined family of operators in a ν -measurable family $\mathfrak{S}$ of Hilbert spaces. Suppose $\mathcal{R}(s)$ is self-adjoint for ν -a.e. s . Then $\mathcal{R}(s) + i$ has an everywhere defined bounded inverse for ν -a.e. s and Lemmas B.3 and B.7 imply that $\mathcal{R}$ is ν -measurable if and only if $s \rightarrow (\mathcal{R}(s) + i)^{-1}\xi(s)$ is a ν -measurable section of $\mathfrak{S}$ for any ν -measurable section ξ of $\mathfrak{S}$. In Ref. 45, the last condition was adopted as the definition of measurability for families of self-adjoint operators.

Remark B.14. A ν -a.e. defined family $\mathcal{R}$ of operators in a ν -measurable family $\mathfrak{S}$ of Hilbert spaces is said to be weakly ν -measurable⁷ if $s \rightarrow \mathcal{R}(s)\xi(s)$ is a ν -measurable section of $\mathfrak{S}$ for every ν -measurable section ξ of $\mathfrak{S}$ satisfying $\xi(s) \in D_{\mathcal{R}(s)}$ for ν -a.e. s . Lemma B.10 states that every ν -measurable family of closed operators is also weakly ν -measurable. In Ref. 7, where Lemma B.10 was originally proved, a question was posed whether this statement can be reverted, i.e., whether every weakly ν -measurable family of closed operators is ν -measurable. In Ref. 46, it was shown by constructing a counterexample that the answer is negative.

5. Direct integrals of Hilbert spaces

Given a ν -measurable family $\mathfrak{S}$ of Hilbert spaces, we denote by $\mathcal{M}(\mathfrak{S}, \nu)$ the set of all ν -equivalence classes $[f]_\nu$, where f is a ν -measurable section of $\mathfrak{S}$. Clearly, $\mathcal{M}(\mathfrak{S}, \nu)$ has a natural structure of a complex vector space (for any ν -measurable sections ξ and η and any $k \in \mathbb{C}$, we set $[\xi]_\nu + [\eta]_\nu = [\xi + \eta]_\nu$ and $k[\xi]_\nu = [k\xi]_\nu$). Now suppose $\mathfrak{S}'$ is a ν -a.e. defined family of subspaces of $\mathfrak{S}$. We denote by $(\mathfrak{S})\text{-}\int^\oplus \mathfrak{S}'(s) d\nu(s)$ the linear subspace of the space $\mathcal{M}(\mathfrak{S}, \nu)$ consisting of all its elements ξ such that $\xi(s) \in \mathfrak{S}'(s)$ for ν -a.e. s and $s \rightarrow \|\xi(s)\|^2$ is a ν -integrable function. The space $(\mathfrak{S})\text{-}\int^\oplus \mathfrak{S}'(s) d\nu(s)$ is endowed with the scalar product defined by the relation

$$\langle \xi, \eta \rangle = \int \langle \xi(s), \eta(s) \rangle d\nu(s).$$

For any ν -measurable family $\mathfrak{S}$, the space $(\mathfrak{S})\text{-}\int^\oplus \mathfrak{S}(s) d\nu(s)$ is complete (the proof is essentially the same as that of completeness of ordinary L_2 -spaces) and, hence, is a Hilbert space.

As a rule, the ν -measurable family $\mathfrak{S}$ can be easily deduced from the context. So we usually omit the prefix $(\mathfrak{S})\text{-}$ and write $\int^\oplus \mathfrak{S}'(s) d\nu(s)$ in place of $(\mathfrak{S})\text{-}\int^\oplus \mathfrak{S}'(s) d\nu(s)$. The simplest examples of direct integrals are L_2 -spaces and countable direct sums of Hilbert spaces, which can be represented using the constant and discrete families of Hilbert spaces respectively (see Sec. B3). Indeed, for any separable Hilbert space $\mathfrak{h}$, we obviously have

$$L_2(S_\nu, \mathfrak{h}, \nu) = \int^\oplus \mathcal{I}_{\mathfrak{h}, \nu}(s) d\nu(s). \quad (\text{B1})$$

Similarly, if $\mathcal{S}$ is a countable set, ν is a counting measure on $\mathcal{S}$, and $\mathfrak{S}$ is a ν -a.e. defined family of Hilbert spaces, then

$$\bigoplus_{s \in \mathcal{S}} \mathfrak{S}(s) = \int^\oplus \mathfrak{S}_{\mathcal{S}}(s) d\nu(s). \quad (\text{B2})$$

Given a ν -a.e. defined section ξ of $\mathfrak{S}$ and a ν -measurable set A , we denote by ξ^A the ν -equivalence class such that $\xi^A(s) = \xi(s)$ for ν -a.e. $s \in A$ and $\xi^A(s) = 0$ for ν -a.e. $s \in S_\nu \setminus A$.

Lemma B.15. *Let $\mathfrak{S}$ be a ν -measurable family of Hilbert spaces and $\mathfrak{S}'$ be a ν -measurable family of subspaces of $\mathfrak{S}$. Then there is a ν -measurable basis in $\mathfrak{S}'$ consisting of elements of $\int^\oplus \mathfrak{S}'(s) d\nu(s)$.*

Proof. Let $\xi_1, \xi_2, \dots$ be a ν -measurable basis in $\mathfrak{S}'$. Multiplying ξ_j by suitable ν -measurable functions, we can ensure that $\|\xi_j(s)\| \leq 1$ for all $j = 1, 2, \dots$ and ν -a.e. s . Let $A_1, A_2, \dots$ be elements of D_ν such that $S_\nu = \bigcup_{k=1}^\infty A_k$. Then $\xi_j^{A_k}$ with $j, k = 1, 2, \dots$ obviously constitute the required ν -measurable basis in $\mathfrak{S}'$. $\square$

Lemma B.16. *Let $\mathfrak{S}$ and $\mathfrak{S}'$ be as in Lemma B.15 and $\mathfrak{H}' = \int^\oplus \mathfrak{S}'(s) d\nu(s)$. Then $\mathfrak{H}'^\perp = \int^\oplus \mathfrak{S}'(s)^\perp d\nu(s)$ and $\overline{\mathfrak{H}'} = \int^\oplus \overline{\mathfrak{S}'(s)} d\nu(s)$.*

Proof. Let $\eta \in \mathfrak{H}'^\perp$ and $\xi \in \mathfrak{H}'$. As $\xi^A \in \mathfrak{H}'$ for any $A \in D_\nu$, we obtain $\int_A \langle \xi(s), \eta(s) \rangle d\nu(s) = \langle \xi_A, \eta \rangle = 0$ and, hence, $\langle \xi(s), \eta(s) \rangle = 0$ for ν -a.e. s . By Lemma B.15, there is a ν -measurable basis $\xi_1, \xi_2, \dots$ in $\mathfrak{S}'$ such that $\xi_j \in \mathfrak{H}'$ for all $j = 1, 2, \dots$. Since $\langle \xi_j(s), \eta(s) \rangle = 0$ for ν -a.e. s and all $j = 1, 2, \dots$, we have $\eta(s) \in \mathfrak{S}'(s)^\perp$ for ν -a.e. s . This implies that $\mathfrak{H}'^\perp = \int^\oplus \mathfrak{S}'(s)^\perp d\nu(s)$. The equality for the closure of $\mathfrak{H}'$ now follows from the relations $\overline{\mathfrak{H}'} = (\mathfrak{H}'^\perp)^\perp$ and $\overline{\mathfrak{S}'(s)} = (\mathfrak{S}'(s)^\perp)^\perp$. $\square$

Let $\mathfrak{S}$ be a ν -measurable family of Hilbert spaces and $\mathfrak{H} = \int^\oplus \mathfrak{S}(s) d\nu(s)$. Given a ν -measurable complex function g , we denote by $\mathcal{T}_g^{\nu, \mathfrak{S}}$ the operator of multiplication by g in $\mathfrak{H}$. By definition, the graph of $\mathcal{T}_g^{\nu, \mathfrak{S}}$ consists of all pairs $(\xi_1, \xi_2) \in \mathfrak{H} \oplus \mathfrak{H}$ such that $\xi_2(s) = g(s)\xi_1(s)$ for ν -a.e. s . The operator $\mathcal{T}_g^{\nu, \mathfrak{S}}$ is closed and densely defined and its adjoint is equal to $\mathcal{T}_{\bar{g}}^{\nu, \mathfrak{S}}$, where $\bar{g}$ is the complex conjugate function of g . In particular, if g is real, then $\mathcal{T}_g^{\nu, \mathfrak{S}}$ is self-adjoint. If g is ν -essentially bounded, then $\mathcal{T}_g^{\nu, \mathfrak{S}}$ is everywhere defined and bounded.

Lemma B.17. *Let $\mathfrak{S}$ be a ν -measurable family of Hilbert spaces, $\mathfrak{H} = \int^\oplus \mathfrak{S}(s) d\nu(s)$, and $\mathfrak{H}'$ be a closed linear subspace of $\mathfrak{H}$ such that $\mathcal{T}_g^{\nu, \mathfrak{S}} \xi \in \mathfrak{H}'$ for any $\xi \in \mathfrak{H}'$ and every ν -essentially bounded complex function g . Then there is a ν -measurable family $\mathfrak{S}'$ of closed subspaces of $\mathfrak{S}$ such that $\mathfrak{H}' = \int^\oplus \mathfrak{S}'(s) d\nu(s)$.*

Proof. By the hypothesis, the projection P of $\mathfrak{H}$ onto $\mathfrak{H}'$ commutes with $\mathcal{T}_g^{\nu, \mathfrak{S}}$ for every ν -essentially bounded function g . By Lemma B.15, there is a ν -measurable basis $\xi_1, \xi_2, \dots$ in $\mathfrak{S}$ consisting of elements of $\mathfrak{H}$. For each $j = 1, 2, \dots$, we set $\eta_j = P\xi_j$. Let $\mathfrak{S}'$ be a ν -measurable family of subspaces of $\mathfrak{S}$ such that $\mathfrak{S}'(s)$ is the closed linear span of $\eta_1(s), \eta_2(s), \dots$ for ν -a.e. s . Let $\tilde{\mathfrak{H}} = \int^\oplus \mathfrak{S}'(s) d\nu(s)$. If $\xi \in \mathfrak{H}$ is such that $\langle \xi_j^A, \xi \rangle = \int_A \langle \xi_j(s), \xi(s) \rangle d\nu(s) = 0$ for every ν -measurable set A and $j = 1, 2, \dots$, then $\langle \xi_j(s), \xi(s) \rangle = 0$ for all j and ν -a.e. s and, therefore, $\xi = 0$. This means that the linear span of all ξ_j^A is dense in $\mathfrak{H}$. Note that $\eta_j^A = \mathcal{T}_{\chi_A}^{\nu, \mathfrak{S}} P\xi_j = P\xi_j^A$, where $\chi_A(s) = 1$ for $s \in A$ and $\chi_A(s) = 0$ for $s \in S_\nu \setminus A$. Hence, the linear span of η_j^A is dense in $\mathfrak{H}'$. Since $\tilde{\mathfrak{H}}$ is closed and contains all η_j^A , we conclude that $\mathfrak{H}' \subset \tilde{\mathfrak{H}}$. Let $\eta \in \mathfrak{H}'^\perp$. Then we have $\langle \eta_j^A, \eta \rangle = \int_A \langle \eta_j(s), \eta(s) \rangle d\nu(s) = 0$ and, hence, $\langle \eta_j(s), \eta(s) \rangle = 0$ for all j and ν -a.e. s . This implies that $\eta(s) \in \mathfrak{S}'(s)^\perp$ for ν -a.e. s , i.e., $\eta \in \tilde{\mathfrak{H}}^\perp$. We therefore obtain $\mathfrak{H}'^\perp \subset \tilde{\mathfrak{H}}^\perp$. As $\mathfrak{H}'$ is closed, it follows that $\tilde{\mathfrak{H}} \subset \mathfrak{H}'$ and, hence, $\tilde{\mathfrak{H}} = \mathfrak{H}'$. $\square$

6. Direct integrals of operators

Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces. Let $\mathfrak{H}_{1,2} = \int^\oplus \mathfrak{S}_{1,2}(s) d\nu(s)$ and $\mathcal{R}$ be a ν -a.e. defined family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$. The direct integral $\int^\oplus \mathcal{R}(s) d\nu(s)$ of the family $\mathcal{R}$ is defined as the linear operator from $\mathfrak{H}_1$ to $\mathfrak{H}_2$ whose graph consists of all pairs $(\xi, \eta) \in \mathfrak{H}_1 \oplus \mathfrak{H}_2$ such that $\xi(s) \in D_{\mathcal{R}(s)}$ and $\mathcal{R}(s)\xi(s) = \eta(s)$ for ν -a.e. s .

Let $\mathfrak{H} = \int^\oplus \mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s) d\nu(s)$. Then there is a unique unitary operator $U_{\mathfrak{S}_1, \mathfrak{S}_2}: \mathfrak{H}_1 \oplus \mathfrak{H}_2 \rightarrow \mathfrak{H}$ such that $(U_{\mathfrak{S}_1, \mathfrak{S}_2}(\xi, \eta))(s) = (\xi(s), \eta(s))$ for any $(\xi, \eta) \in \mathfrak{H}_1 \oplus \mathfrak{H}_2$ and ν -a.e. s . We call $U_{\mathfrak{S}_1, \mathfrak{S}_2}$ the natural isomorphism between $\mathfrak{H}_1 \oplus \mathfrak{H}_2$ and $\mathfrak{H}$.

The next statement follows immediately from the definition of $U_{\mathfrak{S}_1, \mathfrak{S}_2}$.

Lemma B.18. *Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces, $\mathcal{R}$ be a ν -a.e. defined family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$, and $R = \int^\oplus \mathcal{R}(s) d\nu(s)$. Then we have $U_{\mathfrak{S}_1, \mathfrak{S}_2}(G_R) = (\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2) - \int^\oplus G_{\mathcal{R}(s)} d\nu(s)$.*

Lemma B.19. *Let $\mathfrak{H}_{1,2} = \int^\oplus \mathfrak{S}_{1,2}(s) d\nu(s)$, where $\mathfrak{S}_1$ and $\mathfrak{S}_2$ are ν -measurable families of Hilbert spaces, and G be a closed linear subspace of $\mathfrak{H}_1 \oplus \mathfrak{H}_2$. Suppose $\mathcal{G}$ is a ν -measurable family of closed subspaces of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$ such that $U_{\mathfrak{S}_1, \mathfrak{S}_2}(G) = \int^\oplus \mathcal{G}(s) d\nu(s)$. Then G is a graph of an operator from $\mathfrak{H}_1$ to $\mathfrak{H}_2$ if and only if $\mathcal{G}(s)$ is a graph of an operator from $\mathfrak{S}_1(s)$ to $\mathfrak{S}_2(s)$ for ν -a.e. s .*

Proof. Let $U = U_{\mathfrak{S}_1, \mathfrak{S}_2}$ and $\mathfrak{S}'$ be a ν -a.e. defined map such that $\mathfrak{S}'(s) = \{0\} \times \mathfrak{S}_2(s)$ for ν -a.e. s . Clearly, $\mathfrak{S}'$ is a ν -measurable family of closed subspaces of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$. Let $\mathfrak{H}' = \int^\oplus \mathfrak{S}'(s) d\nu(s)$ and $\mathfrak{H}'' = \int^\oplus \mathfrak{S}'(s) \cap \mathcal{G}(s) d\nu(s)$. Since $\mathfrak{H}'' = \mathfrak{H}' \cap U(G)$ and $U^{-1}(\mathfrak{H}') = \{0\} \times \mathfrak{H}_2$, we have $G \cap (\{0\} \times \mathfrak{H}_2) = U^{-1}(\mathfrak{H}'')$. Hence, G is a graph of an operator if and only if $\mathfrak{H}''$ is trivial. By statement 2 of Lemma B.6, $s \rightarrow \mathfrak{S}'(s) \cap \mathcal{G}(s)$ is a ν -measurable family of subspaces of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$ and, therefore, $\mathfrak{H}''$ is trivial if and only if $\mathfrak{S}'(s) \cap \mathcal{G}(s)$ is trivial for ν -a.e. s , i.e., if $\mathcal{G}(s)$ is a graph of an operator for ν -a.e. s . $\square$

Lemma B.20. *Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces, $\mathcal{R}$ be a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$, and $R = \int^\oplus \mathcal{R}(s) d\nu(s)$. Then R is closable if and only if $\mathcal{R}(s)$ is closable for ν -a.e. s , in which case $\overline{R} = \int^\oplus \overline{\mathcal{R}(s)} d\nu(s)$.*

Proof. Let $U = U_{\mathfrak{S}_1, \mathfrak{S}_2}$. As $U(G_R) = \int^\oplus G_{\mathcal{R}(s)} d\nu(s)$, Lemma B.16 yields

$$U(\overline{G_R}) = \int^\oplus \overline{G_{\mathcal{R}(s)}} d\nu(s). \quad (\text{B3})$$

The operator R is closable if and only if $\overline{G_R}$ is a graph of an operator. By Lemma B.19, the latter condition holds if and only if $\overline{G_{\mathcal{R}(s)}}$ is a graph of an operator for ν -a.e. s , i.e., if $\mathcal{R}(s)$ is closable for ν -a.e. s . Suppose now that R is closable and $R' = \int^\oplus \overline{\mathcal{R}(s)} d\nu(s)$. As $\overline{G_R} = G_{\overline{R}}$ and $\overline{G_{\mathcal{R}(s)}} = G_{\overline{\mathcal{R}(s)}}$ for ν -a.e. s , equality (B3) and Lemma B.18 imply that $U(G_{\overline{R}}) = U(G_{R'})$ and, hence, $R = R'$. $\square$

Proposition B.21. *Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces and $\mathcal{R}$ and $\mathcal{R}'$ be ν -measurable families of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$. Let $R = \int^\oplus \mathcal{R}(s) d\nu(s)$, and $R' = \int^\oplus \mathcal{R}'(s) d\nu(s)$. If R is an extension of R' , then $\mathcal{R}(s)$ is an extension of $\mathcal{R}'(s)$ for ν -a.e. s . If $R = R'$, then $\mathcal{R}(s) = \mathcal{R}'(s)$ for ν -a.e. s .*

Proof. Let $U = U_{\mathfrak{S}_1, \mathfrak{S}_2}$. Since $U(G_{R'}) = \int^\oplus G_{\mathcal{R}'(s)} d\nu(s)$, Lemma B.15 implies that there exists a sequence $\zeta_1, \zeta_2, \dots$ of elements of $U(G_{R'})$ such that the linear span of $\zeta_1(s), \zeta_2(s), \dots$ is dense in $G_{\mathcal{R}'(s)}$ for ν -a.e. s . If R is an extension of R' , then $U(G_{R'}) \subset U(G_R)$ and, hence, $\zeta_j \in U(G_R)$ for all $j = 1, 2, \dots$. Since $U(G_R) = \int^\oplus G_{\mathcal{R}(s)} d\nu(s)$, it follows that $\zeta_j(s) \in G_{\mathcal{R}(s)}$ for ν -a.e. s and all $j = 1, 2, \dots$. The linear span of $\zeta_1(s), \zeta_2(s), \dots$ is therefore contained in $G_{\mathcal{R}(s)}$ for ν -a.e. s . As $\mathcal{R}(s)$ is closed, it follows that $G_{\mathcal{R}'(s)} \subset G_{\mathcal{R}(s)}$, i.e., $\mathcal{R}(s)$ is an extension of $\mathcal{R}'(s)$ for ν -a.e. s . If $R = R'$, then, by the above, $\mathcal{R}(s)$ and $\mathcal{R}'(s)$ are extensions of each other and, therefore, are equal for ν -a.e. s . $\square$

Proposition B.22. *Let $\mathfrak{S}_1, \mathfrak{S}_2, \mathfrak{H}_1$, and $\mathfrak{H}_2$ be as in Lemma B.19. Let $\mathcal{R}$ be a ν -measurable family of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$. Then $R = \int^\oplus \mathcal{R}(s) d\nu(s)$ is a closed operator from $\mathfrak{H}_1$ to $\mathfrak{H}_2$ satisfying the condition:*

(M) *If $\xi \in D_R$ and g is a ν -measurable ν -essentially bounded complex function, then $\mathcal{T}_g^{\nu, \mathfrak{S}_1} \xi \in D_R$ and $R \mathcal{T}_g^{\nu, \mathfrak{S}_1} \xi = \mathcal{T}_g^{\nu, \mathfrak{S}_2} R \xi$.*

Conversely, if R is a closed operator from $\mathfrak{H}_1$ to $\mathfrak{H}_2$ satisfying (M), then there is a unique (up to ν -equivalence) ν -measurable family $\mathcal{R}$ of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ such that $R = \int^\oplus \mathcal{R}(s) d\nu(s)$.

Proof. Let $U = U_{\mathfrak{S}_1, \mathfrak{S}_2}$, $\mathcal{R}$ be a ν -measurable family of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$, and $R = \int^\oplus \mathcal{R}(s) d\nu(s)$. By Lemma B.20, R is closed. If $\xi \in D_R$ and g is a ν -measurable ν -essentially bounded function, then $(\mathcal{T}_g^{\nu, \mathfrak{S}_1} \xi, \mathcal{T}_g^{\nu, \mathfrak{S}_2} R \xi)$ belongs to G_R by the very definition of the direct integral of operators. This means that (M) is fulfilled. Conversely, let R be a closed operator from $\mathfrak{H}_1$ to $\mathfrak{H}_2$ satisfying (M). If $(\xi, \eta) \in G_R$ and g is a ν -measurable ν -essentially bounded function, then $(\mathcal{T}_g^{\nu, \mathfrak{S}_1} \xi, \mathcal{T}_g^{\nu, \mathfrak{S}_2} \eta)$ belongs to G_R by (M) and, hence, $\mathcal{T}_g^{\nu, \mathfrak{S}_1} \oplus \mathcal{T}_g^{\nu, \mathfrak{S}_2} U(\xi, \eta) = U(\mathcal{T}_g^{\nu, \mathfrak{S}_1} \xi, \mathcal{T}_g^{\nu, \mathfrak{S}_2} \eta)$ belongs to $U(G_R)$. This means that $U(G_R)$ is invariant under $\mathcal{T}_g^{\nu, \mathfrak{S}_1} \oplus \mathcal{T}_g^{\nu, \mathfrak{S}_2}$. By Lemma B.17, there is a ν -measurable family $\mathcal{G}$ of closed subspaces of $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$ such that $U(G_R) = \int^\oplus \mathcal{G}(s) d\nu(s)$. By Lemma B.19, there is a ν -measurable family $\mathcal{R}$ of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ such that $\mathcal{G}(s) = G_{\mathcal{R}(s)}$ for ν -a.e. s . If $R' = \int^\oplus \mathcal{R}(s) d\nu(s)$, then it follows from Lemma B.18 that $U(G_{R'}) = U(G_R)$ and, hence, $R = R'$. The uniqueness of $\mathcal{R}$ is ensured by Proposition B.21. $\square$

Let $\mathfrak{H}_1$ and $\mathfrak{H}_2$ be Hilbert spaces and R be an operator from $\mathfrak{H}_1$ to $\mathfrak{H}_2$. We define the linear subspaces G_R° and G_R^* of $\mathfrak{H}_2 \oplus \mathfrak{H}_1$ by setting

$$G_R^\circ = \{(-\psi_2, \psi_1) : (\psi_1, \psi_2) \in G_R\}, \quad G_R^* = (G_R^\circ)^\perp.$$

The operator R is densely defined if and only if G_R^* is the graph of an operator, in which case $G_{R^*} = G_R^*$.

Proposition B.23. *Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces, $\mathcal{R}$ be a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$, and $R = \int^\oplus \mathcal{R}(s) d\nu(s)$. Then the following statements hold:*

1. *Suppose $\mathcal{R}(s)$ is closed for ν -a.e. s . The operator R is invertible if and only if $\mathcal{R}(s)$ is invertible for ν -a.e. s , in which case $s \rightarrow \mathcal{R}(s)^{-1}$ is a ν -measurable family of operators from $\mathfrak{S}_2$ to $\mathfrak{S}_1$ and $R^{-1} = \int^\oplus \mathcal{R}(s)^{-1} d\nu(s)$.*

2. The operator R is densely defined if and only if $\mathcal{R}(s)$ is densely defined for ν -a.e. s , in which case $s \rightarrow \mathcal{R}(s)^*$ is a ν -measurable family of operators from $\mathfrak{S}_2$ to $\mathfrak{S}_1$ and $R^* = \int^\oplus \mathcal{R}(s)^* d\nu(s)$.

Proof. Let $\mathfrak{H}_{1,2} = \int^\oplus \mathfrak{S}_{1,2}(s) d\nu(s)$.

1. By statement 2 of Lemma B.7, $s \rightarrow \text{Ker } \mathcal{R}(s)$ is a ν -measurable family of subspaces of $\mathfrak{S}_1$. As $\text{Ker } R = \int^\oplus \text{Ker } \mathcal{R}(s) d\nu(s)$, we conclude that R is invertible if and only if $\text{Ker } \mathcal{R}(s) = \{0\}$ for ν -a.e. s , i.e., if $\mathcal{R}(s)$ is invertible for ν -a.e. s . Let R be invertible and $R' = \int^\oplus \mathcal{R}(s)^{-1} d\nu(s)$. For any $(\eta, \xi) \in \mathfrak{H}_2 \oplus \mathfrak{H}_1$, we have

$$(\eta, \xi) \in G_{R^{-1}} \iff (\xi, \eta) \in G_R \iff (\xi(s), \eta(s)) \in G_{\mathcal{R}(s)} \text{ for } \nu\text{-a.e. } s \iff (\eta(s), \xi(s)) \in G_{\mathcal{R}(s)^{-1}} \text{ for } \nu\text{-a.e. } s \iff (\eta, \xi) \in G_{R'}.$$

We thus have $G_{R^{-1}} = G_{R'}$ and, hence, $R = R'$. By statement 1 of Lemma B.7, $s \rightarrow \mathcal{R}(s)^{-1}$ is a ν -measurable family of operators from $\mathfrak{S}_2$ to $\mathfrak{S}_1$.

2. Let $U_{12} = U_{\mathfrak{S}_1, \mathfrak{S}_2}$ and $U_{21} = U_{\mathfrak{S}_2, \mathfrak{S}_1}$. For ν -a.e. s , let $\mathcal{S}(s)$ be everywhere defined bounded operator from $\mathfrak{S}_1(s) \oplus \mathfrak{S}_2(s)$ to $\mathfrak{S}_2(s) \oplus \mathfrak{S}_1(s)$ taking (ψ_1, ψ_2) to $(-\psi_2, \psi_1)$. Clearly, $\mathcal{S}$ is a ν -measurable family of unitary operators from $\mathfrak{S}_1 \oplus_\nu \mathfrak{S}_2$ to $\mathfrak{S}_2 \oplus_\nu \mathfrak{S}_1$. Note that $G_{\mathcal{R}(s)}^\circ$ is the image of $G_{\mathcal{R}(s)}$ under $\mathcal{S}(s)$ for ν -a.e. s . Hence, $s \rightarrow G_{\mathcal{R}(s)}^\circ$ is a ν -measurable family of subspaces of $\mathfrak{S}_2 \oplus_\nu \mathfrak{S}_1$ by Lemma B.4. Let $S = \int^\oplus \mathcal{S}(s) d\nu(s)$ and $\tilde{S}$ be the unitary operator from $\mathfrak{H}_1 \oplus \mathfrak{H}_2$ to $\mathfrak{H}_2 \oplus \mathfrak{H}_1$ defined by the relation $\tilde{S}(\xi, \eta) = (-\eta, \xi)$. It is straightforward to check that $U_{21}\tilde{S} = SU_{12}$. As $G_R^\circ = \tilde{S}(G_R)$, we have $U_{21}(G_R^\circ) = S(U_{12}(G_R))$. Since $U_{12}(G_R) = \int^\oplus G_{\mathcal{R}(s)} d\nu(s)$ by Lemma B.18 and the image of $\int^\oplus G_{\mathcal{R}(s)} d\nu(s)$ under S is equal to $\int^\oplus G_{\mathcal{R}(s)}^\circ d\nu(s)$, we conclude that $U_{21}(G_R^\circ) = \int^\oplus G_{\mathcal{R}(s)}^\circ d\nu(s)$. It now follows from Lemmas B.6 and B.16 and the unitarity of U_{21} that $s \rightarrow G_{\mathcal{R}(s)}^*$ is a ν -measurable family of subspaces of $\mathfrak{S}_2 \oplus_\nu \mathfrak{S}_1$ and

$$U_{21}(G_R^*) = \int^\oplus G_{\mathcal{R}(s)}^* d\nu(s). \quad (\text{B4})$$

In view of (B4), Lemma B.19 implies that R is densely defined if and only if $G_{\mathcal{R}(s)}^*$ is the graph of an operator for ν -a.e. s , i.e., if $\mathcal{R}(s)$ is densely defined for ν -a.e. s . Suppose R is densely defined and $R' = \int^\oplus \mathcal{R}(s)^* d\nu(s)$. As $G_{\mathcal{R}(s)}^* = G_{\mathcal{R}(s)^*}$ for ν -a.e. s , it follows that $s \rightarrow \mathcal{R}(s)^*$ is a ν -measurable family of operators from $\mathfrak{S}_2$ to $\mathfrak{S}_1$. Finally, since $G_R^* = G_{R^*}$, equality (B4) and Lemma B.18 imply that $U_{21}(G_{R^*}) = U_{21}(G_{R'})$ and, hence, $R^* = R'$. $\square$

Corollary B.24. Let $\mathfrak{S}$ be a ν -measurable family of Hilbert spaces, and $\mathcal{R}$ be a ν -measurable family of closed operators in $\mathfrak{S}$. The operator $\int^\oplus \mathcal{R}(s) d\nu(s)$ is self-adjoint if and only if $\mathcal{R}(s)$ is self-adjoint for ν -a.e. s .

Proof. The statement follows from Propositions B.23 and B.21. $\square$

Proposition B.25. Let $\mathfrak{S}_1, \mathfrak{S}_2$, and $\mathfrak{S}_3$ be ν -measurable families of Hilbert spaces and $\mathcal{R}_1$ and $\mathcal{R}_2$ be ν -measurable families of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ and from $\mathfrak{S}_2$ to $\mathfrak{S}_3$ respectively. Let $R_{1,2} = \int^\oplus \mathcal{R}_{1,2}(s) d\nu(s)$. The operator $R_2 R_1$ is densely defined if and only if the operator $\mathcal{R}_2(s)\mathcal{R}_1(s)$ is densely defined for ν -a.e. s . The operator $\overline{R_2 R_1}$ is closable if and only if the operator $\mathcal{R}_2(s)\mathcal{R}_1(s)$ is closable for ν -a.e. s , in which case $s \rightarrow \overline{\mathcal{R}_2(s)\mathcal{R}_1(s)}$ is a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_3$ and $\overline{R_2 R_1} = \int^\oplus \overline{\mathcal{R}_2(s)\mathcal{R}_1(s)} d\nu(s)$.

Proof. Let $\mathfrak{H}_i = \int^\oplus \mathfrak{S}_i(s) d\nu(s)$, $i = 1, 2, 3$, and $R = \int^\oplus \mathcal{R}_2(s)\mathcal{R}_1(s) d\nu(s)$. Clearly, R is an extension of $R_2 R_1$. Let $(\xi, \eta) \in G_R$. Then $\xi(s) \in D_{\mathcal{R}_2(s)\mathcal{R}_1(s)}$ and, hence, $\xi(s) \in D_{\mathcal{R}_1(s)}$ for ν -a.e. s . Let $\tilde{\eta}$ be a ν -a.e. defined section of $\mathfrak{S}_2$ such that $\tilde{\eta}(s) = \mathcal{R}_1(s)\xi(s)$ for ν -a.e. s . By Lemma B.10, $\tilde{\eta}$ is ν -measurable. Let $A_1 \subset A_2 \subset \dots$ be a sequence of elements of D_ν such that $S_\nu = \bigcup_{k=1}^\infty A_k$ and $\|\tilde{\eta}(s)\| \leq k$ for ν -a.e. $s \in A_k$. Set $\xi_k = \xi^{A_k}$, $\eta_k = \eta^{A_k}$, and $\tilde{\eta}_k = \tilde{\eta}^{A_k}$. We obviously have $\xi_k \in \mathfrak{H}_1$, $\tilde{\eta}_k \in \mathfrak{H}_2$, and $\eta_k \in \mathfrak{H}_3$ for all k . As $\tilde{\eta}(s) \in D_{\mathcal{R}_2(s)}$ and $\eta(s) = \mathcal{R}_2(s)\tilde{\eta}(s)$ for ν -a.e. s , we have $(\xi_k, \tilde{\eta}_k) \in G_{R_1}$ and $(\tilde{\eta}_k, \eta_k) \in G_{R_2}$ and, therefore, $(\xi_k, \eta_k) \in G_{R_2 R_1}$ for all k . Since $(\xi_k, \eta_k) \rightarrow (\xi, \eta)$ in $\mathfrak{H}_1 \oplus \mathfrak{H}_3$ as $k \rightarrow \infty$, we conclude that $G_{R_2 R_1}$ is dense in G_R and, hence, $D_{R_2 R_1}$ is dense in D_R . In view of statement 2 of Proposition B.23, it follows that $R_2 R_1$ is densely defined if and only if $\mathcal{R}_2(s)\mathcal{R}_1(s)$ is densely defined for ν -a.e. s . Since $\overline{G_R} = \overline{G_{R_2 R_1}}$, the operator $R_2 R_1$ is closable if and only if R is closable. In view of Lemma B.20, the latter condition holds if and only if $\mathcal{R}_2(s)\mathcal{R}_1(s)$ is closable for ν -a.e. s (note that $s \rightarrow \mathcal{R}_2(s)\mathcal{R}_1(s)$ is a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_3$ by Lemma B.8). If $R_2 R_1$ is closable, then Lemma B.20 implies that $\overline{R_2 R_1} = \overline{R} = \int^\oplus \overline{\mathcal{R}_2(s)\mathcal{R}_1(s)} d\nu(s)$. $\square$

Proposition B.26. *Let $\mathfrak{S}_1$ and $\mathfrak{S}_2$ be ν -measurable families of Hilbert spaces, $\mathcal{R}_1$ and $\mathcal{R}_2$ be ν -measurable families of closed operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$, and $R_{1,2} = \int^\oplus \mathcal{R}_{1,2}(s) d\nu(s)$. The operator $R_1 + R_2$ is densely defined if and only if the operator $\mathcal{R}_1(s) + \mathcal{R}_2(s)$ is densely defined for ν -a.e. s . The operator $R_1 + R_2$ is closable if and only if the operator $\mathcal{R}_1(s) + \mathcal{R}_2(s)$ is closable for ν -a.e. s , in which case $s \rightarrow \overline{\mathcal{R}_1(s) + \mathcal{R}_2(s)}$ is a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ and $\overline{R_1 + R_2} = \int^\oplus \overline{\mathcal{R}_1(s) + \mathcal{R}_2(s)} d\nu(s)$.*

Proof. Let $\mathfrak{H}_{1,2} = \int^\oplus \mathfrak{S}_{1,2}(s) d\nu(s)$ and $R = \int^\oplus (\mathcal{R}_1(s) + \mathcal{R}_2(s)) d\nu(s)$. Clearly, R is an extension of $R_1 + R_2$. Let $(\xi, \eta) \in G_R$. Then $\xi(s) \in D_{\mathcal{R}_1(s)} \cap D_{\mathcal{R}_2(s)}$ for ν -a.e. s . Let $\eta^{(1)}$ and $\eta^{(2)}$ be ν -a.e. defined sections of $\mathfrak{S}_2$ such that $\eta^{(1,2)}(s) = \mathcal{R}_{1,2}(s)\xi(s)$ for ν -a.e. s . By Lemma B.10, $\eta^{(1)}$ and $\eta^{(2)}$ are ν -measurable. Let $A_1 \subset A_2 \subset \dots$ be a sequence of elements of D_ν such that $S_\nu = \bigcup_{k=1}^\infty A_k$ and $\|\eta^{(1)}(s)\| + \|\eta^{(2)}(s)\| \leq k$ for ν -a.e. $s \in A_k$. Set $\xi_k = \xi^{A_k}$, $\eta_k = \eta^{A_k}$, $\eta_k^{(1,2)} = (\eta^{(1,2)})^{A_k}$. We obviously have $\xi_k \in \mathfrak{H}_1$ and $\eta_k, \eta_k^{(1,2)} \in \mathfrak{H}_2$ for all k . As $(\xi_k, \eta_k^{(1,2)}) \in G_{R_{1,2}}$ and $\eta_k = \eta_k^{(1)} + \eta_k^{(2)}$, we have $(\xi_k, \eta_k) \in G_{R_1 + R_2}$ for all k . Since $(\xi_k, \eta_k) \rightarrow (\xi, \eta)$ in $\mathfrak{H}_1 \oplus \mathfrak{H}_2$ as $k \rightarrow \infty$, we conclude that $G_{R_1 + R_2}$ is dense in G_R and, hence, $D_{R_1 + R_2}$ is dense in D_R . In view of statement 2 of Proposition B.23, it follows that $R_1 + R_2$ is densely defined if and only if $\mathcal{R}_1(s) + \mathcal{R}_2(s)$ is densely defined for ν -a.e. s . Since $\overline{G_R} = \overline{G_{R_1 + R_2}}$, the operator $R_1 + R_2$ is closable if and only if R is closable. In view of Lemma B.20, the latter condition holds if and only if $\mathcal{R}_1(s) + \mathcal{R}_2(s)$ is closable for ν -a.e. s (note that $s \rightarrow \mathcal{R}_1(s) + \mathcal{R}_2(s)$ is a ν -measurable family of operators from $\mathfrak{S}_1$ to $\mathfrak{S}_2$ by Lemma B.9). If $R_1 + R_2$ is closable, then Lemma B.20 implies that $\overline{R_1 + R_2} = \overline{R} = \int^\oplus \overline{\mathcal{R}_1(s) + \mathcal{R}_2(s)} d\nu(s)$. $\square$

Remark B.27. Propositions B.21, B.22, and B.23 were proved in Ref. 7 (see also Ref. 47). They are generalizations of corresponding statements proved in Ref. 6 for bounded operators. Propositions B.25 and B.26 were obtained in Ref. 48 under additional assumption that ν is a finite Borel measure on a metrizable compact space and were proved in Ref. 43 in their general form.

¹S. Albeverio, F. Gesztesy, R. Høegh-Krohn, and H. Holden, *Solvable models in quantum mechanics*, Texts and Monographs in Physics (Springer-Verlag, New York, 1988).

²D. M. Gitman, I. V. Tyutin, and B. L. Voronov, *Self-adjoint extensions in quantum mechanics; general theory and applications to Schrödinger and Dirac equations with singular potentials*, Progress in Mathematical Physics, Vol. 62 (Birkhäuser/Springer, New York, 2012).

³N. I. Akhiezer and I. M. Glazman, *Theory of linear operators in Hilbert space. Vol. I*, Monographs and Studies in Mathematics, Vol. 9 (Pitman (Advanced Publishing Program), Boston, Mass.-London, 1981) translated from the third Russian edition by E. R. Dawson, Translation edited by W. N. Everitt.

⁴A. G. Smirnov, “Generators of von Neumann algebras associated with spectral measures,” *J. Math. Anal. Appl.* **398**, 501–507 (2013).

⁵The counting measure on $\mathcal{S}$ is, by definition, the measure assigning to every finite subset of $\mathcal{S}$ the number of points in this set.

⁶J. von Neumann, “On rings of operators. Reduction theory,” *Ann. of Math. (2)* **50**, 401–485 (1949).

⁷A. E. Nussbaum, “Reduction theory for unbounded closed operators in Hilbert space,” *Duke Math. J.* **31**, 33–44 (1964).

⁸R. Pallu de la Barrière, “Décomposition des opérateurs non bornés dans les sommes continues d’espaces de Hilbert,” *C. R. Acad. Sci. Paris* **232**, 2071–2073 (1951).

⁹Throughout the paper, a.e. means either ‘almost every’ or ‘almost everywhere’.

¹⁰Y. Aharonov and D. Bohm, “Significance of electromagnetic potentials in the quantum theory,” *Phys. Rev. (2)* **115**, 485–491 (1959).

¹¹I. V. Tyutin, “Electron scattering on a solenoid,” arXiv:0801.2167 [quant-ph] (translated from preprint FIAN 27, P.N. Lebedev Physical Institute, Moscow, 1974).

¹²R. Adami and A. Teta, “On the Aharonov-Bohm Hamiltonian,” *Lett. Math. Phys.* **43**, 43–53 (1998).

¹³L. Dąbrowski and P. Šťovíček, “Aharonov-Bohm effect with δ -type interaction,” *J. Math. Phys.* **39**, 47–62 (1998).

¹⁴We endow S with the topology induced from $\mathbb{R}^2$. The Borel σ -algebra of S consists of all Borel subsets of $\mathbb{R}^2$ that are contained in S . The Borel structure on A^ϕ is induced from S .

¹⁵For brevity, we let $\mathcal{W}_\theta^\phi(s, E|x)$ denote the value of the function $\mathcal{W}_\theta^\phi(s, E)$ at the point x : $\mathcal{W}_\theta^\phi(s, E|x) = (\mathcal{W}_\theta^\phi(s, E))(x)$. A similar notation will be used for any maps whose values are also maps.

¹⁶To compute the limit of $u_\theta^\kappa(E|r)$ as $\kappa \rightarrow 0$, one has to apply L’Hôpital’s rule and use the equality $\Gamma'(1+n)/\Gamma(1+n) = c_n - \gamma$ (see Ref. 49, Sec. 1.7.1, formula (9)).

¹⁷See Ref. 49, Sec. 7.2.4, formula (33).

¹⁸A. G. Smirnov, “Eigenfunction expansions for the Schrödinger equation with inverse-square potential,” *Theor. Math. Phys.* (to be published); e-print arXiv:1508.07747 [math-ph].

¹⁹J. Dixmier, *Les algèbres d’opérateurs dans l’espace hilbertien (algèbres de von Neumann)* (Gauthier-Villars Éditeur, Paris, 1969) deuxième édition, revue et augmentée, Cahiers Scientifiques, Fasc. XXV.

²⁰Here and subsequently, S_ν denotes the largest ν -measurable set, see Sec. A 2.

²¹See Sec. A 3.

²²Recall that a closed densely defined linear operator T in a Hilbert space is called normal if the operators TT^* and T^*T have the same domain of definition and coincide thereon. In particular, self-adjoint and unitary operators are normal.

²³B. Fuglede, “A commutativity theorem for normal operators,” *Proc. Nat. Acad. Sci. U. S. A.* **36**, 35–40 (1950).

²⁴M. A. Naïmark, *Linear differential operators. Part II: Linear differential operators in Hilbert space*, With additional material by the author, and a supplement by V. È. Ljance. Translated from the Russian by E. R. Dawson. English translation edited by W. N. Everitt (Frederick Ungar Publishing Co., New York, 1968).

- ²⁵G. Teschl, *Mathematical methods in quantum mechanics, with applications to Schrödinger operators*, Graduate Studies in Mathematics, Vol. 99 (American Mathematical Society, Providence, RI, 2009).
- ²⁶J. Weidmann, *Spectral theory of ordinary differential operators*, Lecture Notes in Mathematics, Vol. 1258 (Springer-Verlag, Berlin, 1987).
- ²⁷Throughout this section and Sec. VII, all equivalence classes are taken with respect to $\lambda_{a,b}$. We shall drop the subscript and write $[f]$ instead of $[f]_{\lambda_{a,b}}$.
- ²⁸F. Gesztesy and M. Zinchenko, “On spectral theory for Schrödinger operators with strongly singular potentials,” *Math. Nachr.* **279**, 1041–1082 (2006).
- ²⁹D. M. Gitman, I. V. Tyutin, and B. L. Voronov, “Self-adjoint extensions and spectral analysis in the Calogero problem,” *J. Phys. A* **43**, 145205, 34 (2010).
- ³⁰A. Kostenko, A. Sakhnovich, and G. Teschl, “Weyl-Titchmarsh theory for Schrödinger operators with strongly singular potentials,” *Int. Math. Res. Not.* **2012**, 1699–1747 (2012).
- ³¹As in Sec. VI, we write $[f]$ in place of $[f]_{\lambda_{a,b}}$, where $\lambda_{a,b}$ is the restriction to (a, b) of the Lebesgue measure λ on $\mathbb{R}$.
- ³²We denote by $\xi'(s|r)$ the derivative of $\xi(s)$ at point r : $\xi'(s|r) = (\xi(s))'(r)$.
- ³³Given $\psi = (\psi_1, \psi_2) \in \mathfrak{h} \oplus \mathfrak{h}$, we set $\bar{\psi} = (\bar{\psi}_1, \bar{\psi}_2)$, where $\bar{\psi}_1$ and $\bar{\psi}_2$ are elements of $\mathfrak{h}$ such that $\bar{\psi}_{1,2}(r) = \overline{\psi_{1,2}(r)}$ for λ -a.e. $r \in (a, b)$.
- ³⁴H. Bauer, *Measure and integration theory*, de Gruyter Studies in Mathematics, Vol. 26 (Walter de Gruyter & Co., Berlin, 2001) translated from the German by Robert B. Burckel.
- ³⁵It is easy to see that the intersection of any set of α -classes is again an α -class (note that such an intersection is always nonempty because $\emptyset$ is an element of every α -class). Hence, given a set of sets $\mathcal{K}$, there exists the smallest α -class containing $\mathcal{K}$.
- ³⁶D. L. Cohn, *Measure theory* (Birkhäuser, Boston, Mass., 1980).
- ³⁷G. W. Mackey, “Borel structure in groups and their duals,” *Trans. Amer. Math. Soc.* **85**, 134–165 (1957).
- ³⁸V. A. Rohlin, “On the fundamental ideas of measure theory,” *Mat. Sbornik N.S.* **25(67)**, 107–150 (1949).
- ³⁹Note that a Hilbert space is separable if it is either finite-dimensional or infinite-dimensional with a countable orthonormal basis. In particular, we consider $\mathbb{C}$ as a one-dimensional Hilbert space with the scalar product $\langle \alpha, \beta \rangle = \bar{\alpha}\beta$, $\alpha, \beta \in \mathbb{C}$.
- ⁴⁰M. S. Birman and M. Z. Solomjak, *Spectral theory of selfadjoint operators in Hilbert space*, Mathematics and its Applications (Soviet Series) (D. Reidel Publishing Co., Dordrecht, 1987) translated from the 1980 Russian original by S. Khrushchëv and V. Peller.
- ⁴¹A detailed account of the reduction theory for bounded operators can be found in Ref. 19.
- ⁴²M. H. Stone, “On unbounded operators in Hilbert space,” *J. Indian Math. Soc. (N.S.)* **15**, 155–192 (1951).
- ⁴³M. J. J. Lennon, “On sums and products of unbounded operators in Hilbert space,” *Trans. Amer. Math. Soc.* **198**, 273–285 (1974).
- ⁴⁴Indeed, let $\mathfrak{H}_1$ and $\mathfrak{H}_2$ be Hilbert spaces, R be a closed operator from $\mathfrak{H}_1$ to $\mathfrak{H}_2$, P be the orthogonal projection of $\mathfrak{H}_1 \oplus \mathfrak{H}_2$ onto G_R , and $\psi_1, \psi_2, \dots$ be elements of $\mathfrak{H}_1$ whose linear span is dense in $\mathfrak{H}_1$. If $(\psi, \psi') \in G_R$ is orthogonal to all vectors $P(\psi_j, 0)$, then $\langle \psi, \psi_j \rangle = \langle (\psi, \psi'), P(\psi_j, 0) \rangle = 0$ for all j and, hence, $\psi = 0$. As G_R is a graph of an operator, it follows that $\psi' = 0$. Thus, the linear span of $P(\psi_j, 0)$ is dense in G_R .
- ⁴⁵M. Reed and B. Simon, *Methods of modern mathematical physics. IV. Analysis of operators* (Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1978).
- ⁴⁶F. Gesztesy, A. Gomilko, F. Sukochev, and Y. Tomilov, “On a question of A. E. Nussbaum on measurability of families of closed linear operators in a Hilbert space,” *Israel J. Math.* **188**, 195–219 (2012).
- ⁴⁷C. Lance, “Direct integrals of left Hilbert algebras,” *Math. Ann.* **216**, 11–28 (1975).
- ⁴⁸P. G. Dixon, “Unbounded operator algebras,” *Proc. London Math. Soc. (3)* **23**, 53–69 (1971).
- ⁴⁹A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher transcendental functions. Vols. I, II* (McGraw-Hill Book Company, Inc., New York-Toronto-London, 1953) based, in part, on notes left by Harry Bateman.